

Low energy effects of neutrino masses

Asmaa Abada,^a Carla Biggio,^b Florian Bonnet,^a Maria B. Gavela^b and Thomas Hambye^{bc}

^a*Laboratoire de Physique Théorique UMR 8627,
 Université de Paris-Sud 11, Bat. 210, 91405 Orsay Cedex, France*

^b*Departamento de Física Teórica and Instituto de Física Teórica IFT-UAM/CSIC,
 Universidad Autónoma de Madrid, 28049 Cantoblanco, Madrid, Spain*

^c*Service de Physique Théorique,
 Université Libre de Bruxelles, 1050 Brussels, Belgium
 E-mail: asmaa.abada@th.u-psud.fr, carla.biggio@uam.es,
 florian.bonnet@th.u-psud.fr, belen.gavela@uam.es, thomas.hambye@uam.es*

ABSTRACT: While all models of Majorana neutrino masses lead to the same dimension five effective operator, which does not conserve lepton number, the dimension six operators induced at low energies conserve lepton number and differ depending on the high energy model of new physics. We derive the low-energy dimension six operators which are characteristic of generic Seesaw models, in which neutrino masses result from the exchange of heavy fields which may be either fermionic singlets, fermionic triplets or scalar triplets. The resulting operators may lead to effects observable in the near future, if the coefficients of the dimension five and six operators are decoupled along a certain pattern, which turns out to be common to all models. The phenomenological consequences are explored as well, including their contributions to $\mu \rightarrow e\gamma$ and new bounds on the Yukawa couplings for each model.

KEYWORDS: Neutrino Physics, Beyond Standard Model, Weak Decays, Rare Decays.

Contents

1. Introduction	2
2. The basic seesaw scenarios	5
2.1 Fermionic singlets: type I seesaw	5
2.1.1 Dimension 5 operator	6
2.1.2 Dimension 6 operator	6
2.2 Scalar triplets: type II seesaw	8
2.2.1 Dimension 4 and 5 operators	9
2.2.2 Dimension 6 operators	10
2.2.3 Renormalization scheme	11
2.3 Fermionic triplets: type III seesaw	12
2.3.1 Dimension 5 operator	12
2.3.2 Dimension 6 operator	13
2.3.3 Parameter counting	15
2.4 Summary	17
3. Low scale seesaw $M \sim \mathcal{O}(TeV)$	17
3.1 Electroweak hierarchy problem	17
3.2 Direct lepton violation	18
4. Phenomenology of seesaw models	21
4.1 Fermionic singlets	21
4.2 Scalar triplets	22
4.2.1 M_W and the ρ parameter	22
4.2.2 $\mu \rightarrow eee$ and $\tau \rightarrow 3l$ decays	24
4.2.3 Complete lagrangian and $\mathbf{l}_1 \rightarrow \mathbf{l}_2 \gamma$	25
4.2.4 Other constraints	26
4.2.5 Collider signatures of scalar triplets	26
4.3 Fermionic triplets	27
4.3.1 W decays	28
4.3.2 Invisible Z decay	28
4.3.3 Universality tests	29
4.3.4 Z decays into charged leptons	29
4.3.5 $\mu \rightarrow eee$ and $\tau \rightarrow 3l$ decays	30
4.3.6 Complete lagrangian and $l_1 \rightarrow l_2 \gamma$	31
4.3.7 Combination of all constraints	33
4.3.8 Signals at colliders from fermionic triplets	34
5. Conclusions	34

A. Non-unitarity	36
B. Lepton mixing in the full type-III seesaw model	38
C. Low scale models of light neutrino masses with large Yukawa couplings	40

1. Introduction

The experimental observation of non-zero neutrino masses and mixings constitutes evidence for physics beyond the Standard Model (SM) and points to the existence of a new, yet unknown, physics scale. It has been already a few years since the breaking of such exciting news and nevertheless little — if anything — is known about the underlying physics. The difficulty lies in both the fact that neutrinos are very weakly interacting particles and, more important, in the tiny value of their masses — orders of magnitude lighter than any other fermion masses — pointing to very suppressed effects. The absence of exotic experimental signals other than neutrino masses, as well as the theoretical criteria of naturalness, point to values of the new physics scale, M , larger than the electroweak scale.

It is worth recalling that the evidence for neutrino masses comes from neutrino oscillations, which detect the interference between the different paths taken by different neutrinos when traveling a long distance. The paths differ because the masses differ and what has been measured is the relative phase shift induced, which is only sizable after extremely long distances. In other words, detection has been possible because neutrino masses affect neutrino propagation. Other possible low-energy effects of the underlying theory, i.e. exotic couplings, are typically zero-distance effects which cannot benefit from such an enhancement. Its suppression is only easily overcome at very high energies, with the particle momenta equal or larger than the scale M , as for instance in leptogenesis scenarios, where the high energies of the early universe allow the heavy fields at the origin of neutrino masses to roam freely.

To see what could be the nature and magnitude of the low energy effects associated to neutrino masses it is convenient to rephrase the above in terms of a generic effective low-energy theory. Effective theories allow rather model-independent analysis based on the fundamental symmetries, while only the coefficient of the effective operators are model-dependent. The impact at low energies of the heavy fields present in the putative high-energy theory can be parametrized, without loss of generality, by an effective Lagrangian including:

- Corrections to the parameters of the SM Lagrangian.
- The addition to the SM Lagrangian of a tower of *non-renormalizable* higher-dimension operators, invariant under the SM gauge group. The latter are made out of the SM fields active at low energies and their coefficients weighted by inverse powers of the high scale M ,

$$\mathcal{L}_{\text{eff}} = \mathcal{L}_{\text{SM}} + \delta\mathcal{L}^{d=5} + \delta\mathcal{L}^{d=6} + \dots \quad (1.1)$$

The only possible dimension 5 ($d = 5$) operator is the famous Weinberg operator [1],

$$\delta\mathcal{L}^{d=5} = \frac{1}{2} c_{\alpha\beta}^{d=5} \left(\overline{\ell}_{L\alpha}^c \tilde{\phi}^* \right) \left(\tilde{\phi}^\dagger \ell_{L\beta} \right) + \text{h.c.}, \quad (1.2)$$

where ℓ_L stands for the lepton weak doublets,¹ greek letters denote flavour indices and $\tilde{\phi}$ is related to the standard Higgs doublet $\phi \equiv (\phi^+, \phi^0)$ by $\tilde{\phi} = i\tau_2 \phi^*$. Finally, $c_{\alpha\beta}^{d=5}$ is a coefficient matrix of inverse mass dimension, i.e. $\mathcal{O}(1/M)$. This operator is not invariant under the $B - L$ symmetry, with B and L denoting respectively baryon and lepton number, which is an accidental symmetry of the SM. Upon electroweak symmetry breaking, $\langle \phi^0 \rangle = v/\sqrt{2}$, $v = 246$ GeV, this term results in Majorana neutrino masses. Such a $d = 5$ operator is characteristic of all theories with Majorana neutrino masses, such as for instance the minimal (type I) Seesaw model [2]. Therefore the knowledge of $c_{\alpha\beta}^{d=5}$ doesn't allow to discriminate between these models. It is very suggestive that the lowest-order effect of high-energy beyond the Standard Model physics may be neutrino masses. There is no hope to see any other low energy effects, e.g. zero distance effects, associated to this operator. These effects are necessarily tiny since neutrino masses — which fix the $c_{\alpha\beta}^{d=5}$ coefficients — are tiny.²

The case of the dimension six ($d = 6$) $SU(3) \times SU(2) \times U(1)$ invariant operators is different, though. There is a plethora of such operators [3]. Different classes of models result in different $d = 6$ operators. Their identification and eventually their experimental selection is then a very important tool to discriminate the origin of neutrino masses. An important property of these operators is that they are not necessarily suppressed as the $d = 5$ operator and, therefore, may lead to observable low-energy effects. The point is that *all of them preserve $B - L$, in contrast with the unique $d = 5$ operator above. This suggests that, from the point of view of symmetries, it may be natural to consider large coefficients for the $d = 6$ operators resulting from the new physics, while having small coefficients for the $B - L$ odd operator.* Such a possibility would require to decouple the coefficients of the $d = 6$ operators from that of the $d = 5$ operator responsible for neutrino masses.

The first purpose of this work is to identify the effective $d = 6$ operators which are characteristic of Seesaw models (section 2). In the latter, the tiny neutrino masses naturally result from the tree-level exchange of heavy particles, which may be either fermions or bosons. The exchange of heavy SM singlet fermions is the essence of the minimal Seesaw model (type I) and its generalizations. Analogously, the exchange of heavy $SU(2)_L$ scalar triplets is another possibility which has been widely explored, as in the type II Seesaw model and its generalizations [4]. $SU(2)_L$ fermionic triplets may also mediate light neutrino masses (type III Seesaw) [5, 6, 6–9]. Most beyond the SM theories with Majorana neutrino masses typically incorporate one of these mechanisms or combinations of them: the lessons

¹The charge-conjugate spinor is denoted $\psi^c \equiv C\overline{\psi}^T$, where T denotes transposition and C charge conjugation.

²Notice that neutrino masses have been detected in neutrino oscillation experiments, which in fact measure differences between the *square* of neutrino masses. That is, if the neutrinos are Majorana particles, the experiments have already measured an effect suppressed as $(c_{\alpha\beta}^{d=5})^2 \sim 1/M^2$ instead of $1/M$ and thus quantitatively alike to that from generic dimension six operators.

learnt from their study should be of extensive relevance. We will thus discuss the effective low-energy Lagrangians for the three generic cases: heavy fermion singlets, heavy scalar triplets and heavy fermionic triplets, illustrated in figure 1.

Next, in a second stage (section 3) we consider the possibility that the $d = 6$ operators are not as suppressed as the $d = 5$ operator, so that observable low-energy effects may be expected. Since these operators are suppressed by $1/M^2$, this requires a value of M not far beyond the electroweak scale. We consider this possibility, which is not excluded at all and may even be supported by hierarchy arguments. It will then be shown that in order to have observable low energy effects, it is necessary and possible to decouple and suppress the coefficient of the $d = 5$ operator relative to the $d = 6$ operator coefficients, in a way which accommodates tiny neutrino masses while allowing large Yukawa couplings. It will be shown that such decoupling requires a common and rather model-independent pattern, which we identify.

In a third stage (section 4) and independently of how large is the scale M , we analyze the long list of phenomenological signals which may arise in each of the three models considered, such as signals associated to non-unitarity or other effects in different observables: neutrino oscillations, lepton and gauge-boson decays. From present data, limits will be set in all models on the coefficients of the $d = 6$ operators. From them, we derive systematic tables of bounds on the Yukawa couplings in each of the Seesaw models. Expectations for the sensitivity of future experiments will be explored, including the contributions to $l_i \rightarrow l_j \gamma$. We show that, in case the decoupling pattern mentioned above occurs, the limits can be saturated if M is still larger than but close to the electroweak scale. The possibilities for direct or indirect discovery of the origin of neutrino masses at the LHC or ILC will be (briefly) discussed.

An important phenomena at the origin of many of the potential low energy effects is non-unitarity of the leptonic mixing matrices. Special emphasis will be set on analyzing whether Seesaw models induce at low energies a non-unitary leptonic mixing matrix. It is expected in all generality [10] that the tree-level exchange of heavy fermions (scalars) will (not) induce it. Indeed, only leptons can mix with other fermions leading to (unitary) mixing matrices of dimension larger than 3, while the submatrix for the light fields needs not be unitary. In a more technical view, the exchange of heavy fermions among light leptons can be understood from the expansion of the heavy field propagator in powers of $1/M$,

$$\frac{1}{\not{p} - M} \sim -\frac{1}{M} + \frac{1}{M} \not{p} \frac{1}{M} + \dots \quad (1.3)$$

The first term in this expansion is a scalar operator, which flips chirality, generating for instance a light neutrino mass term. The second term, instead, preserves chirality and induces a correction to the kinetic term for the light fields. The recovery of canonically normalized kinetic energies for the latter requires in general a flavour-dependent rescaling, which is a non-unitary transformation, surfacing as non-unitary mixing matrices in the leptonic weak currents [10]. Non-unitarity of the leptonic mixing matrix is therefore a basic property of models where masses are induced by heavy fermions. In contrast, in scalar-mediated mechanisms, all terms in the scalar propagator change chirality and thus




Figure 1: The three generic realizations of the Seesaw mechanism, depending on the nature of the heavy fields exchanged: SM singlet fermions (type I Seesaw) on the left, SM triplet scalars (type II Seesaw) and SM triplet fermions (type III Seesaw) on the right.

cannot induce non-unitary mixing at tree-level. The minimal (type I) Seesaw model has been previously shown [11] to induce a non-unitary leptonic mixing matrix. In this work we will explicitly analyze the issue for the other types of Seesaw models.

2. The basic seesaw scenarios

Let us analyze separately the three different minimal models which result from adding either fermionic singlets or scalar triplets or fermionic triplets to the minimal SM field content. It is expected that the lessons obtained from the analysis of the three basic models will hold as well for their possible generalizations, extensions or embeddings in larger theories.

2.1 Fermionic singlets: type I seesaw

As this case has been previously studied [11], only the main results are resumed here for completion. The minimal Seesaw Lagrangian is the most general renormalizable Lagrangian which can be written for the SM gauge group adding only right-handed neutrinos to the SM fermion content of the theory. The leptonic Lagrangian of the Seesaw model is given by

$$\mathcal{L}_{\text{leptons}} = \mathcal{L}_{\text{leptons}}^{\text{KE}} + \mathcal{L}_{\text{leptons}}^{\text{SB}}, \quad (2.1)$$

where

$$\mathcal{L}_{\text{leptons}}^{\text{KE}} = i \bar{\ell}_L \not{D} \ell_L + i \bar{e}_R \not{D} e_R + i \bar{N}_R \not{D} N_R \quad (2.2)$$

contains the kinetic energy and gauge interaction terms of the left-handed lepton doublets ℓ_L , the right-handed charged leptons e_R , the right-handed neutrinos N_R and

$$\mathcal{L}_{\text{leptons}}^{\text{SB}} = -\bar{\ell}_L \phi Y_e e_R - \bar{\ell}_L \tilde{\phi} Y_N^\dagger N_R - \frac{1}{2} \bar{N}_R M_N N_R^c + \text{h.c.} \quad (2.3)$$

contains the Yukawa interactions with coupling Y_N and the Majorana mass term of the gauge-singlet right-handed neutrinos, corresponding to the new physics scale(s) M_N . Flavour indices are implicit in these expressions and we will work in a basis in which M_N is a diagonal complex matrix.

2.1.1 Dimension 5 operator

In the flavour basis, the resulting $d = 5$ operator coefficients are given in terms of the parameters of the high-energy theory as (see figure 1)

$$c^{d=5} = Y_N^T \frac{1}{M_N} Y_N. \quad (2.4)$$

Upon electroweak symmetry breaking, it leads to a Majorana mass matrix for the light neutrinos of the form

$$m_\nu \equiv -\frac{v^2}{2} c^{d=5} = -\frac{1}{2} Y_N^T \frac{v^2}{M_N} Y_N. \quad (2.5)$$

For values of the Yukawa couplings Y_N of order unity, the tiny experimental values of neutrino masses require a scale M_N suggestively close to the Grand Unification scale.

2.1.2 Dimension 6 operator

In ref. [11], the $d = 6$ low-energy effective theory, $\delta\mathcal{L}^{d=6}$, was determined to consist at the tree level of the unique operator

$$\delta\mathcal{L}^{d=6} = c_{\alpha\beta}^{d=6} \left(\overline{\ell_{L\alpha}} \tilde{\phi} \right) i \not{\partial} \left(\tilde{\phi}^\dagger \ell_{L\beta} \right), \quad (2.6)$$

where the $d = 6$ operator coefficients are given in terms of the parameters of the high-energy Seesaw theory by

$$c^{d=6} = Y_N^\dagger \frac{1}{M_N^\dagger} \frac{1}{M_N} Y_N, \quad (2.7)$$

which is of the same order in Yukawa couplings than its $d = 5$ counterpart, eq. (2.4), while quadratically suppressed in $1/M_N$. When the Higgs doublet acquires a vacuum expectation value, this $d = 6$ operator leads to corrections to the $d = 4$ kinetic energy terms for the left-handed Majorana neutrinos, which result in a non-unitary low-energy leptonic mixing matrix [10]. Indeed, the neutrino Lagrangian for the effective theory, including only $d \leq 6$ operators and disregarding couplings to the physical Higgs particle, is given by

$$\mathcal{L}_{\text{neutrino}}^{d \leq 6} = i \overline{\nu_{L\alpha}} \not{\partial} (\delta_{\alpha\beta} + \epsilon_{\alpha\beta}^N) \nu_{L\beta} - \frac{1}{2} \overline{\nu_{L\alpha}}^c m_{\nu\alpha\beta} \nu_{L\beta} - \frac{1}{2} \overline{\nu_{L\alpha}} m_{\nu\alpha\beta}^* \nu_{L\beta}^c, \quad (2.8)$$

where

$$\epsilon^N \equiv \frac{v^2}{2} c^{d=6} \quad (2.9)$$

is the contribution of the $d = 6$ operator coefficient to the left-handed neutrino kinetic energy, which is non-diagonal in flavor space. Let us then go to a basis in which the neutrino field is rescaled, so that the neutrino kinetic energy is canonically normalized: at order $\mathcal{O}(1/M^2)$, the transformation

$$\nu_{L\alpha} \rightarrow \nu'_{L\alpha} \equiv (\delta_{\alpha\beta} + \epsilon_{\alpha\beta}^N)^{\frac{1}{2}} \nu_{L\beta} \quad (2.10)$$

results in a Lagrangian in the flavour basis which, at this order, takes the form (primes will be omitted in the following),

$$\mathcal{L}_{\text{leptons}}^{d \leq 6} = i \overline{\nu_{L\alpha}} \not{\partial} \nu_{L\alpha} + i \overline{l_{L\alpha}} \not{\partial} l_{L\alpha} - \frac{1}{2} [\overline{\nu_{L\alpha}}^c m_{\nu\alpha\beta} \nu_{L\beta} + \text{h.c.}] - \overline{l_{L\alpha}} m_{l\alpha\beta} l_{L\beta} + \mathcal{L}_{CC} + \mathcal{L}_{NC} + \mathcal{L}_{em}, \quad (2.11)$$

where m_l is the charged lepton mass matrix and

$$\begin{aligned}\mathcal{L}_{CC} &= \frac{g}{\sqrt{2}} \bar{l}_{L\alpha} W^- \left(\delta_{\alpha\beta} - \frac{1}{2} \epsilon_{\alpha\beta}^N \right) \nu_{L\beta} + \text{h.c.}, \\ \mathcal{L}_{NC} &= \frac{g}{\cos\theta_W} \left\{ \frac{1}{2} [\bar{\nu}_{L\alpha} \gamma_\mu (\delta_{\alpha\beta} - \epsilon_{\alpha\beta}^N) \nu_{L\beta} - \bar{l}_{L\alpha} \gamma_\mu l_{L\alpha}] - \sin^2\theta_W J_\mu^{em} \right\} Z^\mu, \\ \mathcal{L}_{em} &= e J_\mu^{em} A^\mu,\end{aligned}\tag{2.12}$$

with $J_\mu^{em} = -\bar{l} \gamma_\mu l$ denoting the electromagnetic current. We can now rotate to the basis in which the mass matrices are diagonal,

$$\mathcal{L}_{\text{leptons}}^{d \leq 6} = \frac{1}{2} \bar{\nu}_i (i\not{\partial} - m_{\nu i}^{\text{diag}}) \nu_i + \frac{1}{2} \bar{l}_i (i\not{\partial} - m_{li}^{\text{diag}}) l_i + \mathcal{L}_{CC} + \mathcal{L}_{NC} + \mathcal{L}_{em}.\tag{2.13}$$

Now, because of the flavour-dependent field rescalings involved, the usual U_{PMNS} matrix appearing in the charged-current coupling is replaced by a non-unitary matrix N ,

$$N \equiv \Omega \left(1 - \frac{\epsilon^N}{2} \right) U^\nu,\tag{2.14}$$

where U^ν diagonalizes the neutrino mass matrix and $\Omega \equiv \text{diag}(e^{i\omega_1}, e^{i\omega_2}, e^{i\omega_3})$ reabsorbs three unphysical phases in the definition of the charged lepton fields, as usual. Details of the procedure can be found in appendix A. Notice that, as U^ν does not depend on $c^{d=6}$ at $\mathcal{O}(1/M^2)$, in a flavour basis in which Ω is the identity matrix, N would read

$$N = \left(1 - \frac{\epsilon^N}{2} \right) U_{PMNS}\tag{2.15}$$

and consequently $NN^\dagger = (1 - \epsilon^N)$, $N^\dagger N = U_{PMNS}^\dagger (1 - \epsilon^N) U_{PMNS}$, within the $\mathcal{O}(1/M_N^2)$ considered in this work.

Whatever the flavour basis, in the mass basis the weak currents read now

$$J_\mu^{-CC} \equiv \bar{e}_{L\alpha} \gamma_\mu N_{\alpha i} \nu_i,\tag{2.16}$$

$$J_\mu^{NC} \equiv \frac{1}{2} \bar{\nu}_i \gamma_\mu (N^\dagger N)_{ij} \nu_j,\tag{2.17}$$

where $\sum_\alpha N_{i\alpha}^\dagger N_{\alpha j} \neq \delta_{ij}$ appears in the neutral current since N is not unitary, while the neutral current for charged leptons is the standard one. Accordingly, the Fermi constant measured in experiments, G_F , cannot be identified anymore with the SM tree level combination $G_F^{SM} = \sqrt{2}g^2/(8M_W^2) = \frac{1}{\sqrt{2}v^2}$, due to non-unitarity. For instance, the Fermi constant G_F extracted from the decay $\mu \rightarrow \nu_\mu e \bar{\nu}_e$ is related to G_F^{SM} by [10]

$$G_F = G_F^{SM} \sqrt{(NN^\dagger)_{ee} (NN^\dagger)_{\mu\mu}}.\tag{2.18}$$

The rest of the parameters of the Lagrangian coincide with those in the standard treatment.

It is remarkable that putative departures from unitarity of the leptonic mixing matrix can be now directly related to the $d = 6$ operator coefficients and thus to combinations of

the high-energy parameters,³

$$|NN^\dagger - 1|_{\alpha\beta} = \frac{v^2}{2} |c^{d=6}|_{\alpha\beta} = \frac{v^2}{2} \left| Y_N^\dagger \frac{1}{M_N^\dagger} \frac{1}{M_N} Y_N \right|_{\alpha\beta}. \quad (2.19)$$

In section 4.1 the present numerical constraints on $|c^{d=6}|$ will be explored. For Yukawa couplings $Y_N \sim \mathcal{O}(1)$, the coefficients of the $d = 6$ operator are basically the square of those for the $d = 5$ operator, as eqs. (2.4) and (2.7) show. The smallness of neutrino masses then requires e.g. $M_N \gg v$, which precludes the observation of exotic effects in present and planned facilities for the minimal model discussed in this section. There are, however, situations in which $Y_N \sim \mathcal{O}(1)$ can be accommodated together with $M_N \sim \mathcal{O}(TeV)$ and without fine-tunings, leading to observable effects in the near future, as it will be discussed in section 3.

In ref. [11], it was shown that the low-energy effective theory including only the $d = 5$ and $d = 6$ operators contains an equal (a greater) number of real and imaginary parameters as the high-energy Seesaw model, when the number of right-handed neutrinos in the Seesaw theory is equal to (less than) the number of generations of Standard Model fermions. Thus, the determination of all $d = 5$ and $d = 6$ operator coefficients above would suffice *a priori* to determine all of the parameters of the high-energy Seesaw theory. In consequence, for instance, the leptogenesis rate can be written exclusively in terms of both operator coefficients [11].

Other $d = 6$ operators will be also present in the low-energy Lagrangian, since they are generated by radiative mixing of the above $d = 6$ operator in the renormalization group running between the high-energy and low-energy scales. The effects of these other $d = 6$ operators are in consequence subdominant [11] and will not be further considered. The same statement will hold for all Seesaw theories considered in this work.

2.2 Scalar triplets: type II seesaw

Assume now that the minimal SM matter content is enlarged only by the addition of a SU(2) triplet of scalar fields $\vec{\Delta}$ with hypercharge 2,

$$\vec{\Delta} = (\Delta_1, \Delta_2, \Delta_3), \quad (2.20)$$

whose relation to the physical charge eigenstates,

$$(\Delta^{++}, \Delta^+, \Delta^0), \quad (2.21)$$

is given by

$$\Delta^{++} \equiv \frac{1}{\sqrt{2}}(\Delta^1 - i\Delta^2), \quad \Delta^+ \equiv \Delta^3, \quad \Delta^0 \equiv \frac{1}{\sqrt{2}}(\Delta^1 + i\Delta^2). \quad (2.22)$$

In the minimal Lagrangian, gauge invariance allows a Yukawa coupling of the scalar triplet to two lepton doublets,

$$\mathcal{L}_Y^{\Delta, \nu} = \overline{\ell}_L Y_\Delta (\vec{\tau} \cdot \vec{\Delta}) \ell_L + \text{h.c.}, \quad (2.23)$$

³In the flavour basis above mentioned, in which N is given by eq. (2.15) and the matrix Ω is the identity, the absolute-value bars can be dropped: $(NN^\dagger - 1)_{\alpha\beta} = \frac{v^2}{2} c_{\alpha\beta}^{d=6} = \frac{v^2}{2} (Y_N^\dagger \frac{1}{M_N^\dagger} \frac{1}{M_N} Y_N)_{\alpha\beta}$.

as well as a coupling of the scalar triplet to the Higgs doublet,

$$\mu_\Delta \tilde{\phi}^\dagger (\vec{\tau} \cdot \vec{\Delta})^\dagger \phi + \text{h.c.} \quad (2.24)$$

In these equations τ_i are the Pauli matrices, Y_Δ is a symmetric matrix in generation space and $\tilde{\ell}_L = i\tau_2(\ell_L)^c$ (i.e. $\tilde{\ell}_L = -i\ell_L^T C i\tau_2$). The minimal Lagrangian then writes:

$$\begin{aligned} \mathcal{L}_\Delta = & \left(D_\mu \vec{\Delta} \right)^\dagger \left(D^\mu \vec{\Delta} \right) + \left(\tilde{\ell}_L Y_\Delta (\vec{\tau} \cdot \vec{\Delta}) \ell_L + \mu_\Delta \tilde{\phi}^\dagger (\vec{\tau} \cdot \vec{\Delta})^\dagger \phi + \text{h.c.} \right) \\ & - \left\{ \vec{\Delta}^\dagger M_\Delta^2 \vec{\Delta} + \frac{1}{2} \lambda_2 \left(\vec{\Delta}^\dagger \vec{\Delta} \right)^2 + \lambda_3 \left(\phi^\dagger \phi \right) \left(\vec{\Delta}^\dagger \vec{\Delta} \right) \right. \\ & \left. + \frac{\lambda_4}{2} \left(\vec{\Delta}^\dagger T^i \vec{\Delta} \right)^2 + \lambda_5 \left(\vec{\Delta}^\dagger T^i \vec{\Delta} \right) \phi^\dagger \tau^i \phi \right\}, \end{aligned} \quad (2.25)$$

where summation over the SU(2) indices i is assumed. We choose to work in a basis in which M_Δ is real and diagonal and the covariant derivative D_μ in eq. (2.25) is given by

$$D_\mu \equiv \partial_\mu - ig \vec{T} \vec{W}_\mu - ig' B_\mu \frac{Y}{2}, \quad (2.26)$$

with \vec{T} denoting the dimension-three representations of the SU(2) generators,

$$T_1 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -i \\ 0 & i & 0 \end{pmatrix}, \quad T_2 = \begin{pmatrix} 0 & 0 & i \\ 0 & 0 & 0 \\ -i & 0 & 0 \end{pmatrix}, \quad T_3 = \begin{pmatrix} 0 & -i & 0 \\ i & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}. \quad (2.27)$$

The Lagrangian expressed in terms of the charge components of the $\vec{\Delta}$ field can be found below, in eq. (4.16). Consider the limit in which the triplets are heavy, $M_\Delta \gg v$. To solve the equation of motion for Δ^α in eq. (2.25) and find the dominant terms of the effective low-energy Lagrangian up to $d = 6$ operators, it suffices to solve the problem perturbatively in the quartic couplings of $\vec{\Delta}$, λ_2 and λ_4 . At zero order, it results:

$$\Delta^\alpha = \left[(D_\mu)^2 + \lambda_5 \vec{T} \phi^\dagger \vec{\tau} \phi + \left(M_\Delta^2 + \lambda_3 \left(\phi^\dagger \phi \right) \right) \cdot \mathbf{1}_{\text{isospin}} \right]_{\alpha\beta}^{-1} \left[\mu_\Delta^* \tilde{\phi}^\dagger \tau^\beta \phi + \tilde{\ell}_L Y_\Delta^\dagger \tau^\beta \tilde{\ell}_L \right]. \quad (2.28)$$

2.2.1 Dimension 4 and 5 operators

Expanding now the effective Lagrangian — using eq. (2.28) — in inverse powers of M_Δ , one dimension four operator emerges:

$$\delta \mathcal{L}^{d=4} = \frac{|\mu_\Delta|^2}{M_\Delta^2} \left(\tilde{\phi}^\dagger \vec{\tau} \phi \right) \left(\phi^\dagger \vec{\tau} \tilde{\phi} \right) = 2 \frac{|\mu_\Delta|^2}{M_\Delta^2} \left(\phi^\dagger \phi \right)^2. \quad (2.29)$$

We also obtain $\delta \mathcal{L}^{d=5}$ as given in eq. (1.2), with operator coefficients given by the matrix

$$c^{d=5} = 4Y_\Delta \frac{\mu_\Delta}{M_\Delta^2}, \quad (2.30)$$

which at low energies leads to a light neutrino Majorana mass matrix of the form

$$m_\nu = -2Y_\Delta v^2 \frac{\mu_\Delta}{M_\Delta^2}. \quad (2.31)$$

Notice that neutrino masses turn out to be proportional to both Y_Δ and μ_Δ , see figure 1. This is as expected from the Lagrangian, eq. (2.25), where the breaking of lepton number symmetry L results precisely from the simultaneous presence of the Yukawa and μ_Δ couplings.⁴ It is important that, unlike for the fermionic Seesaw theories, the light neutrino mass matrix in eq. (2.31) is only *linearly* dependent on the Yukawa coupling Y_Δ . This means that the putative determination of the $d = 5$ operator coefficients gives a direct access to the fundamental parameters Y_Δ of the high-energy theory, up to an overall scale μ_Δ/M_Δ^2 . We will analyze in section 4 the experimental access to μ_Δ/M_Δ^2 and to the elements of Y_Δ .

2.2.2 Dimension 6 operators

From eq. (2.28), the $d = 6$ effective Lagrangian can also be obtained,

$$\delta\mathcal{L}_\Delta^{d=6} = \delta\mathcal{L}_{4F} + \delta\mathcal{L}_{\phi D} + \delta\mathcal{L}_{6\phi}, \quad (2.32)$$

where

$$\begin{cases} \delta\mathcal{L}_{4F} = \frac{1}{M_\Delta^2} \left(\widetilde{\ell}_L Y_\Delta \vec{\tau} \ell_L \right) \left(\overline{\ell}_L \vec{\tau} Y_\Delta^\dagger \widetilde{\ell}_L \right) \\ \delta\mathcal{L}_{6\phi} = -2(\lambda_3 + \lambda_5) \frac{|\mu_\Delta|^2}{M_\Delta^4} (\phi^\dagger \phi)^3 \\ \delta\mathcal{L}_{\phi D} = \frac{|\mu_\Delta|^2}{M_\Delta^4} (\phi^\dagger \vec{\tau} \widetilde{\phi}) \left(\overline{D}_\mu \overline{D}^\mu \right) (\widetilde{\phi}^\dagger \vec{\tau} \phi) \end{cases}, \quad (2.33)$$

with the covariant derivative expressed in terms of (3×3) $SU(2)$ generators, as in eq. (2.26).⁵ Two of these operators can be rewritten in a more familiar form. After Fierz transformation, $\delta\mathcal{L}_{4F}$ can be expressed as

$$\delta\mathcal{L}_{4F} = -\frac{1}{M_\Delta^2} Y_{\Delta ij} Y_{\Delta \alpha\beta}^\dagger (\overline{\ell}_{L\beta} \gamma_\mu \ell_{Li}) (\overline{\ell}_{L\alpha} \gamma_\mu \ell_{Lj}), \quad (2.34)$$

while the last operator in eq. (2.33) can be recast as a combination of other operators which have been extensively studied in the literature (e.g. [3]),

$$\delta\mathcal{L}_{\phi D} = 4 \frac{|\mu_\Delta|^2}{M_\Delta^4} (\phi^\dagger \phi) \left[(D_\mu \phi)^\dagger (D_\mu \phi) \right] + 4 \frac{|\mu_\Delta|^2}{M_\Delta^4} \left[\phi^\dagger D_\mu \phi \right]^\dagger \left[\phi^\dagger D_\mu \phi \right], \quad (2.35)$$

where the covariant derivative is meant to be expressed in terms of Pauli matrices,

$$D_\mu = \partial_\mu - ig \frac{\tau_a}{2} W_{a\mu} - ig' B_\mu \frac{Y}{2}. \quad (2.36)$$

⁴In the language of the full theory, this mass results when the neutral component of Δ acquires a vev $\langle \Delta^0 \rangle \equiv u/\sqrt{2} = \mu_\Delta v^2/(\sqrt{2}M_\Delta^2)$, leading to a Majorana mass matrix for the SM neutrinos, $m_\nu = -2Y_\Delta u$.

⁵The first of these operators has already been derived in [12].

2.2.3 Renormalization scheme

Four parameters of the SM are relevant to our discussions (in addition to fermion masses): g, g', v and λ , the latter denoting the quartic self-coupling of the Higgs field,

$$V = -\mu_\phi^2 |\phi|^2 + \lambda |\phi|^4. \quad (2.37)$$

To constrain the first three parameters, we will work in the Z-scheme [13], that is, we will use as input parameters the very-well determined experimental values of the fine structure constant α — as determined from Thompson scattering⁶ — the Fermi constant G_F — as extracted from the muon decay rate by the removal of SM process-dependent radiative corrections — and the very precise measurement of M_Z [14].

The value of α is not affected by the presence of a scalar triplet, unlike the other parameters. M_Z gets a correction from $\delta\mathcal{L}_{\phi D}$ in eq. (2.33)

$$\frac{\delta M_Z^2}{M_Z^2} = 2v^2 \frac{|\mu_\Delta|^2}{M_\Delta^4}. \quad (2.38)$$

Similarly, the 4-fermion operator $\delta\mathcal{L}_{4F}$ affects the extraction of the value of the Fermi constant from muon decay. Defining, as it is customary, this constant as the coefficient in

$$-\frac{4G_F}{\sqrt{2}} \left(\bar{\ell}_{L\nu\beta} \gamma_\mu \ell_{L\mu} \right) \left(\bar{\ell}_{Le} \gamma_\mu \ell_{L\nu\alpha} \right), \quad (2.39)$$

it is easily seen that $\delta\mathcal{L}_{4F}$ in eq. (2.34) induces in turn a shift with respect to the “Standard Model definition” $G_F^{SM} = 1/(\sqrt{2}v^2)$,⁷ which affects the value extracted from muon decay,

$$\delta G_F = \frac{1}{\sqrt{2}M_\Delta^2} |Y_{\Delta e\mu}|^2 \quad (2.40)$$

$$G_F = G_F^{SM} + \delta G_F. \quad (2.41)$$

The quartic self-coupling of the Higgs field is also renormalized by the dimension four operator obtained in the effective theory, eq. (2.29),

$$\delta\lambda = -2 \frac{|\mu_\Delta|^2}{M_\Delta^2}, \quad (2.42)$$

influencing the location of the minimum of the Higgs potential. Another $d = 6$ operators, $\delta\mathcal{L}_{6\phi}$ in eq. (2.33), also modifies the Higgs potential, which all in all becomes

$$V = -\mu_\phi^2 |\phi|^2 + (\lambda + \delta\lambda) |\phi|^4 + 2(\lambda_3 + \lambda_5) \frac{|\mu_\Delta|^2}{M_\Delta^4} |\phi|^6, \quad (2.43)$$

inducing a shift in the vacuum expectation value of the Higgs field,

$$\frac{\delta v^2}{v^2} = -3v^2 \frac{|\mu_\Delta|^2}{M_\Delta^4} \frac{(\lambda_3 + \lambda_5)}{\lambda + \delta\lambda}. \quad (2.44)$$

⁶An even more precise determination is now available from $g - 2$ of the electron [14].

⁷Note that with a scalar triplet $\frac{1}{2v^2} \neq \frac{g^2}{8M_W^2}$ due to the scalar triplet induced M_W shift, see below.

Using all these renormalized parameters, in section 4.2 we will consider the deviations — with respect to the SM predictions — induced by the new physics on the values taken by a variety of physical observables.

Finally, as regards the relative number of parameters in the high and low energy theories, the inclusion in the latter of only the $d = 5$ and $d = 6$ operators above does not suffice to match the number of free parameters of the full scalar-triplet Seesaw theory, as can be easily deduced from the comparison of eq. (2.25) with eqs. (2.32) and (2.33). Up to $d = 8$ operators would have to be considered for this purpose, which is beyond the scope of the present work.

2.3 Fermionic triplets: type III seesaw

Consider now the SM field content extended by the only addition of fermions which are triplets of $SU(2)$ with zero hypercharge, hereafter denoted by $\vec{\Sigma}$, where the vectorial character refers to its three $SU(2)$ -components, $\vec{\Sigma} = (\Sigma^1, \Sigma^2, \Sigma^3)$. Being $\vec{\Sigma}$ in the adjoint representation of the gauge group, its Majorana mass term is gauge invariant and the interactions are described by the Lagrangian

$$\mathcal{L}_\Sigma = i \vec{\Sigma}_R^\dagger \not{D} \vec{\Sigma}_R - \left[\frac{1}{2} \vec{\Sigma}_R^\dagger M_\Sigma \vec{\Sigma}_R + \vec{\Sigma}_R^\dagger Y_\Sigma (\tilde{\phi}^\dagger \tilde{\tau} \ell_L) + \text{h.c.} \right]. \quad (2.45)$$

In this equation, the covariant derivative is given by eqs. (2.26) and (2.27) and the three $SU(2)$ -components of the field $\vec{\Sigma}$ have (identical) Majorana mass terms. They are not eigenstates of the electric charge, which would be given instead by the combinations

$$\Sigma^\pm \equiv \frac{\Sigma^1 \mp i \Sigma^2}{\sqrt{2}}, \quad \Sigma^0 \equiv \Sigma^3. \quad (2.46)$$

We will work throughout in a basis in which M_Σ is a diagonal matrix in generation space. The Yukawa coupling Y_Σ in eq. (2.45) is then a general matrix in generation space. After electroweak symmetry breaking, this term induces Majorana neutrino masses for the left-handed neutrino fields of the SM through the exchange of $\vec{\Sigma}$ particles, see figure 1.

2.3.1 Dimension 5 operator

Solving the equations of motion, it results that

$$\begin{aligned} \vec{\Sigma}_R &= P_R [i \not{D} - M_\Sigma]^{-1} \left[Y_\Sigma^* \phi^\dagger \tilde{\tau} \tilde{\ell}_L + Y_\Sigma \tilde{\phi}^\dagger \tilde{\tau} \ell_L \right] \\ &= -\frac{1}{M_\Sigma} Y_\Sigma^* \phi^\dagger \tilde{\tau} \tilde{\ell}_L - \frac{1}{M_\Sigma^\dagger} i \not{D} \frac{1}{M_\Sigma} Y_\Sigma \tilde{\phi}^\dagger \tilde{\tau} \ell_L + \mathcal{O} \left(\frac{1}{M_\Sigma^3} \right), \end{aligned} \quad (2.47)$$

where i, j are $SU(2)$ indices, $i, j = 1, 2, 3$. This allows to obtain the $d = 5$ operator in eq. (1.2), with coefficient matrix given in this case by

$$c^{d=5} = Y_\Sigma^T \frac{1}{M_\Sigma} Y_\Sigma, \quad (2.48)$$

which leads at low energies to a light neutrino Majorana mass matrix of the form

$$m_\nu = -\frac{v^2}{2} Y_\Sigma^T \frac{1}{M_\Sigma} Y_\Sigma. \quad (2.49)$$

2.3.2 Dimension 6 operator

At the next order in the effective Lagrangian, we obtain a unique operator:⁸

$$\delta\mathcal{L}^{d=6} = c_{\alpha\beta}^{d=6} \left(\overline{\ell}_{L\alpha} \vec{\tau} \tilde{\phi} \right) i \not{D} \left(\tilde{\phi}^\dagger \vec{\tau} \ell_{L\beta} \right), \quad (2.50)$$

where the $d = 6$ operator coefficients are given in terms of the parameters of the high-energy Seesaw theory by

$$c^{d=6} = Y_\Sigma^\dagger \frac{1}{M_\Sigma^\dagger} \frac{1}{M_\Sigma} Y_\Sigma. \quad (2.51)$$

Notice the large parallelism between the results for this Seesaw scenario mediated by fermionic triplets and those for the minimal Seesaw based on the exchange of fermionic singlets, eqs. (2.4) and (2.6)–(2.7). The main difference is that, now, in the $d = 6$ operator in eq. (2.50) the interaction terms in the covariant derivative are active, as the quantities in brackets are SU(2) triplets, which amounts to a richer interaction pattern.

A first consequence is that, when the Higgs doublet acquires a vacuum expectation value, the $d = 6$ operator corrects both the $d = 4$ kinetic energy terms of light leptons *and* their couplings to W bosons, while no corrections to the hypercharge boson B_μ appeared, because the combinations in brackets in eq. (2.50) have zero hypercharge. After electroweak symmetry breaking, the part of the effective Lagrangian concerning leptons is, in the flavour basis,

$$\begin{aligned} \mathcal{L}_{\text{leptons}}^{d \leq 6} = & i \overline{\nu}_{L\alpha} \not{\partial} (\delta_{\alpha\beta} + \epsilon_{\alpha\beta}^\Sigma) \nu_{L\beta} + i \overline{l}_{L\alpha} \not{\partial} (\delta_{\alpha\beta} + 2\epsilon_{\alpha\beta}^\Sigma) l_{L\beta} + i \overline{l}_{R\alpha} \not{\partial} l_{R\alpha} \\ & - \frac{1}{2} [\overline{\nu}_{L\alpha}^c m_{\nu\alpha\beta} \nu_{L\beta} + \text{h.c.}] - [\overline{l}_{R\alpha} m_{l\alpha\beta} l_{L\beta} + \text{h.c.}] \\ & + \frac{1}{\sqrt{2}} g [\overline{l}_{L\alpha} W^- (\delta_{\alpha\beta} + 2\epsilon_{\alpha\beta}^\Sigma) \nu_{L\beta} + \text{h.c.}] - \frac{g}{2} \overline{l}_{L\alpha} W^3 (\delta_{\alpha\beta} + 4\epsilon_{\alpha\beta}^\Sigma) l_{L\beta} \\ & + \frac{g}{2} \overline{\nu}_{L\alpha} W^3 \nu_{L\alpha} - \frac{g'}{2} \overline{l}_{L\alpha} \not{B} l_{L\beta} - \frac{g'}{2} \overline{\nu}_{L\alpha} \not{B} \nu_{L\alpha}, \end{aligned} \quad (2.52)$$

where

$$\epsilon^\Sigma \equiv \frac{v^2}{2} c^{d=6}, \quad (2.53)$$

with $c^{d=6}$ as defined in eq. (2.51) and m_l denoting the charged lepton mass matrix. We assume hereafter a choice of basis in which both m_l and M_Σ are diagonal. The neutrino *and* charged lepton fields need now to be normalized in order to acquire canonically normalized kinetic terms. At order $1/M^2$, i.e. linear in the parameters $\epsilon_{\alpha\beta}^\Sigma$, the redefinitions

$$\begin{aligned} \nu_{L\alpha} &\rightarrow \nu'_{L\alpha} \equiv \left(\delta_{\alpha\beta} + \frac{1}{2} \epsilon_{\alpha\beta}^\Sigma \right) \nu_{L\beta}, \\ l_{L\alpha} &\rightarrow l'_{L\alpha} \equiv (\delta_{\alpha\beta} + \epsilon_{\alpha\beta}^\Sigma) l_{L\beta}, \end{aligned} \quad (2.54)$$

⁸We thank S. Antusch for helping to clarify the derivation of this operator in an early stage.

results in a Lagrangian in the flavour basis which, at order $\mathcal{O}(1/M^2)$, takes the form (primes on the fields will be disregarded),

$$\begin{aligned} \mathcal{L}_{\text{leptons}}^{d \leq 6} = & i\overline{\nu}_{L\alpha}\not{\partial}\nu_{L\alpha} + i\overline{l}_{L\alpha}\not{\partial}l_{L\alpha} + i\overline{l}_{R\alpha}\not{\partial}l_{R\alpha} - \frac{1}{2} \left[\overline{\nu}_{L\alpha}^c m'_{\nu\alpha\beta} \nu_{L\beta} + \text{h.c.} \right] \\ & - \left[\overline{l}_{R\alpha} m'_{l\alpha\beta} l_{L\beta} + \text{h.c.} \right] + \mathcal{L}_{CC} + \mathcal{L}_{NC} + \mathcal{L}_{em}, \end{aligned} \quad (2.55)$$

where $m'_\nu \equiv (1 - \epsilon^*/2)m_\nu(1 - \epsilon/2)$, $m'_l \equiv m_l(1 - \epsilon)$ and

$$\mathcal{L}_{CC} = \frac{g}{\sqrt{2}} \overline{l}_{L\alpha} W^- \left(\delta_{\alpha\beta} + \frac{1}{2} \epsilon_{\alpha\beta}^\Sigma \right) \nu_{L\beta} + \text{h.c.}, \quad (2.56)$$

$$\begin{aligned} \mathcal{L}_{NC} = & \frac{g}{\cos\theta_W} \left\{ \frac{1}{2} \left[\overline{\nu}_{L\alpha} \gamma_\mu (\delta_{\alpha\beta} - \epsilon_{\alpha\beta}^\Sigma) \nu_{L\beta} - \overline{l}_{L\alpha} \gamma_\mu (\delta_{\alpha\beta} + 2\epsilon_{\alpha\beta}^\Sigma) l_{L\beta} \right] - \sin^2\theta_W J_\mu^{em} \right\} Z^\mu, \\ \mathcal{L}_{em} = & e J_\mu^{em} A^\mu, \end{aligned}$$

where $J_\mu^{em} = -\overline{l}\gamma_\mu l$ is the electromagnetic current. We can finally rotate to the basis in which both the lepton kinetic energies and their mass matrices are diagonalized (for details see appendix A),

$$\mathcal{L}_{\text{leptons}}^{d \leq 6} = \frac{1}{2} \overline{\nu}_i \left(i\not{\partial} - m_{\nu i}^{\text{diag}} \right) \nu_i + \frac{1}{2} \overline{l}_i \left(i\not{\partial} - m_{l i}^{\text{diag}} \right) l_i + \mathcal{L}_{CC} + \mathcal{L}_{NC} + \mathcal{L}_{em}. \quad (2.57)$$

A non-unitary mixing matrix N replaces now the usual unitary U_{PMNS} matrix in the charged current couplings contained in eq. (2.57), because of the flavour-dependent field rescaling involved, while the couplings to the Z boson acquire also a non-unitary mixing pattern,

$$J_\mu^{CC} \equiv \overline{l}_L \gamma_\mu N \nu, \quad (2.58)$$

$$J_\mu^3(\text{neutrinos}) \equiv \frac{1}{2} \overline{\nu} \gamma_\mu (N^\dagger N)^{-1} \nu, \quad (2.59)$$

$$J_\mu^3(\text{leptons}) \equiv \frac{1}{2} \overline{l} \gamma_\mu (N N^\dagger)^2 l. \quad (2.60)$$

The non-unitary mixing matrix N is a function of the $d = 6$ coefficient matrix

$$N \equiv \Omega U_L^{l\dagger} \left(1 + \frac{1}{2} \epsilon^\Sigma \right) U^\nu, \quad (2.61)$$

where, once again, $\Omega \equiv \text{diag}(e^{i\omega_1}, e^{i\omega_2}, e^{i\omega_3})$ reabsorbs three unphysical phases in the definition of the charged lepton fields, and the matrices U^ν and U_L^l diagonalize the effective leptonic mass matrices,⁹ $m_\nu^{\text{diag}} \equiv U^{\nu T} m_\nu U^\nu$, $m_l^{\text{diag}} \equiv U_R^{l\dagger} m_l (1 - \epsilon) U_L^l$ (see appendix A). When the flavour basis chosen is such that both U_L^l and Ω are equal to the identity matrix, and taking into account that U^ν does not receive corrections from $c^{d=6}$ at $\mathcal{O}(1/M_\Sigma^2)$, N simplifies to

$$N \equiv \left(1 + \frac{1}{2} \epsilon^\Sigma \right) U_{PMNS} \quad (2.62)$$

⁹Within the order ϵ^Σ used throughout, the mass eigenvalues are defined at first order in it and thus the eigenvectors should be consistently defined at order zero in that expansion. As a consequence, any representation of the leptonic matrices $U_{L,R}^l$ which diagonalizes the mass matrix has to be physically equivalent to the identity.

and, consequently, $NN^\dagger = 1 + \epsilon^\Sigma$, $N^\dagger N = U_{PMNS}^\dagger(1 + \epsilon^\Sigma)U_{PMNS}$. These expressions can be compared with the equivalent ones for the singlet-fermion Seesaw theory, eq. (2.15) and below it.

Whatever the basis, the currents in eqs. (2.58)–(2.60) can also be compared with the corresponding ones for the singlet-fermion Seesaw theory, eq. (2.16) and (2.17). *A non-unitary mixing pattern has appeared in both cases, although the modified Z-neutrino couplings differ and non-unitary flavour mixing is now also present in the Z-charged lepton couplings.*

An important consequence of the flavour-changing W and Z-lepton couplings is their contribution to muon decay into electron plus missing energy, which modifies the definition of G_F as extracted from muon decay, as follows:

$$G_F = G_F^{SM} \sqrt{(NN^\dagger)_{ee}(NN^\dagger)_{\mu\mu} + \frac{3}{4}[(NN^\dagger)_{e\mu}^2]^2} \sim G_F^{SM} \sqrt{(NN^\dagger)_{ee}(NN^\dagger)_{\mu\mu}}, \quad (2.63)$$

where higher order correction, $\mathcal{O}((\epsilon^\Sigma)^2)$, have been neglected in the last step. Its phenomenological consequences will be explored in section (4.3).

Finally, in analogy with the case of the fermionic singlet Seesaw theory, it is remarkable that departures from unitarity of the leptonic mixing matrix can be now directly related to the $d = 6$ operator coefficients and thus to combinations of the high-energy parameters,

$$|NN^\dagger - 1|_{\alpha\beta} = |\epsilon^\Sigma| = \frac{v^2}{2} |c^{d=6}|_{\alpha\beta} = \frac{v^2}{2} \left| Y_\Sigma^\dagger \frac{1}{M_\Sigma^\dagger} \frac{1}{M_\Sigma} Y_\Sigma \right|_{\alpha\beta}. \quad (2.64)$$

Once again, in the flavour basis in which Ω and U_L^l equal the identity matrix, the absolute-value bars in this equation can be dropped. In section 4.3 the present numerical constraints on $|c^{d=6}|$ will be explored.

2.3.3 Parameter counting

Finally, it can be shown that the low-energy effective theory, including only the $d = 5$ and $d = 6$ operators, contains in this case an equal (greater) number of real and imaginary parameters as the high-energy Seesaw model, when the number of right-handed fermionic triplet generations in the Seesaw theory is equal to (less than) the number of generations of Standard Model fermions. The demonstration is equivalent to that in ref. [11] for the case of singlet-fermion Seesaw theory. The kinetic energy terms in the Lagrangian, eq. (2.45), are invariant under the chiral transformations

$$\begin{aligned} \ell_L &\rightarrow V_\ell \ell_L, \\ e_R &\rightarrow V_e e_R, \\ \Sigma_R &\rightarrow V_\Sigma \Sigma_R, \end{aligned} \quad (2.65)$$

where the V 's are unitary transformations. Consider first the complete theory with n lepton families and n' right-handed fermionic triplets. The Yukawa terms and the Majorana mass term are not invariant under such chiral symmetry, but invariance can be recovered if they

Seesaw Model

Matrix	Moduli	Phases
Y_e	$n \times n$	$n \times n$
Y_Σ	$n \times n'$	$n \times n'$
M_Σ	$\frac{n'(n'+1)}{2}$	$\frac{n'(n'+1)}{2}$
V_e	$\frac{n(n-1)}{2}$	$\frac{n(n+1)}{2}$
V_ℓ	$\frac{n(n-1)}{2}$	$\frac{n(n+1)}{2}$
V_Σ	$\frac{n'(n'-1)}{2}$	$\frac{n'(n'+1)}{2}$
N_{phys}	$n + n' + nn'$	$n(n' - 1)$

Table 1: Number of physical parameters, for n light and n' heavy neutrino generations.

are considered as spurion fields transforming as

$$\begin{aligned}
 Y_e &\rightarrow Y'_e \equiv V_\ell Y_e V_e^\dagger, \\
 Y_\Sigma &\rightarrow Y'_\Sigma \equiv V_\Sigma Y_\Sigma V_\ell^\dagger, \\
 M_\Sigma &\rightarrow M'_\Sigma \equiv V_\Sigma M_\Sigma V_\Sigma^t.
 \end{aligned}
 \tag{2.66}$$

Counting how many physical parameters N_{phys} are needed to describe the Yukawa and Majorana mass terms in the Seesaw Lagrangian is tantamount to counting how many equivalence classes there exist with respect to these transformations. The result is given by

$$N_{\text{phys}} = N_{\text{order}} - (N_G - N_H), \tag{2.67}$$

where N_{order} is the total number of parameters contained in the Yukawa and Majorana mass matrices, N_G is the number of parameters contained in the matrices of the chiral symmetry group $G = \text{U}(n)_\ell \times \text{U}(n)_e \times \text{U}(n')_N$. N_H is the number of parameters contained in the matrices of the subgroup H of the chiral symmetry group which remains unbroken by the Yukawa and Majorana mass matrices: in the present model there is no unbroken subgroup H because of lepton number violation. table 1 summarizes the result for the high-energy theory. This is to be compared with the effective low-energy Lagrangian including operators of $d \leq 6$. It is invariant under the chiral transformations only if

$$\begin{aligned}
 c^{d=5} &\rightarrow V_\ell^* c^{d=5} V_\ell^\dagger, \\
 c^{d=6} &\rightarrow V_\ell c^{d=6} V_\ell^\dagger.
 \end{aligned}
 \tag{2.68}$$

$c^{d=5}$ is a complex symmetric matrix and $c^{d=6}$ is a complex hermitian matrix and since the dimension 5 operators breaks lepton number, there is no unbroken subgroup that remains. The corresponding counting of parameters is shown in table 2, to be compared with that in table 1 for the high-energy theory.

Thus, the determination of all $d = 5$ and $d = 6$ operator coefficients above would again suffice *a priori* to determine all of the parameters of the high-energy Seesaw theory with

Effective Theory ($d \leq 6$)

Matrix	Moduli	Phases
Y_e	$n \times n$	$n \times n$
$c^{d=5}$	$\frac{n(n+1)}{2}$	$\frac{n(n+1)}{2}$
$c^{d=6}$	$\frac{n(n+1)}{2}$	$\frac{n(n-1)}{2}$
V_e	$\frac{n(n-1)}{2}$	$\frac{n(n+1)}{2}$
V_ℓ	$\frac{n(n-1)}{2}$	$\frac{n(n+1)}{2}$
N_{phys}	$n(n+2)$	$n(n-1)$

Table 2: Number of physical parameters, for n light lepton generations.

Model	Effective Lagrangian $\mathcal{L}_{\text{eff}} = c_i \mathcal{O}_i$		
	$c^{d=5}$	$c_i^{d=6}$	$\mathcal{O}_i^{d=6}$
Fermionic Singlet	$Y_N^T \frac{1}{M_N} Y_N$	$\left(Y_N^\dagger \frac{1}{M_N^\dagger} \frac{1}{M_N} Y_N \right)_{\alpha\beta}$	$\left(\overline{\ell_{L\alpha}} \tilde{\phi} \right) i \not{\partial} \left(\tilde{\phi}^\dagger \ell_{L\beta} \right)$
Scalar Triplet	$4Y_\Delta \frac{\mu_\Delta}{M_\Delta^2}$	$\frac{1}{M_\Delta^2} Y_{\Delta\alpha\beta} Y_{\Delta\gamma\delta}^\dagger$	$\left(\overline{\ell_{L\alpha}} \vec{\tau} \ell_{L\beta} \right) \left(\overline{\ell_{L\gamma}} \vec{\tau} \ell_{L\delta} \right)$
		$\frac{ \mu_\Delta ^2}{M_\Delta^4}$	$\left(\phi^\dagger \vec{\tau} \tilde{\phi} \right) \left(\overline{D_\mu} D^\mu \right) \left(\tilde{\phi}^\dagger \vec{\tau} \phi \right)$
		$-2(\lambda_3 + \lambda_5) \frac{ \mu_\Delta ^2}{M_\Delta^4}$	$(\phi^\dagger \phi)^3$
Fermionic Triplet	$Y_\Sigma^T \frac{1}{M_\Sigma} Y_\Sigma$	$\left(Y_\Sigma^\dagger \frac{1}{M_\Sigma^\dagger} \frac{1}{M_\Sigma} Y_\Sigma \right)_{\alpha\beta}$	$\left(\overline{\ell_{L\alpha}} \vec{\tau} \tilde{\phi} \right) i \not{\partial} \left(\tilde{\phi}^\dagger \vec{\tau} \ell_{L\beta} \right)$

Table 3: Coefficients of the $d = 5$ operator, $c^{d=5}$, and $d = 6$ operators and their coefficients, $c^{d=6}$, in the three basic Seesaw theories.

two or three heavy neutrino generations. In consequence, for instance, the leptogenesis rate could be written exclusively in terms of both operator coefficients [15].

2.4 Summary

To conclude this section, we have gathered in table 3 the $d = 6$ operators obtained for the three basic Seesaw scenarios, together with the corresponding expressions for the elements $d = 6$ coefficient matrices. The elements of $d = 5$ coefficient matrices are included as well.

3. Low scale seesaw $M \sim \mathcal{O}(TeV)$

3.1 Electroweak hierarchy problem

If the Seesaw scale is far above the electroweak scale, the theory clashes with the electroweak

hierarchy problem, that is, the fact that data indicate a value for the Higgs mass of the order of the electroweak scale, $v \sim \mathcal{O}(100)\text{GeV}$. Such a mass is unnaturally light if there is new physics beyond the SM and at a higher scale, to which the Higgs boson is sensitive. The three minimal scenarios considered in the previous section do face this problem if the new scales are much larger than the electroweak scale v .

Indeed, for the Seesaw Type I, the one-loop contribution to the Higgs mass has been computed long ago [16],

$$\delta m_H^2 = -\frac{Y_N^\dagger Y_N}{16\pi^2} \left[2\Lambda^2 + 2M_N^2 \log \frac{M_N^2}{\Lambda^2} \right], \quad (3.1)$$

while in the case of the scalar-triplet (type II), we find that the contribution is given by¹⁰

$$\delta m_H^2 = \frac{1}{16\pi^2} \left[3\lambda_3 \left(\Lambda^2 - M_\Delta^2 \log \frac{\Lambda^2}{M_\Delta^2} \right) - 12|\mu_\Delta|^2 \log \frac{\Lambda^2}{M_\Delta^2} \right], \quad (3.2)$$

and, finally, for the fermionic-triplet Seesaw (type III), we obtain

$$\delta m_H^2 = -3 \frac{Y_\Sigma^\dagger Y_\Sigma}{16\pi^2} \left[2\Lambda^2 + 2M_\Sigma^2 \log \frac{M_\Sigma^2}{\Lambda^2} \right], \quad (3.3)$$

where Λ is the regulator cutoff. In these equations, terms proportional to v^2 and m_H^2 have been neglected. eqs. (3.1)–(3.3) all show a quadratic sensitivity to the new scales characteristic of Seesaw theories, implying that large fine-tunings would be necessary to accommodate the experimental data if *any* of the new scales introduced is much larger than v (or the Yukawa couplings are not extremely fine-tuned in Type I and III Seesaw).

For instance, imposing that the one-loop correction is not larger than the Higgs mass itself, let's say $m_H = 150\text{ GeV}$ for definiteness, M_N and M_Σ should be below $\sim 10^7\text{ GeV}$ for Yukawa couplings of order $m_\nu^{1/2} M_{N,\Sigma}^{1/2}/v$, while M_Δ should be below a scale which depends on λ_3 and μ_Δ . In any case, for scales not much larger than the electroweak one, the contribution of the Seesaw theory to the hierarchy problem would be obviously avoided. As a by-product, new exciting physics signals would then be expected at present and future experimental facilities.

The question we wish to analyze now is whether it is indeed possible that nature has chosen the high energy scale M of the Seesaw scenario close to the electroweak scale, rather than to the Grand Unified scale, with $\mathcal{O}(1)$ Yukawa couplings, without fine-tuning the parameters and in particular the Yukawa couplings.

3.2 Direct lepton violation

After all, the analysis of the previous sections has shown that, while neutrino masses result from a lepton-number odd $d = 5$ operator, other manifestations of the new physics behind are encoded in lepton-number conserving $d = 6$ operators (as well as in higher dimensional operators). As lepton number appears to be an approximate symmetry of

¹⁰No dependence on the quartic coupling λ_5 of the Lagrangian eq. (2.25) appears, as the Higgs fields are combined in this term in a triplet of $\text{SU}(2)$, while the Higgs mass is a singlet.

nature, it is natural to assume that it may be broken through small parameters — such as those responsible for neutrino masses — while other beyond the SM effects of the high-energy theory, which are lepton-number preserving, need not be strongly suppressed. The choice of such a L-odd small parameter may be thus a natural one, as it corresponds to the breaking of a symmetry and its value cannot be destabilized by other large scales of the theory through radiative corrections, because by nature it can only be multiplicatively renormalized.

Assume thus M (M_N , M_Δ , M_Σ) to be higher but not far from the TeV scale. The issue is then whether it is possible to decouple and further suppress the coefficients of the $d = 5$ operators from those of the fermionic $d = 6$ operators, without appealing to fine-tunings and cancellations in the Yukawa parameters or heavy mass matrices.¹¹ If this is possible, the tiny values of the neutrino masses could be accommodated, while the effects of the those $d = 6$ operators — suppressed only as $1/M^2$ — would be close to observability.¹²

As a guideline to achieve such a scenario recall that, because Majorana neutrino masses are forbidden in the SM, light neutrinos inheritate their Majorana character from a Majorana source in the high-energy theory. This implies that light Majorana neutrino masses have to vanish either when the new Majorana scale goes to infinity and the new physics decouples, or proportionally to it. A quick look at table 3, together with the pattern of operator coefficients found for the case of scalar-triplet mediated Seesaw mechanisms, suggests the following *ansatz*:

When the breaking of L symmetry takes place in the full theory through a small mass scale μ , distinct from the high-energy scale $M \sim \mathcal{O}(\text{TeV})$, $\mu \ll M$, the coefficient of the $d = 5$ operator necessarily acquires an extra suppression in powers of μ/M , while the fermionic $d = 6$ operators keep its unsuppressed $1/M^2$ dependence.

As an example, a typical decoupling pattern goes qualitatively as follows:

$$c^{d=5} = f(Y) \frac{\mu}{M^2}, \quad (3.4)$$

$$c^{d=6} = g(Y) \frac{1}{|M|^2}, \quad (3.5)$$

where f and g are some functions of the Yukawa couplings, implying a light neutrino mass matrix of the form

$$m_\nu = -\frac{f(Y)v^2}{2} \frac{\mu}{M^2}, \quad (3.6)$$

while the effects of the $d = 6$ operator, eq. (3.5), are independent of μ and may be sizable for generic Yukawa couplings, which may remain large and even $\mathcal{O}(1)$. Notice that such dependence has already been found above for the minimal version of the scalar-triplet mediated Seesaw scenario (with $f \sim Y_\Delta$, $g \sim Y_\Delta^\dagger Y_\Delta$, $\mu = \mu_\Delta$), see eqs. (2.30) and (2.33),

¹¹Although operators of dimension higher than six are increasingly relevant as the scale is lowered toward the electroweak scale, an analysis restrained to the $d = 5$ and $d = 6$ operators should still convey the main physical aspects, as long as the scale keeps being larger than $\mathcal{O}(v)$.

¹²Note that Ref. [17] studied the effects of various higher-order operators in a completely different context: the dissociation of flavour violation scale and lepton number violation scale in extended theories, while we focus on the dissociation of $d=5$ and $d=6$ operators characteristic of the minimal seesaw models.

suggesting the possibility $\mu_\Delta \ll M_\Delta$, $Y_\Delta \sim 1$. We call this universal pattern *direct lepton violation*, since the neutrino masses are proportional to the (small) lepton number violating quantity μ , rather than inversely proportional to the large lepton number violating heavy field mass.

Multiple seesaw models. Let us consider, as illustration and support of our general ansatz, models existing in the literature and based on extensions of the type-I Seesaw scenario with a low scale. The examples considered below can be straightforwardly applied and extended to the type-III — triplet-fermion mediated — Seesaw scenarios. We are thus interested in a class of models which, to lead to sufficiently suppressed neutrino masses and large $d = 6$ operators, do not require any precise cancellations between the various (a priori independent) entries of the singlet neutrino mass matrix and/or of the Yukawa matrix.¹³ The cases considered below just require that some of the entries of these mass matrices carry Majorana character and are much smaller than other ones. For simplicity, only one left-handed neutrino and two singlet fermions will be included in the analysis (ν_L, N_1, N_2).

For instance, in the “inverse Seesaw model” [19], the following texture is assumed:¹⁴

$$\begin{pmatrix} 0 & m_{D_1} & 0 \\ m_{D_1} & 0 & M_{N_1} \\ 0 & M_{N_1} & \mu \end{pmatrix}, \quad (3.7)$$

where μ is a small Majorana mass, $\mu \ll M_{N_1}$. All other entries in the matrix are of Dirac character: for $\mu = 0$, assigning $L = 1, -1, 1$ to ν_L, N_1, N_2 respectively, lepton number is indeed conserved by the Lagrangian and no Majorana mass results. Expanding the eigenvalues of eq. (3.7) in powers of μ/M_{N_1} , a light eigenvalue is obtained:

$$m_\nu = \frac{m_{D_1}^2}{M_{N_1}} \frac{\mu}{M_{N_1}} \frac{M_{N_1}^2}{M_{N_1}^2 + m_{D_1}^2} + O(\mu^3) \xrightarrow{(m_{D_1} \ll M_{N_1})} \frac{m_{D_1}^2}{M_{N_1}} \frac{\mu}{M_{N_1}} + O(\mu^3), \quad (3.8)$$

where higher order terms have been neglected. As m_{D_1} is a typical Dirac mass term, $m_{D_1} \sim Y_1 v/\sqrt{2}$ with Y_1 a Yukawa coupling, eq. (3.8) shows that the neutrino mass is suppressed by an extra factor μ/M_{N_1} with respect to the result for the minimal type-I Seesaw model, eq. (2.5), exactly as expected from the general argument above, see eq. (3.6):

$$m_\nu \sim \mu \frac{Y_1^2 v^2}{M_1^2}. \quad (3.9)$$

The smallness of neutrino masses, and the argument of no fine-tuning, do not require tiny Yukawa couplings. For instance, if the Yukawa coupling Y_{ν_1} is of order unity, i.e. $m_{D_1} = Y_{\nu_1} v \sim v$, and if $M_{N_1} \sim 1$ TeV, this requires $\mu/M_{N_1} \sim 10^{-12}$. Similarly, for $M_{N_1} \sim 1$ TeV, a rather “large” Yukawa coupling of order 10^{-3} requires $\mu/M_{N_1} \sim 10^{-6}$. On the other hand, the $d = 6$ operator coefficient is independent of μ , as in eq. (3.5), and

¹³Consequently, we don’t consider cases such as, for example, that in ref. [18], based on the relation $Y_\nu Y_\nu^T = 0$ and $(M_N)_{ij} = m_N \delta_{ij}$.

¹⁴We acknowledge interesting discussions on this topic with S. Antusch.

low-energy effects associated to it — such as non-unitary mixings in the weak currents and other signals — could be discovered in the near future.

These results can be generalized to the case with the most general matrices which, with large Yukawa couplings, still lead to vanishing ν masses from extra small entries, therefore leading to suppressed $d = 5$ operator coefficients together with large $d = 6$ operator coefficients. For instance, in the two N plus one ν case above, the most general Majorana texture is the one in eq. (3.7) with an additional non-zero value for the 22 element. This can be justified for instance in the context of extended models (see e.g. ref. [20]). A non-zero 22 entry has the interesting feature of being a source of lepton number violation without inducing by itself neutrino masses: for $\mu = 0$ the determinant still vanishes leading to massless neutrinos. We will postpone the discussion of scenarios with a non-zero value for that entry to appendix C, where a generalization to the 3 left-handed plus 3 right-handed neutrino case can also be found. Analogous extensions of fermion-triplet mediated type-III Seesaw models are straightforward. The interesting textures are just the same as in the type I (that is, singlet-fermion Seesaw) scenario.

In conclusion, irrespective of whether the Seesaw mechanism results from the exchange of heavy fermions or heavy scalars, to have large effects from $d = 6$ operators requires first to lower the scale M toward the TeV range and second a decoupling of the values of the $d = 5$ and $d = 6$ coefficients along the pattern developed above, i.e. eq. (3.4) and eq. (3.5). This allows to account for the experimental values of neutrino masses without neither fine-tuning the Yukawa couplings nor assuming cancellations in combinations of them. For a Seesaw scale of $\mathcal{O}(\text{TeV})$, observable effects are then possible. The next section — which deals with the phenomenological aspects of Seesaw models including bounds for any value of M — will focus on those effects.

4. Phenomenology of seesaw models

4.1 Fermionic singlets

The models where the heavy fields are SM singlets are most difficult to test, as they lead to fewer and rarer experimental signals at low energies, even for low Seesaw scales. There exist, though, bounds on combinations of the Yukawa couplings which can be saturated for the type-I inverse Seesaw and similar extensions, as well as for models with extra dimensions containing Kaluza-Klein replicas which are SM singlets [21].

The bounds stem from important indirect signals which may be induced from the fact that the leptonic mass matrix appearing in the charged current is no longer unitary, see section 2.1. This subject, as well as the determination of the corresponding bounds on $|NN^\dagger|_{\alpha\beta}$, has been studied at length recently [10]. In a nutshell, deviations from the values expected in a unitary analysis are constrained to be of order 1% or smaller. Indeed, a global fit to the constraints resulting from W decays, Z decays, universality tests and rare lepton decays proved [10] that the NN^\dagger elements agree with those expected in the unitary

case, within a precision better than a few percent, at the 90% CL:

$$|NN^\dagger| \approx \begin{pmatrix} 0.994 \pm 0.005 & < 7.0 \cdot 10^{-5} & < 1.6 \cdot 10^{-2} \\ < 7.0 \cdot 10^{-5} & 0.995 \pm 0.005 & < 1.0 \cdot 10^{-2} \\ < 1.6 \cdot 10^{-2} & < 1.0 \cdot 10^{-2} & 0.995 \pm 0.005 \end{pmatrix}. \quad (4.1)$$

The off-diagonal constraints in eq. (4.1) result from the experimental bounds existing on the radiative processes $\mu \rightarrow e\gamma$, $\tau \rightarrow e\gamma$ and $\tau \rightarrow \mu\gamma$, while the diagonal ones come from the combined analysis of all other processes mentioned above. Using now the relation obtained in eq. (2.19) between the elements of the coefficient matrix $c^{d=6}$ and those of NN^\dagger , it follows that

$$\frac{v^2}{2} |c^{d=6}|_{\alpha\beta} = \frac{v^2}{2} \left| Y_N^\dagger \frac{1}{|M_N|^2} Y_N \right|_{\alpha\beta} \lesssim \begin{pmatrix} 10^{-2} & 7.0 \cdot 10^{-5} & 1.6 \cdot 10^{-2} \\ 7.0 \cdot 10^{-5} & 10^{-2} & 1.0 \cdot 10^{-2} \\ 1.6 \cdot 10^{-2} & 1.0 \cdot 10^{-2} & 10^{-2} \end{pmatrix}. \quad (4.2)$$

When obtaining the numerical bounds in eqs. (4.1) and (4.2), the effective theory was used to compute $\mu \rightarrow e\gamma$ and $\tau \rightarrow \mu\gamma$, that is, the analysis was done in terms of $c^{d=5}$ and $c^{d=6}$. It is to be noticed that the computation of such one-loop transitions in the effective theory does not coincide exactly with that done in the full theory (i.e. type I Seesaw model), as higher dimension effective operators have to be taken into account in the matching between both. Numerically, the differences are of $\mathcal{O}(1)$ and irrelevant for the precision attempted here, though. Furthermore, when computing these $l_1 \rightarrow l_2\gamma$ transitions - here and in the chapters to follow - we will not take into account the electromagnetic radiative corrections [22], as their inclusion would correspond to a two-loop calculation and numerically they will not change the order of magnitude of the bounds obtained.

Notice that the bounds above are valid for any value of M_N and apply to any (type I) Seesaw theory. In consequence, they apply to the inverse Seesaw model considered above, in which M_N could be near the TeV scale while the Yukawa couplings may be large, and signals could appear at the edge of the experimental limits above. New signals of CP-violation in neutrino oscillations, sensitive to the phases of $c^{d=6}$ may also be observable in future facilities [23].

As for direct detection of the heavy singlets in future accelerators in case $M_N \sim 1$ TeV, several studies exist of the associated production of the heavy singlets and the Higgs particle, with difficult prospects for a positive signal [24].

4.2 Scalar triplets

Using the experimental values for the renormalized parameters α , G_F and M_Z as defined in the Z-scheme in section 2.2, we will now consider deviations from the SM predictions for a set of observables.

4.2.1 M_W and the ρ parameter

The operator $\mathcal{L}_{\phi D}$ induces corrections to the predicted value of M_W and to the ρ parameter, through the term $4 \frac{|\mu_\Delta|^2}{|M_\Delta|^4} [\phi^\dagger D_\mu \phi]^\dagger [\phi^\dagger D_\mu \phi]$ in eq. (2.35). When the ρ parameter is

extracted from data using only hadronic transitions,¹⁵ its predicted value is shifted from the SM prediction by

$$\delta\rho_{\text{had}} = -\frac{|\mu_\Delta|^2}{M_\Delta^4} \frac{\sqrt{2}}{G_F}, \quad (4.3)$$

a result previously obtained in the literature [25].

We find that the mass of the W boson is also predicted to acquire a shift from both $\mathcal{L}_{\phi D}$ and \mathcal{L}_{4F} in eq. (2.33), which is given by

$$\begin{aligned} \delta M_W^2 &= -\frac{M_W^2}{2M_W^2 - M_Z^2} \left[\delta\rho_{\text{had}} M_W^2 - \frac{\delta G_F}{G_F} (M_W^2 - M_Z^2) \right] \\ &= -\frac{M_W^2}{2M_W^2 - M_Z^2} \left[\frac{|\mu_\Delta|^2}{M_\Delta^4} \frac{M_W^2}{G_F \sqrt{2}} - \frac{M_W^2 - M_Z^2}{\sqrt{2} G_F M_\Delta^2} Y_{\Delta e\mu} Y_{\Delta e\mu}^\dagger \right]. \end{aligned} \quad (4.4)$$

In this equation, M_W is to be identified with the SM prediction for the W -boson mass in the Z -scheme,

$$(M_W^{SM})^2 = \frac{M_Z^2}{2} \left(1 + \sqrt{1 - \frac{4\pi\alpha}{\sqrt{2} G_F M_Z^2}} \right) \frac{1}{(1 - \Delta r)}, \quad (4.5)$$

where Δr accounts for the dominant SM one-loop radiative corrections¹⁶, and G_F is extracted from muon decay, see eqs. (2.40) and (2.41).

The very precise experimental determination of the W boson mass allows to set stringent bounds on both terms in eq. (4.4), barring extreme cancellations between both. From the difference between the experimental value and the SM prediction of M_W obtained in the Z -scheme ($M_W^{SM} = (80.4887 \pm 0.0515) \text{ GeV}$) we obtain:

$$|Y_{\Delta e\mu}|^4 = (0.00023 \pm 0.00109) \left(\frac{M_\Delta}{1 \text{ TeV}} \right)^{-4}, \quad (4.6)$$

and

$$-v^2 \frac{|\mu_\Delta|^2}{M_\Delta^4} = 0.0001368 \pm 0.00032, \quad \text{or} \quad (4.7)$$

$$\frac{|\mu_\Delta|}{M_\Delta^2} < 8.7 \times 10^{-2} \text{ TeV}^{-1}. \quad (4.8)$$

Notice however that the hadronic data on the ρ parameter allow to independently constraint μ_Δ/M_Δ^2 . As an estimate, taking at face value the average experimental value of the ρ parameter ($\rho = 1.0002 \pm_{0.0004}^{0.0007}$) as if it were indeed dominated by the hadronic contributions — which do not depend on the leptonic Yukawa couplings — it would follow that

$$-v^2 \frac{|\mu_\Delta|^2}{M_\Delta^4} < 0.0001 \pm_{0.0002}^{0.00035}. \quad (4.9)$$

¹⁵It is customary to extract the value of ρ from a global fit to data, including simultaneously hadronic and leptonic transitions; if the latter were considered in the analysis, further corrections would appear in $\delta\rho$, induced by δG_F in eq. (2.40).

¹⁶While these corrections are important when compared to the total value of M_W , they can be dropped in eq. (4.4), as we work at first order in all corrections. ,PDG

The neutrino masses in eq. (2.31) are proportional to this ratio multiplied by Y_Δ . For instance, $m_\nu \sim 1\text{eV}$ and $Y_\Delta \sim \mathcal{O}(1)$ requires $\mu_\Delta/M_\Delta^2 \sim 10^{-11}$, well below the bound in eq. (4.8), while the lower limit of the bound can be saturated for values of $Y_\Delta \sim 10^{-7}$.

4.2.2 $\mu \rightarrow eee$ and $\tau \rightarrow 3l$ decays

$\delta\mathcal{L}_{4F}$ in eqs. (2.33) and (2.34) induces exotic four-lepton couplings contributing to lepton-flavor violating processes. Notice that this operator does not depend on the scale μ_Δ , so that the discussion is independent of it. Besides its impact on the determination of G_F from muon decay, eqs. (2.40) and (2.41), it modifies the branching ratios for rare leptonic decays. The constraints implied by the present experimental bounds on these processes have been studied in models with a scalar triplet in refs. [26]–[35].

Important decays are $\mu^- \rightarrow e^+e^-e^-$ and $\tau \rightarrow 3l$, considered in the full theory with a scalar triplet in refs. [27, 28] and from the generic leptonic $d = 6$ effective operator $\delta\mathcal{L}_{4F}$ in Refs.[36]–[39]. In terms of the coefficient $c_{\alpha\beta\gamma\delta}^{4F}$ of the leptonic $d = 6$ operator coefficient \mathcal{L}_{4F} in eq. (2.33),

$$c_{\alpha\beta\gamma\delta}^{4F} \equiv \frac{1}{M_\Delta^2} Y_{\Delta_{\alpha\beta}} Y_{\Delta_{\gamma\delta}}^\dagger, \quad (4.10)$$

we obtain

$$\Gamma(\mu^- \rightarrow e^+e^-e^-) = \frac{m_\mu^5}{192\pi^3} |c_{\mu e e e}^{4F}|^2 = \frac{m_\mu^5}{192\pi^3} \frac{1}{M_\Delta^4} |Y_{\Delta_{\mu e}}|^2 |Y_{\Delta_{ee}}|^2 \quad (4.11)$$

which gives

$$\text{Br}(\mu^- \rightarrow e^+e^-e^-) \simeq \frac{\Gamma(\mu^- \rightarrow e^+e^-e^-)}{\Gamma(\mu^- \rightarrow e^- \nu_\mu \bar{\nu}_e)} = \frac{|c_{\mu e e e}^{4F}|^2}{G_F^2} = \frac{1}{M_\Delta^4 G_F^2} |Y_{\Delta_{e\mu}}|^2 |Y_{\Delta_{ee}}|^2, \quad (4.12)$$

Similarly we obtain

$$\Gamma(\tau^- \rightarrow l_i^+ l_j^- l_j^-) = \frac{m_\tau^5}{192\pi^3} |c_{\tau i j j}^{4F}|^2 = \frac{m_\tau^5}{192\pi^3} \frac{1}{M_\Delta^4} |Y_{\Delta_{\tau i}}|^2 |Y_{\Delta_{jj}}|^2 \quad (4.13)$$

for any i and j , while

$$\Gamma(\tau^- \rightarrow l_i^+ l_j^- l_k^-) = \frac{m_\tau^5}{96\pi^3} |c_{\tau i j k}^{4F}|^2 = \frac{m_\tau^5}{96\pi^3} \frac{1}{M_\Delta^4} |Y_{\Delta_{\tau i}}|^2 |Y_{\Delta_{jk}}|^2 \quad (4.14)$$

for any i, j, k with $j \neq k$. Using all experimental branching ratios or upper limits on branching ratios as given in ref. [14], the corresponding bounds on the Yukawa couplings are given in table 4.

For Yukawa couplings of order unity, the present non-observation of those LFV transitions, in particular of the most stringent one, $\mu \rightarrow eee$, implies a lower bound on the scalar triplet Seesaw scale,

$$M_\Delta \geq 294 \text{ TeV}, \quad \text{for } Y_\Delta \sim \mathcal{O}(1). \quad (4.15)$$

Process	Constraint on	Bound $\left(\times \left(\frac{M_\Delta}{1\text{TeV}}\right)^2\right)$
M_W	$ Y_{\Delta\mu e} ^2$	$< 7.3 \times 10^{-2}$
$\mu^- \rightarrow e^+ e^- e^-$	$ Y_{\Delta\mu e} Y_{\Delta ee} $	$< 1.2 \times 10^{-5}$
$\tau^- \rightarrow e^+ e^- e^-$	$ Y_{\Delta\tau e} Y_{\Delta ee} $	$< 1.3 \times 10^{-2}$
$\tau^- \rightarrow \mu^+ \mu^- \mu^-$	$ Y_{\Delta\tau\mu} Y_{\Delta\mu\mu} $	$< 1.2 \times 10^{-2}$
$\tau^- \rightarrow \mu^+ e^- e^-$	$ Y_{\Delta\tau\mu} Y_{\Delta ee} $	$< 9.3 \times 10^{-3}$
$\tau^- \rightarrow e^+ \mu^- \mu^-$	$ Y_{\Delta\tau e} Y_{\Delta\mu\mu} $	$< 1.0 \times 10^{-2}$
$\tau^- \rightarrow \mu^+ \mu^- e^-$	$ Y_{\Delta\tau\mu} Y_{\Delta\mu e} $	$< 1.8 \times 10^{-2}$
$\tau^- \rightarrow e^+ e^- \mu^-$	$ Y_{\Delta\tau e} Y_{\Delta\mu e} $	$< 1.7 \times 10^{-2}$
$\mu \rightarrow e\gamma$	$ \Sigma_{l=e,\mu,\tau} Y_{\Delta l_\mu}^\dagger Y_{\Delta el} $	$< 4.7 \times 10^{-3}$
$\tau \rightarrow e\gamma$	$ \Sigma_{l=e,\mu,\tau} Y_{\Delta l_\tau}^\dagger Y_{\Delta el} $	< 1.05
$\tau \rightarrow \mu\gamma$	$ \Sigma_{l=e,\mu,\tau} Y_{\Delta l_\tau}^\dagger Y_{\Delta \mu l} $	$< 8.4 \times 10^{-1}$

Table 4: Bounds on $Y_{\Delta ij}$ from M_W , eq. (4.6), from tree level $\ell_1^- \rightarrow \ell_2^- \ell_3^+ \ell_4^-$ decays and from one loop $l_1 \rightarrow l_2 \gamma$ processes.

4.2.3 Complete lagrangian and $l_1 \rightarrow l_2 \gamma$

It is useful to consider also the bounds which arise from the radiative decays $\ell_1 \rightarrow \ell_2 \gamma$, although these processes cannot be obtained completely from the $d = 6$ operators because they are one-loop processes. Consider then instead the full high-energy Lagrangian for the scalar triplet in eq. (4.16), expanded into charge components:

$$\begin{aligned}
 \mathcal{L}_\Delta = & D_\mu \Delta^{0*} D^\mu \Delta^0 + D_\mu \Delta^+ D^\mu \Delta^- + D_\mu \Delta^{++} D^\mu \Delta^{--} \\
 & + \left\{ (\bar{l}_L^c Y_\Delta \nu_L) \Delta^+ + (\bar{\nu}_L^c Y_\Delta e_L) \Delta^+ + \sqrt{2} (\bar{l}_L^c Y_\Delta l_L) \Delta^{++} - \sqrt{2} (\bar{\nu}_L^c Y_\Delta \nu_L) \Delta^0 + \text{h.c.} \right\} \\
 & + \left\{ \mu_\Delta \left(2\phi^0 \phi^+ \Delta^- + \sqrt{2} \phi^0 \phi^0 \Delta^{0*} - \sqrt{2} \phi^+ \phi^0 \Delta^{--} \right) + \text{h.c.} \right\} \\
 & - M_\Delta^2 \left(\Delta^{0*} \Delta^0 + \Delta^- \Delta^+ + \Delta^{--} \Delta^{++} \right) - \frac{\lambda_2}{2} \left(\Delta^{0*} \Delta^0 + \Delta^- \Delta^+ + \Delta^{--} \Delta^{++} \right)^2 \\
 & - \lambda_3 (\phi^- \phi^+ + \phi^{0*} \phi^0) \left(\Delta^{0*} \Delta^0 + \Delta^- \Delta^+ + \Delta^{--} \Delta^{++} \right) - \lambda_4 \left[\frac{1}{2} (\Delta^{0*} \Delta^0)^2 + \frac{1}{2} (\Delta^{++} \Delta^{--})^2 \right. \\
 & \left. + \Delta^{++} \Delta^{--} (\Delta^+ \Delta^- - \Delta^{0*} \Delta^0) + \Delta^+ \Delta^- \Delta^{0*} \Delta^0 - \{ \Delta^{++} \Delta^- \Delta^- \Delta^0 + \text{h.c.} \} \right] \\
 & - \lambda_5 \left[(\Delta^{++} \Delta^{--} - \Delta^{0*} \Delta^0) (\phi^+ \phi^- - \phi^{0*} \phi^0) - \sqrt{2} \{ \phi^- \phi^0 (\Delta^{++} \Delta^- - \Delta^{0*} \Delta^+) + \text{h.c.} \} \right].
 \end{aligned}$$

$\mu \rightarrow e\gamma, \tau \rightarrow e\gamma, \tau \rightarrow \mu\gamma$. Radiative processes are due to the exchange between lepton fields of both the Δ^{++} and Δ^+ fields, as given in eq. (4.16), and the branching ratios

Combined bounds		
Process	Yukawa	Bound $\left(\times \left(\frac{M_\Delta}{1\text{TeV}}\right)^4\right)$
$\mu \rightarrow e\gamma$	$ Y_{\Delta\mu\mu}^\dagger Y_{\Delta\mu e} + Y_{\Delta\tau\mu}^\dagger Y_{\Delta\tau e} $	$< 4.7 \times 10^{-3}$
$\tau \rightarrow e\gamma$	$ Y_{\Delta\tau\tau}^\dagger Y_{\Delta\tau e} $	< 1.05
$\tau \rightarrow \mu\gamma$	$ Y_{\Delta\tau\tau}^\dagger Y_{\Delta\tau\mu} $	$< 8.4 \times 10^{-1}$

Table 5: Bounds on combinations of $Y_{\Delta ij}$.

read [29, 31, 34]:

$$\text{Br}(l_1 \rightarrow l_2 \gamma) = \frac{\alpha}{48\pi} \frac{25}{16} \frac{\left| \sum_l Y_{\Delta l l_1}^\dagger Y_{\Delta l l_2} \right|^2}{G_F^2 M_\Delta^4} \text{Br}(l_1 \rightarrow e \nu_1 \bar{\nu}_e). \quad (4.16)$$

The corresponding bounds are also given in table 4. Combining all bounds of this table, we have obtained new bounds for combinations of Yukawa parameters, not considered previously in the literature and gathered in table 5. They show that, for low values of the Seesaw scale, the Yukawa couplings may be of $\mathcal{O}(1)$, while they should be sizeably smaller by up to 2 orders of magnitude for some specific flavours, for an $\mathcal{O}(\text{TeV})$ Seesaw scale.

4.2.4 Other constraints

There are also other bounds which arise from other processes, from Bhabha scattering [27, 32, 34] (leading to the bound $|Y_{\Delta ee}| < 1.0 \cdot (M_\Delta/1\text{TeV})$), from muonium to antimuonium conversion [27, 35] (leading to the bound $|Y_{\Delta ee}| |Y_{\Delta\mu\mu}| < 0.1 \cdot (M_\Delta/1\text{TeV})^2$), and from the anomalous magnetic moment of the muon. The latter constraint comes from the fact that the doubly charged scalar $\Delta^{\pm\pm}$ as well as the simply charged Δ^\pm induce a shift in the anomalous magnetic moment of the muon [32, 35],

$$\delta(a_\mu) = -\frac{m_\mu^2}{3\pi M_\Delta^2} \sum_{j=e,\mu'\tau} |Y_{\Delta\mu j}|^2, \quad \delta(a_\mu) = -\frac{m_\mu^2}{24\pi M_\Delta^2} \sum_{j=e,\mu'\tau} |Y_{\Delta\mu j}|^2, \quad (4.17)$$

respectively. This contribution has opposite sign with respect to the observed deviation. Taking for instance $\delta(a_\mu) < 20 \times 10^{-10}$ we get $\sum_{j=e,\mu'\tau} |Y_{\Delta\mu j}|^2 < 1.9 \cdot (M_\Delta/1\text{TeV})^2$.

4.2.5 Collider signatures of scalar triplets

Scalar triplet Seesaw opens the possibility of observing new signals at present and/or future facilities. For instance, for $Y_\Delta \sim \mathcal{O}(1)$, a positive observation of $\mu \rightarrow e\gamma$ by the MEG experiment [40] (which aims to achieve 10^{-13} sensitivity for the branching ratio) would require

$$15\text{ TeV} < M_\Delta < 50\text{ TeV}, \quad (4.18)$$

while $Y_\Delta \sim \mathcal{O}(10^{-2})$ would require $0.15\text{ TeV} < M_\Delta < 0.5\text{ TeV}$.

If M_Δ turns out to be as low as $\mathcal{O}(\text{TeV})$, the non-vanishing electroweak charge of the $\vec{\Delta}$ field offers the possibility of clean signals in hadronic accelerators (Tevatron, LHC). The production (or associated production) of Δ^{++} and Δ^{--} particles, and their subsequent decay in pairs of same-sign leptons, would constitute striking signals, free from SM backgrounds [41, 42]. A lower bound on the mass of $\Delta^{\pm\pm}$ of the order of 136 GeV has been obtained at CDF [43]. Assuming that a boson with those characteristics is indeed observed in an accelerator, one still needs to ascertain whether a scalar-mediated Seesaw mechanism is indeed at work. For that, and as a first step, it is necessary to measure and disentangle the Yukawa couplings appearing in tables 4 and 5. In order to extract values for the individual $Y_{\Delta ij}$, it would be necessary to observe in addition at least three lepton flavor violating processes.

The first term in the Lagrangian in eq. (2.25) generates tree-level vertices $\Delta^{\pm\pm}W^\pm W^\pm$ and $\Delta^\pm W^\mp Z$, which would be detected by observing for instance $\Delta^{++} \rightarrow W^+W^+$, $W^+W^+ \rightarrow \Delta^{++}$, $Z^* \rightarrow \Delta^+W^-$, or $\Delta^+ \rightarrow ZW^+$. The analysis of some of these processes has already been covered in [44, 45], for the LHC. Once produced by Drell-Yann processes ($\bar{q}q \rightarrow \Delta^{--}\Delta^{++}$), with production cross-section given in refs. [41, 42], the Δ^{++} (Δ^{--}) can decay into pairs of W s of the same charge ($\Delta^{\mp\mp} \rightarrow W^\mp W^\mp$), for which the decay rate is proportional to $v^2 \frac{M_\Delta^3}{M_W^2} \frac{|\mu_\Delta|^2}{M_\Delta^4}$, or into leptons l_i, l_j of the same charge, with a decay rate proportional to $M_\Delta |Y_{\Delta ij}|^2$. Finally, the Δ^{++} (Δ^{--}) particle can also decay into a charged Higgs and an off-shell W gauge boson, as in $\Delta^{--} \rightarrow \phi^- W^{*-}$. The decay rate of the latter process is suppressed when compared to the previous ones, unless the λ_5 coupling in eq. (2.25) takes an unnaturally large value [44, 46]. Due to the constraint obtained in eq. (4.8), the process $\Delta^{--} \rightarrow W^-W^-$ will be suppressed and the only relevant channel in our scenario will be $\Delta^{\pm\pm} \rightarrow l^\pm l^\pm$, which will be background-free. The related branching ratio will give access to $|Y_{\Delta ij}|$, which is directly related to neutrino mass matrix elements up to the global factor $\frac{|\mu_\Delta|}{M_\Delta^2}$, i.e. to the effective theory coefficient $c^{d=5}$ in eq. (2.30).

Other interesting signals can be also searched for in accelerators. $\mathcal{L}_{6\phi}$ (eq. (2.33)) and the first term of $\mathcal{L}_{\phi D}$ in eq. (2.35), besides modifying the Higgs potential and renormalizing the scalar kinetic energy term, induce new couplings: $HHW, HZZ, HHWW, HHZZ, H^3$ and H^4 -where H stands for the physical Higgs-. Consequently, the Higgs production cross sections at the future facilities ILC and CLIC [47] get corrections. Nevertheless, the strong limit in eq. (4.8) precludes observable effects, except maybe from $\mathcal{L}_{6\phi}$ for very large values of λ_3 and/or λ_5 [47].

Similarly, $\mathcal{L}_{\phi D}$ also affects Higgs physics. Its impact on the Higgs decay branching ratios has been analyzed [48], although again the bound from the ρ parameter discussed above excludes observation in the planned future facilities such as ILC.

4.3 Fermionic triplets

We have argued that non-unitary flavour-changing matrices are to be expected in this case for the couplings of light leptons to the W and Z gauge boson, see eqs. (2.58)–(2.60). The putative departures from unitarity can be re-expressed directly in terms of the $d = 6$ operator coefficients, that is to say in terms of the Yukawa couplings, see eq. (2.64).

Specifically, notice that¹⁷

$$|NN^\dagger - 1| = |\epsilon^\Sigma|, \quad (4.19)$$

$$(N^\dagger N)^{-1} = U_\nu^\dagger (1 - \epsilon^\Sigma) U_\nu \sim 1 - U_{PMNS}^\dagger \epsilon^\Sigma U_{PMNS}. \quad (4.20)$$

For values of M_Σ close to the electroweak scale, the deviations of these quantities from their standard values can be at the edge of the present experimental bounds on non-unitarity. Taking into account the shift induced on G_F as extracted from muon decay, eq. (2.63) for the effective theory, we proceed to compute below the departures predicted on leptonic processes in the effective and full theories.

As we will see, all transitions considered below result in constraints on the elements of the NN^\dagger matrix — and thus on the $d = 6$ operator coefficients — analogously to the situation for fermionic singlet Seesaw theories [10], see for instance eq. (4.2). Indeed, even if we could have expected that Z -mediated processes are sensitive also to U_{PMNS} through eq. (4.20), this is not the case, as we will show in section 4.3.2.

4.3.1 W decays

The non-unitary mixing matrix N appearing now in the charged weak couplings, eq. (2.58), results in a leptonic W decay width of the form

$$\Gamma(W \rightarrow l_\alpha \nu_\alpha) = \sum_i \Gamma(W \rightarrow l_\alpha \nu_i) = \frac{G_F^{SM} M_W^3}{6\sqrt{2}\pi} (NN^\dagger)_{\alpha\alpha}. \quad (4.21)$$

Using the value of G_F extracted from the decay $\mu \rightarrow \nu_\mu e \bar{\nu}_e$, as given in eq. (2.63), the following combinations can be defined:

$$\frac{(NN^\dagger)_{\alpha\alpha}}{\sqrt{(NN^\dagger)_{ee}(NN^\dagger)_{\mu\mu}}} = \frac{\Gamma(W \rightarrow l_\alpha \nu_\alpha) 6\sqrt{2}\pi}{G_F M_W^3} \equiv f_\alpha. \quad (4.22)$$

With the experimental values of the W decay widths and mass from ref. [14] and $G_F = (1.16637 \pm 0.00001) \times 10^{-5}$, the parameters f_α take the values:

$$\begin{aligned} f_e &= 1.000 \pm 0.024, \\ f_\mu &= 0.986 \pm 0.028, \\ f_\tau &= 1.002 \pm 0.032. \end{aligned} \quad (4.23)$$

4.3.2 Invisible Z decay

The modified neutral weak couplings in eqs. (2.59) and (2.60) lead to

$$\Gamma(Z \rightarrow \text{invisible}) = \sum_{i,j} \Gamma(Z \rightarrow \bar{\nu}_i \nu_j) = \frac{G_F^{SM} M_Z^3}{12\sqrt{2}\pi} (1 + \rho_t) \sum_{i,j} |(N^\dagger N)^{-1}]_{ij}|^2, \quad (4.24)$$

where $\rho_t \approx 0.008$ [14] takes into account radiative corrections mainly stemming from loops including the top quark. As the dominant radiative corrections do not involve leptons, the

¹⁷Again, the absolute-value bars in eq. (4.19) can be dropped when choosing the appropriate basis in flavour space, see the discussion in section 2.3.

Constraints on	Process	Bound
$\frac{(NN^\dagger)_{\mu\mu}}{(NN^\dagger)_{ee}}$	$\frac{\Gamma(W \rightarrow \mu\bar{\nu}_\mu)}{\Gamma(W \rightarrow e\bar{\nu}_e)}$	0.997 ± 0.010
$\frac{(NN^\dagger)_{\tau\tau}}{(NN^\dagger)_{ee}}$	$\frac{\Gamma(W \rightarrow \tau\bar{\nu}_\tau)}{\Gamma(W \rightarrow e\bar{\nu}_e)}$	1.034 ± 0.0014
$\frac{(NN^\dagger)_{\mu\mu}}{(NN^\dagger)_{ee}}$	$\frac{\Gamma(\pi \rightarrow \mu\bar{\nu}_\mu)}{\Gamma(\pi \rightarrow e\bar{\nu}_e)}$	1.0017 ± 0.0015
$\frac{(NN^\dagger)_{\tau\tau}}{(NN^\dagger)_{\mu\mu}}$	$\frac{\Gamma(\tau \rightarrow \pi\bar{\nu}_\tau)}{\Gamma(\pi \rightarrow \mu\bar{\nu}_\mu)}$	0.9999 ± 0.0036
$\frac{(NN^\dagger)_{\mu\mu}}{(NN^\dagger)_{ee}}$	$\frac{\Gamma(\tau \rightarrow \nu_\tau \mu\bar{\nu}_\mu)}{\Gamma(\tau \rightarrow \nu_\tau e\bar{\nu}_e)}$	0.9999 ± 0.0020
$\frac{(NN^\dagger)_{\tau\tau}}{(NN^\dagger)_{\mu\mu}}$	$\frac{\Gamma(\tau \rightarrow \nu_\tau e\bar{\nu}_e)}{\Gamma(\mu \rightarrow \nu_\mu e\bar{\nu}_e)}$	1.0004 ± 0.0023
$\frac{(NN^\dagger)_{\tau\tau}}{(NN^\dagger)_{ee}}$	$\frac{\Gamma(\tau \rightarrow \nu_\tau \mu\bar{\nu}_\mu)}{\Gamma(\mu \rightarrow \nu_\mu e\bar{\nu}_e)}$	1.0002 ± 0.0022

Table 6: Constraints on $(NN^\dagger)_{\alpha\alpha}$ from a selection of processes.

dependence on the mixing matrix in eq. (4.24) appears as a global factor to an excellent approximation. Using the data provided in ref. [14] and the following approximation valid at first order in ϵ^Σ

$$\sum_{i,j} |[(N^\dagger N)^{-1}]_{ij}|^2 = \text{Tr}(1 - 2\epsilon^\Sigma) = 9 - 2 \sum_{\alpha} (NN^\dagger)_{\alpha\alpha}, \quad (4.25)$$

the following constraint is then obtained:

$$\frac{9 - 2 \sum_{\alpha} (NN^\dagger)_{\alpha\alpha}}{\sqrt{(NN^\dagger)_{ee}(NN^\dagger)_{\mu\mu}}} = \frac{12\sqrt{2}\pi \Gamma(Z \rightarrow \text{invisible})}{G_F^M M_Z^3 (1 + \rho_t)} = 2.984 \pm 0.009. \quad (4.26)$$

As it is well known, this number should correspond to the number of active neutrinos at LEP. Its 2σ departure from the value of 3 is not (yet) significant enough to be interpreted as a signal of new physics.

4.3.3 Universality tests

The existing constraints on the universality of weak interactions can be turned into bounds on non-unitarity if the weak couplings are indeed universal, as it is the case in Seesaw models. The results of our analysis, always at order ϵ^Σ , are displayed in table 6, where the bounds have been extracted from ref. [49]. For the leptonic decays, the following expression has been used (for $\alpha \neq \beta$):

$$\Gamma(l_\alpha \rightarrow \nu_\alpha l_\beta \bar{\nu}_\beta) = \frac{G_F^{SM} m_\alpha^5}{192\pi^3} (NN^\dagger)_{\alpha\alpha} (NN^\dagger)_{\beta\beta}. \quad (4.27)$$

Charged pion decays to a lepton pair are also considered in that table.

4.3.4 Z decays into charged leptons

While the processes analyzed in the previous sections permit to put bounds on the diagonal elements of (NN^\dagger) , as in the case of the fermionic singlets, the additional presence of flavour changing effects in the coupling of charged fermions to the Z boson allows to constrain the

Constraints on	Process	Bound
$ (NN^\dagger)_{e\mu} $	$\text{Br}(Z \rightarrow e^\pm \mu^\mp)$	$< 2.5 \cdot 10^{-3}$
$ (NN^\dagger)_{e\tau} $	$\text{Br}(Z \rightarrow e^\pm \tau^\mp)$	$< 6.1 \cdot 10^{-3}$
$ (NN^\dagger)_{\mu\tau} $	$\text{Br}(Z \rightarrow \mu^\pm \tau^\mp)$	$< 6.7 \cdot 10^{-3}$

Table 7: Constraints on $(NN^\dagger)_{\alpha\beta}$ from tree-level Z decays into charged leptons.

off-diagonal elements of (NN^\dagger) with tree-level processes, at variance with the fermionic singlet case. The leptonic width of the Z gauge boson is given by

$$\Gamma(Z \rightarrow l_\alpha \bar{l}_\alpha) = \frac{G_F M_Z^3}{3\sqrt{2}\pi} (|\sin^2 \theta_W|^2 + |\sin^2 \theta_W - \frac{1}{2}|[(NN^\dagger)^2]_{\alpha\alpha}|^2), \quad (4.28)$$

where the first (second) term in the parenthesis is the contribution of right-handed (left-handed) leptons. For $\alpha \neq \beta$ it follows that:

$$\Gamma(Z \rightarrow l_\alpha \bar{l}_\beta) = \frac{G_F^{SM} M_Z^3}{3\sqrt{2}\pi} \frac{1}{4} |[(NN^\dagger)^2]_{\alpha\beta}|^2. \quad (4.29)$$

It is now possible to obtain the branching ratios at leading order in ϵ^Σ :

$$\begin{aligned} \text{Br}(Z \rightarrow l_\alpha \bar{l}_\beta) &= \frac{\Gamma(Z \rightarrow l_\alpha \bar{l}_\beta)}{\Gamma(Z \rightarrow l_\gamma \bar{l}_\gamma)} \text{Br}(Z \rightarrow l_\gamma \bar{l}_\gamma) = \\ &= \frac{|(NN^\dagger)_{\alpha\beta}|^2}{2\sin^4 \theta_W - \sin^2 \theta_W + 1/4} \text{Br}(Z \rightarrow l_\gamma \bar{l}_\gamma), \end{aligned} \quad (4.30)$$

where we have used $|[(NN^\dagger)^2]_{\alpha\beta}|^2 = 4|(NN^\dagger)_{\alpha\beta}|^2$ and $\sin^2 \theta_W = 0.23$ is the Weinberg angle. From this, the bounds in table 7 have been derived.

4.3.5 $\mu \rightarrow eee$ and $\tau \rightarrow 3l$ decays

The non-unitarity of the leptonic mixing matrix N results, in the case of the fermionic triplet Seesaw theory under study, in tree-level $\mu \rightarrow 3e$ transitions given by (at leading order in ϵ^Σ):

$$\begin{aligned} \text{Br}(\mu^- \rightarrow e^+ e^- e^-) &\simeq \frac{\Gamma(\mu^- \rightarrow e^+ e^- e^-)}{\Gamma(\mu^- \rightarrow e^- \nu_\mu \bar{\nu}_e)} \\ &= |[(NN^\dagger)^2]_{e\mu}|^2 \left(3\sin^4 \theta_W - 2\sin^2 \theta_W + \frac{1}{2} \right). \end{aligned} \quad (4.31)$$

Analogously, τ decays in $3e$ or 3μ are non-zero and given by:

$$\begin{aligned} \text{Br}(\tau^- \rightarrow l_\alpha^+ l_\alpha^- l_\alpha^-) &= \frac{\Gamma(\tau^- \rightarrow l_\alpha^+ l_\alpha^- l_\alpha^-)}{\Gamma(\tau^- \rightarrow e^- \nu_\tau \bar{\nu}_e)} \text{Br}(\tau^- \rightarrow e^- \nu_\tau \bar{\nu}_e) \\ &= |[(NN^\dagger)^2]_{\alpha\tau}|^2 \left(3\sin^4 \theta_W - 2\sin^2 \theta_W + \frac{1}{2} \right) \text{Br}(\tau^- \rightarrow e^- \nu_\tau \bar{\nu}_e), \end{aligned} \quad (4.32)$$

Constraints on	Process	Bound
$ (NN^\dagger)_{e\mu} $	$\mu^- \rightarrow e^+ e^- e^-$	$< 1.1 \cdot 10^{-6}$
$ (NN^\dagger)_{e\tau} $	$\tau^- \rightarrow e^+ e^- e^-$	$< 1.2 \cdot 10^{-3}$
$ (NN^\dagger)_{\mu\tau} $	$\tau^- \rightarrow \mu^+ \mu^- \mu^-$	$< 1.2 \cdot 10^{-3}$
$ (NN^\dagger)_{\tau e} $	$\tau^- \rightarrow \mu^+ \mu^- e^-$	$< 1.6 \cdot 10^{-3}$
$ (NN^\dagger)_{\tau\mu} (NN^\dagger)_{e\mu} $	$\tau^- \rightarrow e^+ \mu^- \mu^-$	$< 3.1 \cdot 10^{-4}$
$ (NN^\dagger)_{\tau\mu} $	$\tau^- \rightarrow e^+ e^- \mu^-$	$< 1.5 \cdot 10^{-3}$
$ (NN^\dagger)_{\tau e} (NN^\dagger)_{\mu e} $	$\tau^- \rightarrow \mu^+ e^- e^-$	$< 2.9 \cdot 10^{-4}$
$ (NN^\dagger)_{e\mu} $	$\mu \rightarrow e\gamma$	$2.8 \cdot 10^{-5}$
$ (NN^\dagger)_{\mu\tau} $	$\tau \rightarrow \mu\gamma$	$5.2 \cdot 10^{-3}$
$ (NN^\dagger)_{e\tau} $	$\tau \rightarrow e\gamma$	$6.6 \cdot 10^{-3}$

Table 8: Constraints on $(NN^\dagger)_{\alpha\beta}$ from charged leptons decays.

where $\alpha = \mu, e$. On the other side, τ decays in $2e(\mu) + 1\mu(e)$ are given by:

$$\begin{aligned} \text{Br}(\tau^- \rightarrow l_\alpha^+ l_\alpha^- l_\beta^-) &= \frac{\Gamma(\tau^- \rightarrow l_\alpha^+ l_\alpha^- l_\beta^-)}{\Gamma(\tau^- \rightarrow l_\delta^- \nu_\tau \bar{\nu}_e)} \text{Br}(\tau^- \rightarrow l_\delta^- \nu_\tau \bar{\nu}_e) \\ &= |[(NN^\dagger)^2]_{\beta\tau}|^2 \left(2\sin^4 \theta_W - \sin^2 \theta_W + \frac{1}{4} \right) \text{Br}(\tau^- \rightarrow l_\delta^- \nu_\tau \bar{\nu}_e), \end{aligned} \quad (4.33)$$

$$\begin{aligned} \text{Br}(\tau^- \rightarrow l_\beta^+ l_\alpha^- l_\alpha^-) &= \frac{\Gamma(\tau^- \rightarrow l_\beta^+ l_\alpha^- l_\alpha^-)}{\Gamma(\tau^- \rightarrow e^- \nu_\tau \bar{\nu}_e)} \text{Br}(\tau^- \rightarrow e^- \nu_\tau \bar{\nu}_e) \\ &= \frac{1}{2} |[(NN^\dagger)^2]_{\alpha\tau}|^2 |[(NN^\dagger)^2]_{\alpha\beta}|^2 \text{Br}(\tau^- \rightarrow e^- \nu_\tau \bar{\nu}_e), \end{aligned} \quad (4.34)$$

where $\alpha, \beta = \mu, e$ with $\alpha \neq \beta$. The bounds resulting from these processes for combinations of NN^\dagger elements are contained in table 8.

4.3.6 Complete lagrangian and $l_1 \rightarrow l_2 \gamma$

As the phenomenological consequences of Seesaw scenarios mediated by SU(2) fermionic triplets remain almost unexplored in the literature, it is worth to study in detail the complete Lagrangian for the high-energy theory in eq. (2.45), developing it in terms of the electrically charged components¹⁸ of $\vec{\Sigma}$,

$$\begin{aligned} \mathcal{L} &= \overline{\Sigma_R^+} i \not{\partial} \Sigma_R^+ + \overline{\Sigma_R^-} i \not{\partial} \Sigma_R^- + \overline{\Sigma_R^0} i \not{\partial} \Sigma_R^0 \\ &+ g \left(W_\mu^+ \overline{\Sigma_R^0} \gamma_\mu \Sigma_R^- - W_\mu^+ \overline{\Sigma_R^+} \gamma_\mu \Sigma_R^0 + \text{h.c.} \right) + g \left(W_\mu^3 \overline{\Sigma_R^+} \gamma_\mu \Sigma_R^+ - W_\mu^3 \overline{\Sigma_R^-} \gamma_\mu \Sigma_R^- \right) \\ &- \frac{1}{2} \left(\overline{\Sigma_R^+} M_\Sigma \Sigma_R^{-c} + \overline{\Sigma_R^-} M_\Sigma \Sigma_R^{+c} + \overline{\Sigma_R^0} M_\Sigma \Sigma_R^{0c} + \text{h.c.} \right) \\ &- \left(\phi^0 \overline{\Sigma_R^0} Y_\Sigma \nu_L + \sqrt{2} \phi^0 \overline{\Sigma_R^-} Y_\Sigma l_L + \phi^+ \overline{\Sigma_R^0} Y_\Sigma l_L - \sqrt{2} \phi^+ \overline{\Sigma_R^+} Y_\Sigma \nu_L + \text{h.c.} \right). \end{aligned} \quad (4.35)$$

This Lagrangian, in which the charged components of the triplets are expressed in terms of 2-component fields, is not convenient when considering mixing with the charged leptons,

¹⁸Note that the charged conjugate of Σ_R^\pm is not Σ_R^\mp but $\Sigma_R^{\pm c}$.

which as usual are expressed in 4-component notation. As the charged triplet components have 4 degrees of freedom they can all be written in terms of a 4-component unique Dirac spinor,

$$\Psi \equiv \Sigma_R^{+c} + \Sigma_R^- . \quad (4.36)$$

The neutral fermionic triplet components on the other hand can be left in 2-component notation, since they have only two degrees of freedom and mix with the neutrinos, which are also described by 2-component fields. This leads to the Lagrangian

$$\begin{aligned} \mathcal{L} = & \overline{\Psi} i \not{\partial} \Psi + \overline{\Sigma}_R^0 i \not{\partial} \Sigma_R^0 - \overline{\Psi} M_\Sigma \Psi - \left(\overline{\Sigma}_R^0 \frac{M_\Sigma}{2} \Sigma_R^{0c} + \text{h.c.} \right) \\ & + g \left(W_\mu^+ \overline{\Sigma}_R^0 \gamma_\mu P_R \Psi + W_\mu^+ \overline{\Sigma}_R^{0c} \gamma_\mu P_L \Psi + \text{h.c.} \right) - g W_\mu^3 \overline{\Psi} \gamma_\mu \Psi \\ & - \left(\phi^0 \overline{\Sigma}_R^0 Y_\Sigma \nu_L + \sqrt{2} \phi^0 \overline{\Psi} Y_\Sigma l_L + \phi^+ \overline{\Sigma}_R^0 Y_\Sigma l_L - \sqrt{2} \phi^+ \overline{\nu}_L^c Y_\Sigma^T \Psi + \text{h.c.} \right) . \end{aligned} \quad (4.37)$$

The mass term of the charged sector shows then the usual aspect for Dirac particles (omitting flavor indices):

$$\mathcal{L} \ni -(\overline{l}_R \ \overline{\Psi}_R) \begin{pmatrix} m_l & 0 \\ Y_\Sigma v & M_\Sigma \end{pmatrix} \begin{pmatrix} l_L \\ \Psi_L \end{pmatrix} - (\overline{l}_L \ \overline{\Psi}_L) \begin{pmatrix} m_l & Y_\Sigma^\dagger v \\ 0 & M_\Sigma \end{pmatrix} \begin{pmatrix} l_R \\ \Psi_R \end{pmatrix} , \quad (4.38)$$

The — symmetric — mass matrix for the neutral states is on the other hand given by:

$$\begin{aligned} \mathcal{L} \ni & -(\overline{\nu}_L \ \overline{\Sigma}^{0c}) \begin{pmatrix} 0 & Y_\Sigma^\dagger v / 2\sqrt{2} \\ Y_\Sigma^* v / 2\sqrt{2} & M_\Sigma / 2 \end{pmatrix} \begin{pmatrix} \nu_L^c \\ \Sigma^0 \end{pmatrix} \\ & -(\overline{\nu}_L^c \ \overline{\Sigma}^0) \begin{pmatrix} 0 & Y_\Sigma^T v / 2\sqrt{2} \\ Y_\Sigma v / 2\sqrt{2} & M_\Sigma / 2 \end{pmatrix} \begin{pmatrix} \nu_L \\ \Sigma^{0c} \end{pmatrix} . \end{aligned} \quad (4.39)$$

The corresponding mixing matrices, necessary to calculate $\mu \rightarrow e\gamma$ and similar processes, are explicitly given in appendix B.

$\mu \rightarrow e\gamma$, $\tau \rightarrow e\gamma$ and $\tau \rightarrow \mu\gamma$. $l_1 \rightarrow l_2\gamma$ transitions result from Z and W -mediated one-loop processes, depicted in figure 2. The amplitude of the matrix element, computed within the complete theory, eqs. (2.45) and (4.37), is given by (the details of the computation will be given in a separate publication [15]):

$$\begin{aligned} \mathcal{A}_{l_1 \rightarrow l_2 \gamma} = & -i \frac{G_F^{SM}}{\sqrt{2}} \frac{e}{16\pi^2} m_\mu \overline{u}_e(p-q) P_R \sigma_{\lambda\nu} q^\nu \epsilon^\lambda u_\mu(p) \times \\ & \left\{ C \epsilon_{21}^\Sigma + \sum_i x_{\nu_i} (U_{0\nu\nu})_{2i} \left((U_{0\nu\nu})^\dagger \right)_{i1} + \mathcal{O} \left(\frac{1}{M_\Sigma^4} \right) \right\} \end{aligned} \quad (4.40)$$

In this equation, $C = 2.23$, $x_{\nu_i} = \frac{m_{\nu_i}^2}{M_W^2}$ and ϵ_{21}^Σ corresponds to the $d = 6$ operator coefficients $\epsilon_{e\mu}^\Sigma$, $\epsilon_{e\tau}^\Sigma$ and $\epsilon_{\mu\tau}^\Sigma$ in eq. (2.53), when considering $\mu \rightarrow e\gamma$, $\tau \rightarrow e\gamma$ and $\tau \rightarrow \mu\gamma$ transitions, respectively. $U_{0\nu\nu}$ is the unitary matrix which diagonalizes the neutral lepton mass matrix




Figure 2: Diagrams contributing to $\mu \rightarrow e\gamma$. ϕ^- is the Goldstone boson associated with the W^- boson, η is the Goldstone boson associated with the Z boson and H stands for the physical Higgs boson.

for the fields (ν_L, Σ^{0c}) , see appendix B for details. Using these results, the branching ratio for the $l_1 \rightarrow l_2\gamma$ transition is given by (at order $1/M_\Sigma^4$):

$$BR(l_1 \rightarrow l_2\gamma) = \frac{3}{32} \frac{\alpha}{\pi} \frac{\left| C \epsilon_{21}^\Sigma + \sum_i x_{\nu_i} (U_{0\nu\nu})_{2i} \left((U_{0\nu\nu})^\dagger \right)_{i1} \right|^2}{(NN^\dagger)_{11}(NN^\dagger)_{22}} \quad (4.41)$$

The experimental bounds on these processes result in constraints given in table. 8. These are comparable to those stemming from tree-level purely leptonic decays.

4.3.7 Combination of all constraints

From all constraints obtained above we have performed a global fit, and the following bounds on the NN^\dagger elements have been derived, at the 90% CL.:

$$|NN^\dagger| \approx \begin{pmatrix} 1.001 \pm 0.002 & < 1.1 \cdot 10^{-6} & < 1.2 \cdot 10^{-3} \\ < 1.1 \cdot 10^{-6} & 1.002 \pm 0.002 & < 1.2 \cdot 10^{-3} \\ < 1.2 \cdot 10^{-3} & < 1.2 \cdot 10^{-3} & 1.002 \pm 0.002 \end{pmatrix}. \quad (4.42)$$

Using now the relation obtained in eq. (2.64) between the elements of the coefficient matrix $c^{d=6}$ and those of NN^\dagger , it follows that

$$\frac{v^2}{2} |c^{d=6}|_{\alpha\beta} = \frac{v^2}{2} \left| Y_\Sigma^\dagger \frac{1}{M_\Sigma^\dagger} \frac{1}{M_\Sigma} Y_\Sigma \right|_{\alpha\beta} \lesssim \begin{pmatrix} 3 \cdot 10^{-3} & < 1.1 \cdot 10^{-6} & < 1.2 \cdot 10^{-3} \\ < 1.1 \cdot 10^{-6} & 4 \cdot 10^{-3} & < 1.2 \cdot 10^{-3} \\ < 1.2 \cdot 10^{-3} & < 1.2 \cdot 10^{-3} & 4 \cdot 10^{-3} \end{pmatrix}. \quad (4.43)$$

Notice that these bounds are stronger than those obtained in the case of the fermionic singlet Seesaw theory, eq. (4.2). This is due to the fact that now flavour changing processes with charged fermions are allowed already at tree level.

4.3.8 Signals at colliders from fermionic triplets

As for direct production and detection, alike to the case of the generic type-II Seesaw model, the non-zero electroweak charge of the triplet results in gauge production from photon and Z couplings. Only particles with electric charge ± 1 exist in this case, though, and the experimental signals are less clean. Anyway, if light enough, triplet fermions can be produced in forthcoming colliders through Drell-Yan production. In ref. [6, 8], the following channels have been analyzed:

- Σ decays into gauge bosons plus light leptons: $\Sigma^- \rightarrow Zl^-$, $\Sigma^- \rightarrow W^-\nu$, $\Sigma^0 \rightarrow Z\nu$, $\Sigma^0 \rightarrow W^\pm l^\mp$;
- Σ decays into Higgs plus light leptons: $\Sigma^- \rightarrow \phi^0 l^-$, $\Sigma^0 \rightarrow \phi^0 \nu$.

5. Conclusions

While the unique dimension five effective operator is common to all Seesaw models of Majorana neutrinos, dimension six operators discriminate among them. We have determined the latter for the three families of Seesaw models: fermionic singlet (type I), scalar triplet (type II) and fermionic triplet (type III). They should be the low-energy tell-tale of the Seesaw mechanism, for any generic beyond the SM theory whose typical scale is larger than the electroweak scale and which accounts for Majorana neutrino masses. These results have been gathered in table 1.

For fermionic Seesaw theories, the effective operators obtained result in non-unitary leptonic mixing matrices affecting the couplings of leptons to gauge bosons, in very precise patterns. Denoting by N the non-unitary matrix which replaces the usual U_{PMNS} matrix in the charged current, the neutrino- Z and charged lepton- Z currents have now a flavour structure given by

$$\begin{aligned} J_\mu^{3-\nu} &\propto N^\dagger N, & J_\mu^{3-l} &\propto 1, & \text{for singlet-fermion Seesaw,} \\ J_\mu^{3-\nu} &\propto (N^\dagger N)^{-1}, & J_\mu^{3-l} &\propto (NN^\dagger)^2, & \text{for triplet-fermion Seesaw.} \end{aligned}$$

For scalar-triplet Seesaw theories the mixing matrices remain unitary, while the dimension six operators indicate instead correlations between exotic four-fermion couplings and gauge and Higgs potential parameters, as well as with Higgs transitions.

For all families of Seesaw theories, it turns out that the coefficient matrix of the dimension six leptonic operators is of the generic form $|c^{d=6}| = Y^\dagger \frac{1}{M^2} Y$, where Y denote the new Yukawa couplings and M the high scale of the new theory. Irrespective of the value of M , we have set bounds on the Y/M ratios for the three theories, resulting in an overall constraint $|Y| \sim 10^{-1} \frac{M}{TeV}$, with more stringent constraints for specific channels, specially for type II and III theories due the richness of their phenomenology. The specific

results have been collected in eq. (4.2), tables 4 and 5, and eq. (4.43), for the fermionic singlet, scalar triplet and fermionic triplet Seesaw theories, respectively. To achieve them, we took into account the experimental data on many tree-level processes as well as on radiative one-loop processes ($\mu \rightarrow e\gamma$, $\tau \rightarrow \mu\gamma$ and $\tau \rightarrow e\gamma$), all of which we computed in all 3 theories.

Independently of the above, we have also discussed the possible values of M from a theoretical, albeit model independent, point of view. There is no clue at present on whether there is a relationship between the source of $B - L$ violation in nature and the origin of the flavour structure of the SM, which displays Yukawa couplings for charged fermions ranging from $Y \sim 1$ for the top quark to $\sim 10^{-6}$ for the electron. The values of neutrino Yukawa couplings could also be in that same range within Seesaw theories, if the high energy scale lies in the range between the typical Grand Unification scale down to the TeV scale. Indeed, the electroweak hierarchy problem prefers new physics scales closer to the electroweak scale than to the hypothetical Grand Unified scale, if the new physics involves the Higgs field. To illustrate this point in the present context, we have explicitly computed the one-loop contributions to the Higgs mass in the three families of Seesaw theories, showing its quadratic sensitivity to the new scales.

We have then also addressed in this work the question of whether it is possible to simultaneously allow a “low” scale $M \sim \text{TeV}$ and large, order one, Yukawa couplings, without fine-tuning neither any Yukawa coupling nor combinations of them. The answer is positive and guided by symmetry considerations. Indeed, while neutrino masses correspond to the dimension five operator which violates $B - L$, all dimension six operators preserve it. From the point of view of symmetries, it is then a sensible option to expect large effects of the new physics associated to the latter, while the dimension five operator is further suppressed.

A natural ansatz proposed is that, if in the new theory the Majorana character is associated to some tiny parameter μ which heralds the breaking of $B - L$, the dimension five operator coefficient is necessarily proportional to it, $c^{d=5} \sim \frac{\mu}{M^2}$ and thus suppressed. This mechanism and pattern is stable under radiative corrections, as they have to be proportional to μ , which is the parameter responsible for the small breaking of a global symmetry. We call this pattern *direct lepton violation* since the neutrino masses are proportional to the (small) lepton number violating quantity μ , rather than inversely proportionnal to the large lepton number violating heavy field mass.

It turns out that such an ansatz and pattern is already incorporated in the minimal scalar-triplet Seesaw theory (type II). We have also argued that fermionic Seesaw theories at the TeV scale (as for instance the so-called inverse Seesaw mechanism) include it as well and we have explored the corresponding possible textures and realizations. Would this ansatz happen in nature, new beautiful signals may be expected near the present experimental bounds and in accelerators sensitive to the TeV scale, such as the LHC or ILC. The loose bounds we have obtained for the dimension six operator coefficients, above mentioned, show that it is indeed possible to have such strong signals, with $M \sim \text{TeV}$, Yukawa couplings of $\mathcal{O}(10^{-2}-1)$ and no unnatural fine-tunings.

Acknowledgments

We specially thank Stefan Antusch, who participated in the early stages of this work. We also acknowledge illuminating discussions with José Ramón Espinosa, Enrique Fernández-Martínez, Jacobo López-Pavón and Stefano Rigolin. Furthermore, the authors received partial support from CICYT through the project FPA2006-05423, as well as from the Comunidad Autónoma de Madrid through Proyecto HEPHACOS; P-ESP-00346. T.H. thanks the FNRS-FRS for support. A.A and F.B acknowledge the support of the Agence Nationale de la Recherche ANR through the project JC05-43009-NEUPAC.

A. Non-unitarity

We give here in detail the transformations leading to the currents in eqs. (2.16), (2.17) and eqs. (2.58), (2.59), (2.60), corresponding to the singlet and triplet fermionic Seesaws, respectively.

Singlet fermion seesaw. Consider the Lagrangians in eqs. (2.11) and (2.12). We can rotate to the basis in which the mass matrices are diagonal. In this basis, the neutrino light eigenstates are redefined as

$$\nu_i = V_{i\alpha}^{\text{eff}} \nu_{\alpha L} + V_{i\alpha}^{\text{eff}*} \nu_{\alpha L}^c, \quad (\text{A.1})$$

$$(\text{A.2})$$

where V^{eff} are not unitary matrices because of the field rescaling involved. V^{eff} can be expressed in terms of the matrix which diagonalizes the neutrino mass matrix¹⁹ U^ν ,

$$V^{\text{eff}} = \left(1 - \frac{1}{2}\epsilon^N\right) U^\nu. \quad (\text{A.3})$$

$$(\text{A.4})$$

In terms of the light mass eigenstates, the leptonic Lagrangian now becomes

$$\mathcal{L}_{\text{leptons}}^{d \leq 6} = \frac{1}{2} \bar{\nu}_i \left(i \not{\partial} - m_{\nu i}^{\text{diag}} \right) \nu_i + \frac{1}{2} \bar{l}_i \left(i \not{\partial} - m_{l i}^{\text{diag}} \right) l_i + \mathcal{L}_{CC} + \mathcal{L}_{NC}, \quad (\text{A.5})$$

where, in this mass basis, the charged and neutral currents read

$$\mathcal{L}_{CC} = \frac{g}{\sqrt{2}} \bar{l}_L W^- \left[\Omega \left(1 - \frac{1}{2} \epsilon^N \right) U^\nu \right] \nu_L + \text{h.c.}, \quad (\text{A.6})$$

$$\mathcal{L}_{NC} = \frac{g}{\cos \theta_W} \left\{ \frac{1}{2} \left[\bar{\nu}_L \gamma_\mu \left[U^{\nu\dagger} (1 - \epsilon^N) U^\nu \right] \nu_L - \bar{l}_L \gamma_\mu l_L \right] - \sin^2 \theta_W J_\mu^{\text{em}} \right\} Z^\mu + e J_\mu^{\text{em}} A^\mu, \quad (\text{A.7})$$

where $\Omega \equiv \text{diag}(e^{i\omega_1}, e^{i\omega_2}, e^{i\omega_3})$ reabsorbs three unphysical phases in the definition of the charged lepton fields. The above expressions look quite complicated, but the measurable

¹⁹Notice that U^ν does not depend on $c^{d=6}$ at $\mathcal{O}(1/M^2)$.

effects can be expressed in a compact way, denoting by N the non-unitary matrix appearing in the charged current coupling,

$$N \equiv \Omega \left(1 - \frac{1}{2} \epsilon^N \right) U^\nu. \quad (\text{A.8})$$

In terms of the matrix N , the charged and neutral currents read

$$J_\mu^{-CC} \equiv \overline{e}_{L\alpha} \gamma_\mu N_{\alpha i} \nu_i, \quad (\text{A.9})$$

$$J_\mu^{NC} \equiv \frac{1}{2} \overline{\nu}_i \gamma_\mu (N^\dagger N)_{ij} \nu_j, \quad (\text{A.10})$$

with $N_{i\alpha}^\dagger N_{\alpha j} \neq \delta_{ij}$ appearing in the neutral current since N is not unitary.

Triplet fermion seesaw. Consider now the Lagrangians in eqs. (2.55)–(2.56). We can rotate to the basis in which both the leptonic kinetic energies and their mass matrices are diagonalized. This requires to redefine the leptonic light fields as

$$\nu_i = V_{i\alpha}^{\text{eff}} \nu_{\alpha L} + V_{i\alpha}^{\text{eff}*} \nu_{\alpha L}^c, \quad (\text{A.11})$$

$$l_{Li} = K_{i\alpha}^{\text{eff}} l_{L\alpha}, \quad (\text{A.12})$$

$$l_{Ri} = (U_R^l)_{i\alpha} l_{R\alpha}, \quad (\text{A.13})$$

where U_R^l is unitary while V^{eff} and K^{eff} are not unitary matrices because of the field rescaling involved,

$$V^{\text{eff}} = (1 - \frac{1}{2} \epsilon^\Sigma) U^\nu \quad (\text{A.14})$$

$$K^{\text{eff}} = (1 - \epsilon^\Sigma) U_L^l, \quad (\text{A.15})$$

with the matrices U_ν and U^l diagonalizing the neutrino and charged lepton mass terms, respectively,²⁰ with

$$m_\nu^{\text{diag}} \equiv U^{\nu T} m_\nu U^\nu, \quad m_l^{\text{diag}} \equiv U_R^{l\dagger} m_l (1 - \epsilon) U_L^l. \quad (\text{A.16})$$

In terms of the light mass eigenstates, the leptonic Lagrangian becomes

$$\mathcal{L}_{\text{leptons}}^{d \leq 6} = \frac{1}{2} \overline{\nu}_i \left(i \not{\partial} - m_{\nu i}^{\text{diag}} \right) \nu_i + \frac{1}{2} \overline{l}_i \left(i \not{\partial} - m_{li}^{\text{diag}} \right) l_i + \mathcal{L}_{CC} + \mathcal{L}_{NC}, \quad (\text{A.17})$$

in which the charged and neutral currents now are given by

$$\mathcal{L}_{CC} = \frac{g}{\sqrt{2}} \overline{l}_L W^- \left[\Omega U_L^{l\dagger} \left(1 + \frac{1}{2} \epsilon^\Sigma \right) U^\nu \right] \nu_L + \text{h.c.}, \quad (\text{A.18})$$

$$\begin{aligned} \mathcal{L}_{NC} = \frac{g}{\cos \theta_W} \Big\{ & \frac{1}{2} \left[\overline{\nu}_L \gamma_\mu \left[U^{\nu\dagger} (1 - \epsilon^\Sigma) U^\nu \right] \nu_L - \overline{l}_L \gamma_\mu \left[\Omega U_L^{l\dagger} (1 + 2\epsilon^\Sigma) U_L^l \Omega^\dagger \right] l_L \right] \\ & - \sin^2 \theta_W J_\mu^{em} \Big\} Z^\mu + e J_\mu^{em} A^\mu, \end{aligned} \quad (\text{A.19})$$

²⁰While U^ν does not depend on $c^{d=6}$ at $\mathcal{O}(1/M^2)$, U_L^l and U_R^l do.

with, once again, $\Omega \equiv \text{diag}(e^{i\omega_1}, e^{i\omega_2}, e^{i\omega_3})$ reabsorbing three unphysical phases in the definition of the charged lepton fields.

Because of the flavour-dependent field rescaling involved, a non-unitary mixing matrix N has appeared in the charged-current couplings, replacing the usual unitary U_{PMNS} matrix, while non-unitary flavour mixing appears as well in the couplings of leptons to the Z boson. The above expressions look quite cumbersome, but the measurable effects can be cast in a compact way. Indeed, we denote by N the non-unitary matrix appearing in the charged current coupling,

$$N \equiv \Omega U_L^{\dagger} \left(1 + \frac{1}{2} \epsilon^{\Sigma} \right) U^{\nu}. \quad (\text{A.20})$$

Working at order $\mathcal{O}(1/M^2)$, i.e. at first order in the ϵ^{Σ} parameters, the charged and neutral currents can then be neatly expressed in the mass basis as

$$J_{\mu}^{-CC} \equiv \overline{l}_L \gamma_{\mu} N \nu, \quad (\text{A.21})$$

$$J_{\mu}^3(\text{neutrinos}) \equiv \frac{1}{2} \overline{\nu} \gamma_{\mu} (N^{\dagger} N)^{-1} \nu, \quad (\text{A.22})$$

$$J_{\mu}^3(\text{leptons}) \equiv \frac{1}{2} \overline{l} \gamma_{\mu} (N N^{\dagger})^2 l. \quad (\text{A.23})$$

B. Lepton mixing in the full type-III seesaw model

As the type-III model has not been properly presented in an extensive way for what concerns notations, mass matrices, mixing matrices, etc, it is useful to discuss it also in the context of the full theory where the triplets of fermions are not integrated out. This will also allow to establish the precise tree-level connection between the effective and full theories. This model is defined by eq. (2.45) in a vector notation. It can be equivalently rewritten in terms of the usual and compact 2 by 2 notation for triplets (with implicit flavour summation):

$$\mathcal{L} = \text{Tr}[\overline{\Sigma} i \not{D} \Sigma] - \frac{1}{2} \text{Tr}[\overline{\Sigma} M_{\Sigma} \Sigma^c + \overline{\Sigma}^c \Sigma] - \tilde{\phi}^{\dagger} \overline{\Sigma} \sqrt{2} Y_{\Sigma} L - \overline{L} \sqrt{2} Y_{\Sigma}^{\dagger} \Sigma \tilde{\phi} \quad (\text{B.1})$$

with, for each fermionic triplet,

$$\begin{aligned} \Sigma &= \begin{pmatrix} \Sigma^0/\sqrt{2} & \Sigma^+ \\ \Sigma^- & -\Sigma^0/\sqrt{2} \end{pmatrix}, & \Sigma^c &= \begin{pmatrix} \Sigma^{0c}/\sqrt{2} & \Sigma^{-c} \\ \Sigma^{+c} & -\Sigma^{0c}/\sqrt{2} \end{pmatrix}, \\ D_{\mu} &= \not{\partial}_{\mu} - i\sqrt{2}g \begin{pmatrix} W_{\mu}^3/\sqrt{2} & W_{\mu}^+ \\ W_{\mu}^- & -W_{\mu}^3/\sqrt{2} \end{pmatrix}. \end{aligned} \quad (\text{B.2})$$

Either way, eq. (2.45) or eq. (B.1), lead to the same Lagrangian expressed in terms of charge components, as given in eq. (4.35) or, in terms of the more convenient spinor field Ψ (eq. (4.36)), in eq. (4.37). This leads to the mass matrices for both neutral and charged leptons in eqs. (4.38)–(4.39). As it happens with any Dirac mass matrix, the charged lepton

mass matrix can be diagonalized by a bi-unitary matrix transformation (6 by 6 if there are 3 triplets of fermions),

$$\begin{pmatrix} l_{L,R} \\ \Psi_{L,R} \end{pmatrix} = U_{L,R} \begin{pmatrix} l'_{L,R} \\ \Psi'_{L,R} \end{pmatrix}, \quad (\text{B.3})$$

while the symmetric neutral lepton mass matrix can be diagonalized by a single unitary matrix:

$$\begin{pmatrix} \nu_L \\ \Sigma^{0c} \end{pmatrix} = U_0 \begin{pmatrix} \nu'_L \\ \Sigma'^{0c} \end{pmatrix}. \quad (\text{B.4})$$

Writing the mixing matrices in terms of 3 by 3 blocks

$$U_L = \begin{pmatrix} U_{LL} & U_{L\Sigma} \\ U_{L\Sigma l} & U_{L\Sigma\Sigma} \end{pmatrix}, \quad U_R = \begin{pmatrix} U_{Rl} & U_{R\Sigma} \\ U_{R\Sigma l} & U_{R\Sigma\Sigma} \end{pmatrix}, \quad U_0 = \begin{pmatrix} U_{0\nu\nu} & U_{0\nu\Sigma} \\ U_{0\Sigma\nu} & U_{0\Sigma\Sigma} \end{pmatrix}, \quad (\text{B.5})$$

at first order in Yv/M_Σ we obtain:

$$U_{L\Sigma} = Y^\dagger v M_\Sigma^{-1}, \quad U_{L\Sigma l} = -M_\Sigma^{-1} Y v, \quad U_{0\nu\Sigma} = \frac{Y^\dagger}{\sqrt{2}} v M^{-1}, \quad U_{0\Sigma\nu} = -M_\Sigma^{-1} \frac{Y}{\sqrt{2}} v U_{PMNS}, \quad (\text{B.6})$$

while $U_{Ll} = U_{L\Sigma\Sigma} = U_{0\Sigma\Sigma} = 1$, $U_{0\nu\nu} = U_{PMNS}$, $U_R = 1$.²¹ In these equations U_{PMNS} is the lowest order neutrino mass matrix, which is unitary. Note that we have neglected smaller corrections in Yvm_l/M_Σ^2 . The 6 by 6 mixing matrices $U_{L,R,0}$ are unitary but the various 3 by 3 ones are not. This leads to non-unitary effects in the gauge interactions of leptons. Re-expressing the gauge interactions in the mass eigenstate basis we get eqs. (2.58)–(2.60) with

$$(NN^\dagger)^2 = 1 + U_{L\Sigma l}^\dagger U_{L\Sigma l} = 1 + 2\epsilon^\Sigma, \quad (\text{B.7})$$

$$(N^\dagger N)^{-1} = 1 - U_{0\Sigma\nu}^\dagger U_{0\Sigma\nu} = 1 - U_{PMNS}^\dagger \epsilon^\Sigma U_{PMNS}, \quad (\text{B.8})$$

$$N = (U_{Ll}^\dagger U_{0\nu\nu} + \sqrt{2} U_{L\Sigma l}^\dagger U_{0\Sigma\nu}) = \left(1 + \frac{\epsilon^\Sigma}{2}\right) U_{PMNS}. \quad (\text{B.9})$$

In obtaining these results recall that, in the full high-energy theory, all the analysis has been performed in the flavour basis in which the initial charged lepton mass matrix is diagonal and the light charged lepton fields have reabsorbed three arbitrary phases. In the last equalities of eqs. (B.7) and (B.8) we have used eq. (B.6), while the last equality of eq. (B.9) can be obtained from combining eqs. (B.7) and (B.8). The results we get in terms of the Yukawa couplings are fully in agreement with the ones obtained in the effective theory, eqs. (2.58)–(2.62) and eq. (2.64). Note that in eqs. (B.6)–(B.8) (although not in eq. (B.9)), U_{PMNS} can be replaced by N since the difference is of higher order in Yv/M_Σ .

²¹It should be remarked that, in the one-loop computations of the radiative processes $l_1 \rightarrow l_2 \gamma$, it has been necessary to consider the expansion of U_R at $\mathcal{O}(\frac{1}{M_\Sigma^2})$: $U_{Rl\Sigma} = m_l Y^\dagger v M_\Sigma^{-2}$, $U_{R\Sigma l} = -M_\Sigma^{-2} Y m_l v$.

C. Low scale models of light neutrino masses with large Yukawa couplings

In the following, we consider models based on type-I Seesaw mechanism which lead to large dimension 6 operators. The examples considered can be straightforwardly applied to the type-III Seesaw too, as the textures are exactly the same. Such a situation arises for particular patterns of the singlet neutrino mass matrix and/or of the Yukawa matrix. As already explained in section 3 we are interested in a class of models which, to lead to sufficiently suppressed neutrino masses and large $d = 6$ operators, do not require any precise cancellations between the various (a priori independent) entries of these 2 matrices. The cases we consider just require that some of the entries of these mass matrices are much smaller than other ones.

For simplicity, let us first consider — as in section 3 — only one left-handed neutrino and two singlet fermions. In full generality, in this case there are 3 mass matrix textures which automatically lead to a vanishing light neutrino mass [in the basis (ν_L, N_1, N_2)]:

$$\begin{pmatrix} 0 & m_{D_1} & 0 \\ m_{D_1} & M_{N_2} & M_{N_1} \\ 0 & M_{N_1} & 0 \end{pmatrix}, \quad \begin{pmatrix} 0 & 0 & m_{D_1} \\ 0 & 0 & M_{N_1} \\ m_{D_1} & M_{N_1} & M_{N_2} \end{pmatrix}, \quad \begin{pmatrix} 0 & m_{D_1} & m_{D_2} \\ m_{D_1} & 0 & 0 \\ m_{D_2} & 0 & 0 \end{pmatrix} \quad (\text{C.1})$$

In the following we will consider only the first mass matrix since the second one is equivalent to the first one under $N_1 \leftrightarrow N_2$, and since the third one which is of the Dirac type doesn't lead to any interesting case for our purposes. Assuming $M_{N_1} > m_{D_1}$ the eigenstate which is predominantly a ν_L is massless. For $M_{N_2} = 0$, which corresponds to the well-known inverse Seesaw model [19] considered in section 3, this can be understood easily from the fact that assigning $L = 1, -1, 1$ to ν_L, N_1, N_2 respectively, lepton number is conserved. For $M_{N_2} \neq 0$, which can also be justified from a symmetry in specific extended models [20], this remains true because the determinant of the mass matrix still vanishes in this case. This case has the interesting feature to have a large source of lepton number violation (i.e. M_{N_2}) with a vanishing neutrino mass.²²

In order to induce a naturally small neutrino mass, even if the Yukawa coupling in m_{D_1} is large, and without fine-tuning, one must introduce a small mass parameter μ in the mass matrix. This can be done in 2 ways (plus combination of them), either from introducing an extra small Majorana mass, or from introducing an extra small Dirac mass term:

$$\begin{pmatrix} 0 & m_{D_1} & 0 \\ m_{D_1} & M_{N_2} & M_{N_1} \\ 0 & M_{N_1} & \mu \end{pmatrix}, \quad \begin{pmatrix} 0 & m_{D_1} & \mu \\ m_{D_1} & M_{N_2} & M_{N_1} \\ \mu & M_{N_1} & 0 \end{pmatrix}. \quad (\text{C.2})$$

Expanding in powers of μ , in the first case in eq. (C.2) we obtain:

$$m_\nu = \frac{m_{D_1}^2}{M_{N_1}} \frac{\mu}{M_{N_1}} \frac{M_{N_1}^2}{M_{N_1}^2 + m_{D_1}^2} + \mathcal{O}(m_{D_1}^2 \mu^2 M_{N_2} / M_{N_1}^4, \mu^3), \quad (\text{C.3})$$

²²Note that since the 22 element breaks lepton number, it could induce neutrino mass in presence of extra interactions coupling to the N_i 's. This contribution would be suppressed by loop factors, couplings of the extra interactions, as well as the masses of the new states involved, but wouldn't be necessarily negligible with respect to the contribution of eq. (C.3) below. We thank S. Antusch, M. Frigerio and J. Kersten for discussions on this point.

while the second case leads to:

$$m_\nu = -2 \frac{m_{D_1} \mu}{M_{N_1}} \frac{M_{N_1}^2}{M_{N_1}^2 + m_{D_1}^2} + \frac{\mu^2}{M_{N_1}} \frac{M_{N_2}}{M_{N_1}} \frac{(M_{N_1}^2 - m_{D_1}^2)^2}{(M_{N_1}^2 + m_{D_1}^2)^2} + \mathcal{O}(\mu^3). \quad (\text{C.4})$$

Eq. (C.3) shows that the neutrino mass is suppressed by an extra factor μ/M_{N_1} , so that the smallness of neutrino masses, and the argument of no fine tuning, do not require tiny Yukawa couplings.

As for the first term in eq. (C.4), it has the standard neutrino mass form, i.e. with 2 Dirac masses in the numerator and one Majorana mass in the denominator, but unlike the usual Seesaw formula, it involves only the product of 2 different Dirac masses. Therefore, if one of them is smaller than the other, e.g. $\mu \ll m_{D_1}$, a small neutrino mass can be obtained here too with a large Yukawa coupling in m_{D_1} , and no fine-tuning. As for the second term in eq. (C.4), which involves the independent parameter M_{N_2} , it also leads to suppressed neutrino masses, even if M_{N_2} largely breaks lepton number.

Now, in the limit $\mu \rightarrow 0$ the point is that the coefficient of the $d = 5$ operator vanishes but that of the $d = 6$ operator does not. This can be seen from the fact that the $d = 6$ operator takes the form $(Y_N)^\dagger (M_N^{-2}) (Y_N)$, see above, and doesn't vanish in this limit. Eq. (2.7) in all cases above, with for example $m_{D_1} = Y_1 v \sim v$ and $M_{N_1} \sim 1 \text{ TeV}$, becomes simply $|Y_1|^2/M_{N_1}^2 \sim 1/M_{N_1}^2$ which is large. The one left-handed plus two right-handed neutrino example above can be generalized to the 3 left-handed plus 3 right-handed neutrino above. The condition for having vanishing neutrino masses is to start with a 6 by 6 mass matrix which has rank 3. Assuming that all entries of the Yukawa coupling matrix are independent (i.e. barring cancellations between the various entries), it turns out that there is only one possibility to have large Yukawa couplings with three massless light neutrinos and three massive right-handed neutrinos. In the basis $(\nu_e, \nu_\mu, \nu_\tau, N_1, N_2, N_3)$ it is

$$\begin{pmatrix} 0 & 0 & 0 & c & 0 & 0 \\ 0 & 0 & 0 & d & 0 & 0 \\ 0 & 0 & 0 & e & 0 & 0 \\ c & d & e & f & g & a \\ 0 & 0 & 0 & g & b & 0 \\ 0 & 0 & 0 & a & 0 & 0 \end{pmatrix}, \quad (\text{C.5})$$

plus permutations. This matrix has the particularity that only one of the 3 right-handed neutrinos couples to light neutrinos at leading order (just as the 1 ν plus 2 N case above). From a simple lepton number assignment there is only one way to justify this pattern, which gives in addition $f = g = 0$, i.e. by taking $L_{\nu_e} = L_{\nu_\mu} = L_{\nu_\tau} = L_{N_1} = -L_{N_3} = 1$ and

$L_{N_2} = 0$.^{23 24}. The matrix of eq. (C.5) can be perturbed in many ways:

$$\begin{pmatrix} 0 & 0 & 0 & c & \varepsilon_1 & \varepsilon_2 \\ 0 & 0 & 0 & d & \varepsilon_3 & \varepsilon_4 \\ 0 & 0 & 0 & e & \varepsilon_5 & \varepsilon_6 \\ c & d & e & f & g & a \\ \varepsilon_1 & \varepsilon_3 & \varepsilon_5 & g & b & \varepsilon_7 \\ \varepsilon_2 & \varepsilon_4 & \varepsilon_6 & a & \varepsilon_7 & \varepsilon_8 \end{pmatrix}, \quad (\text{C.6})$$

To have two massless light neutrinos, at least one ε_i among $\varepsilon_{1,\dots,7}$ must be different from 0. To have 3 massless light neutrinos, at least two well chosen ε_i must be different from 0, for example ε_3 and ε_6 . It is beyond the scope of the present analysis to determine all possible perturbations textures which may accommodate the neutrino data along these lines, see also ref. [50]. There are many possibilities, and the point is that all of them do lead to unsuppressed $d = 6$ operators (i.e. with non-vanishing coefficients in the limit in which all $\varepsilon_i = 0$) as long as a and b , together with at least one parameter among c, d, e , are different from 0.

References

- [1] S. Weinberg, *Baryon and lepton nonconserving processes*, *Phys. Rev. Lett.* **43** (1979) 1566.
- [2] P. Minkowski, $\mu \rightarrow e\gamma$ at a rate of one out of 1-billion muon decays?, *Phys. Lett.* **B 67** (1977) 421;
M. Gell-Mann, P. Ramond and R. Slansky, in *Supergravity*, edited by P. van Nieuwenhuizen and D. Freedman, North-Holland (1979);
T. Yanagida, in *Proceedings of the Workshop on the unified theory and the baryon number in the universe*, edited by O. Sawada and A. Sugamoto, Tsukuba (1979);
R.N. Mohapatra and G. Senjanovic, *Neutrino mass and spontaneous parity nonconservation*, *Phys. Rev. Lett.* **44** (1980) 912.
- [3] W. Buchmuller and D. Wyler, *Effective lagrangian analysis of new interactions and flavor conservation*, *Nucl. Phys.* **B 268** (1986) 621.
- [4] M. Magg and C. Wetterich, *Neutrino mass problem and gauge hierarchy*, *Phys. Lett.* **B 94** (1980) 61;
J. Schechter and J.W.F. Valle, *Neutrino masses in $SU(2) \times U(1)$ theories*, *Phys. Rev.* **D 22** (1980) 2227;
C. Wetterich, *Neutrino masses and the scale of B-L violation*, *Nucl. Phys.* **B 187** (1981) 343;
G. Lazarides, Q. Shafi and C. Wetterich, *Proton lifetime and fermion masses in an $SO(10)$ model*, *Nucl. Phys.* **B 181** (1981) 287;
R.N. Mohapatra and G. Senjanovic, *Neutrino masses and mixings in gauge models with spontaneous parity violation*, *Phys. Rev.* **D 23** (1981) 165.

²³For completeness, it can be noted that the 3 light ν s plus 2 heavy N case also leads to a unique possible texture. It corresponds to take no N_3 , i.e. $a = 0$, and requires to take $b = 0$ in addition. It can be justified from a L assignment if moreover $f = 0$ with $L = 1$ for all particles except N_2 which has $L = -1$.

²⁴During the completion of this work, ref. [50] appeared, which also considers this particular texture.

- [5] R. Foot, H. Lew, X.G. He and G.C. Joshi, *Seesaw neutrino masses induced by a triplet of leptons*, *Z. Physik C* **44** (1989) 441;
E. Ma, *Pathways to naturally small neutrino masses*, *Phys. Rev. Lett.* **81** (1998) 1171 [[hep-ph/9805219](#)].
- [6] E. Ma and D.P. Roy, *Heavy triplet leptons and new gauge boson*, *Nucl. Phys. B* **644** (2002) 290 [[hep-ph/0206150](#)].
- [7] T. Hambye, Y. Lin, A. Notari, M. Papucci and A. Strumia, *Constraints on neutrino masses from leptogenesis models*, *Nucl. Phys. B* **695** (2004) 169 [[hep-ph/0312203](#)].
- [8] B. Bajc and G. Senjanovic, *Seesaw at LHC*, *JHEP* **08** (2007) 014 [[hep-ph/0612029](#)];
B. Bajc, M. Nemevsek and G. Senjanovic, *Probing seesaw at LHC*, *Phys. Rev. D* **76** (2007) 055011 [[hep-ph/0703080](#)].
- [9] I. Dorsner and P. Fileviez Perez, *Upper bound on the mass of the type-III seesaw triplet in an SU(5) model*, *JHEP* **06** (2007) 029 [[hep-ph/0612216](#)];
P.F. Perez, *Renormalizable adjoint SU(5)*, *Phys. Lett. B* **654** (2007) 189 [[hep-ph/0702287](#)];
Supersymmetric adjoint SU(5), *Phys. Rev. D* **76** (2007) 071701 [[arXiv:0705.3589](#)].
- [10] S. Antusch, C. Biggio, E. Fernandez-Martinez, M.B. Gavela and J. Lopez-Pavon, *Unitarity of the leptonic mixing matrix*, *JHEP* **10** (2006) 084 [[hep-ph/0607020](#)].
- [11] A. Broncano, M.B. Gavela and E.E. Jenkins, *The effective lagrangian for the seesaw model of neutrino mass and leptogenesis*, *Phys. Lett. B* **552** (2003) 177 [*Erratum ibid.* **636** (2006) 330] [[hep-ph/0210271](#)]; *Neutrino physics in the seesaw model*, *Nucl. Phys. B* **672** (2003) 163 [[hep-ph/0307058](#)].
- [12] W. Chao and H. Zhang, *One-loop renormalization group equations of the neutrino mass matrix in the triplet seesaw model*, *Phys. Rev. D* **75** (2007) 033003 [[hep-ph/0611323](#)].
- [13] A. De Rujula, M.B. Gavela, P. Hernandez and E. Masso, *The selfcouplings of vector bosons: does LEP-1 obviate LEP-2?*, *Nucl. Phys. B* **384** (1992) 3.
- [14] PARTICLE DATA GROUP collaboration, W.M. Yao et al., *Review of particle physics*, *J. Phys. G* **33** (2006) 1 (and 2007 partial update for edition 2008).
- [15] A. Abada, F. Bonnet, C. Biggio, M.B. Gavela and T. Hambye, in preparation.
- [16] J.A. Casas, J.R. Espinosa and I. Hidalgo, *Implications for new physics from fine-tuning arguments. I: application to SUSY and seesaw cases*, *JHEP* **11** (2004) 057 [[hep-ph/0410298](#)].
- [17] V. Cirigliano, B. Grinstein, G. Isidori and M.B. Wise, *Minimal flavor violation in the lepton sector*, *Nucl. Phys. B* **728** (2005) 121 [[hep-ph/0507001](#)].
- [18] A. Pilaftsis, *Resonant tau leptogenesis with observable lepton number violation*, *Phys. Rev. Lett.* **95** (2005) 081602 [[hep-ph/0408103](#)].
- [19] M.C. Gonzalez-Garcia and J.W.F. Valle, *Fast decaying neutrinos and observable flavor violation in a new class of Majoron models*, *Phys. Lett. B* **216** (1989) 360.
- [20] E. Dudas and C.A. Savoy, *Multiple seesaw at low energy*, *Acta Phys. Polon. B* **33** (2002) 2547 [[hep-ph/0205264](#)].
- [21] A. de Gouvêa, G.F. Giudice, A. Strumia and K. Tobe, *Phenomenological implications of neutrinos in extra dimensions*, *Nucl. Phys. B* **623** (2002) 395 [[hep-ph/0107156](#)].

- [22] A. Czarnecki and E. Jankowski, *Electromagnetic suppression of the decay $\mu \rightarrow e\gamma$* , *Phys. Rev. D* **65** (2002) 113004 [[hep-ph/0106237](#)].
- [23] E. Fernandez-Martinez, M.B. Gavela, J. Lopez-Pavon and O. Yasuda, *CP-violation from non-unitary leptonic mixing*, *Phys. Lett. B* **649** (2007) 427 [[hep-ph/0703098](#)].
- [24] F. del Aguila, J.A. Aguilar-Saavedra and R. Pittau, *Neutrino physics at large colliders*, *J. Phys. Conf. Ser.* **53** (2006) 506 [[hep-ph/0606198](#)].
- [25] J.F. Gunion, H.E. Haber, G.L. Kane and S. Dawson, *The Higgs hunter's guide*, Westview Press (2000).
- [26] H.M. Georgi, S.L. Glashow and S. Nussinov, *Unconventional model of neutrino masses*, *Nucl. Phys. B* **193** (1981) 279.
- [27] V.D. Barger, H. Baer, W.Y. Keung and R.J.N. Phillips, *Decays of weak vector bosons and T quarks into doubly charged Higgs scalars*, *Phys. Rev. D* **26** (1982) 218.
- [28] P.B. Pal, *Constraints on a muon neutrino mass around 100 KeV*, *Nucl. Phys. B* **227** (1983) 237.
- [29] A. Pich, A. Santamaria and J. Bernabeu, *$\mu^- \rightarrow e^- + \gamma$ decay in the scalar triplet model*, *Phys. Lett. B* **148** (1984) 229.
- [30] J. Bernabeu, A. Pich and A. Santamaria, *CP phases in the charged current and Higgs sectors for majorana neutrinos*, *Z. Physik C* **30** (1986) 213.
- [31] S.M. Bilenky and S.T. Petcov, *Massive neutrinos and neutrino oscillations*, *Rev. Mod. Phys.* **59** (1987) 671.
- [32] J.F. Gunion, J. Grifols, A. Mendez, B. Kayser and F.I. Olness, *Higgs bosons in left-right symmetric models*, *Phys. Rev. D* **40** (1989) 1546.
- [33] M.L. Swartz, *Limits on doubly charged Higgs bosons and lepton flavor violation*, *Phys. Rev. D* **40** (1989) 1521.
- [34] R.N. Mohapatra, *Rare decays of the tau lepton as a probe of the left-right symmetric theories of weak interactions*, *Phys. Rev. D* **46** (1992) 2990.
- [35] E. Accomando et al., *Workshop on CP studies and non-standard Higgs physics*, [hep-ph/0608079](#).
- [36] R.N. Cahn and H. Harari, *Bounds on the masses of neutral generation changing gauge bosons*, *Nucl. Phys. B* **176** (1980) 135.
- [37] C.S. Lim and T. Inami, *Lepton flavor nonconservation and the mass generation mechanism for neutrinos*, *Prog. Theor. Phys.* **67** (1982) 1569.
- [38] A.G. Akeroyd, M. Aoki and Y. Okada, *Lepton flavour violating tau decays in the left-right symmetric model*, *Phys. Rev. D* **76** (2007) 013004 [[hep-ph/0610344](#)].
- [39] V. Cirigliano and B. Grinstein, *Phenomenology of minimal lepton flavor violation*, *Nucl. Phys. B* **752** (2006) 18 [[hep-ph/0601111](#)].
- [40] MEG collaboration, S. Ritt, *Status of the meg experiment $\mu \rightarrow e \gamma$* , *Nucl. Phys.* **162** (Proc. Suppl.) (2006) 279.
- [41] A.G. Akeroyd and M. Aoki, *Single and pair production of doubly charged Higgs bosons at hadron colliders*, *Phys. Rev. D* **72** (2005) 035011 [[hep-ph/0506176](#)].

- [42] E. Accomando et al., *Workshop on CP studies and non-standard Higgs physics*, [hep-ph/0608079](#).
- [43] CDF collaboration, D. Acosta et al., *Search for doubly-charged Higgs bosons decaying to dileptons in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV*, *Phys. Rev. Lett.* **93** (2004) 221802 [[hep-ex/0406073](#)].
- [44] K. Huitu, J. Maalampi, A. Pietila and M. Raidal, *Doubly charged Higgs at LHC*, *Nucl. Phys.* **B 487** (1997) 27 [[hep-ph/9606311](#)];
E. Ma, M. Raidal and U. Sarkar, *Phenomenology of the neutrino-mass-giving Higgs triplet and the low-energy seesaw violation of lepton number*, *Nucl. Phys.* **B 615** (2001) 313 [[hep-ph/0012101](#)].
- [45] A. Hektor, M. Kadastik, M. Muntel, M. Raidal and L. Rebane, *Testing neutrino masses in little Higgs models via discovery of doubly charged Higgs at LHC*, [arXiv:0705.1495](#).
- [46] E.J. Chun, K.Y. Lee and S.C. Park, *Testing Higgs triplet model and neutrino mass patterns*, *Phys. Lett.* **B 566** (2003) 142 [[hep-ph/0304069](#)].
- [47] V. Barger, T. Han, P. Langacker, B. McElrath and P. Zerwas, *Effects of genuine dimension-six Higgs operators*, *Phys. Rev.* **D 67** (2003) 115001 [[hep-ph/0301097](#)].
- [48] S. Kanemura, D. Nomura and K. Tsumura, *Dimension-six top Higgs interaction and its effect in collider phenomenology*, *Phys. Rev.* **D 74** (2006) 076007 [[hep-ph/0607333](#)].
- [49] A. Pich, *The standard model of electroweak interactions*, [hep-ph/0502010](#).
- [50] J. Kersten and A.Y. Smirnov, *Right-handed neutrinos at LHC and the mechanism of neutrino mass generation*, *Phys. Rev.* **D 76** (2007) 073005 [[arXiv:0705.3221](#)].

Erratum

The correct address of the Instituto de Física Teórica is:

*Instituto de Física Teórica UAM-CSIC, Facultad de Ciencias C-XVI,
Universidad Autónoma de Madrid, Cantoblanco, Madrid 28049, Spain*