

# A precision constraint on multi-Higgs-doublet models

W Grimus<sup>1</sup>, L Lavoura<sup>2</sup>, O M Ogreid<sup>3</sup> and P Osland<sup>4</sup>

<sup>1</sup> Fakultät für Physik, Universität Wien, Boltzmannngasse 5, 1090 Wien, Austria

<sup>2</sup> Universidade Técnica de Lisboa and Centro de Física Teórica de Partículas, Instituto Superior Técnico, 1049-001 Lisboa, Portugal

<sup>3</sup> Bergen University College, Bergen, Norway

<sup>4</sup> Department of Physics and Technology, University of Bergen, Postboks 7803, N-5020 Bergen, Norway

E-mail: [walter.grimus@univie.ac.at](mailto:walter.grimus@univie.ac.at), [balio@cftp.ist.utl.pt](mailto:balio@cftp.ist.utl.pt), [omo@hib.no](mailto:omo@hib.no) and [per.osland@ift.uib.no](mailto:per.osland@ift.uib.no)

Received 15 February 2008

Published 8 May 2008

Online at [stacks.iop.org/JPhysG/35/075001](http://stacks.iop.org/JPhysG/35/075001)

## Abstract

We derive a general expression for  $\Delta\rho$  (or, equivalently, for the oblique parameter  $T$ ) in the  $SU(2) \times U(1)$  electroweak model with an arbitrary number of scalar  $SU(2)$  doublets, with hypercharge  $\pm 1/2$ , and an arbitrary number of scalar  $SU(2)$  singlets. The experimental bound on  $\Delta\rho$  constitutes a strong constraint on the masses and mixings of the scalar particles in that model.

## 1. Introduction

In the Standard Model (SM), the parameter

$$\rho = \frac{m_W^2}{m_Z^2 \cos^2 \theta_W}, \quad (1)$$

where  $m_W$  and  $m_Z$  are the masses of the  $W^\pm$  and  $Z^0$  gauge bosons, respectively, and  $\theta_W$  is the weak mixing angle, gives the relative strength of the neutral-current and charged-current interactions in four-fermion processes at zero momentum transfer [1]. At tree level  $\rho$  is equal to one, and it remains one even if additional scalar  $SU(2)$  doublets, with hypercharge  $\pm 1/2$ , are added to the SM<sup>5</sup>. At the one-loop level, the vacuum-polarization effects, which are sensitive to any field that couples either to the  $W^\pm$  or to the  $Z^0$ , produce the vacuum-polarization tensors ( $V = W, Z$ )

$$\Pi_{VV}^{\mu\nu}(q) = g^{\mu\nu} A_{VV}(q^2) + q^\mu q^\nu B_{VV}(q^2), \quad (2)$$

where  $q^\mu$  is the 4-momentum of the gauge boson. Then, deviations of  $\rho$  from unity arise, which are determined by the self-energy difference [1, 2]

$$\frac{A_{WW}(0)}{m_W^2} - \frac{A_{ZZ}(0)}{m_Z^2}. \quad (3)$$

<sup>5</sup> Other scalar  $SU(2) \times U(1)$  representations are also allowed, as long as they have vanishing vacuum expectation values.

The precise measurement [3], at LEP, of the  $W^\pm$  and  $Z^0$  self-energies is in striking agreement with the SM predictions [4] and provides a strong constraint on extended electroweak models. For instance, one can constrain the two-Higgs-doublet model (2HDM) in this way [5, 6].

In this paper we are interested in the contributions to the  $\rho$  parameter generated by an extension of the SM. Therefore, we define a  $\Delta\rho$  which refers to the non-SM part of the quantity (3):

$$\Delta\rho = \left[ \frac{A_{WW}(0)}{m_W^2} - \frac{A_{ZZ}(0)}{m_Z^2} \right]_{\text{SM extension}} - \left[ \frac{A_{WW}(0)}{m_W^2} - \frac{A_{ZZ}(0)}{m_Z^2} \right]_{\text{SM}}. \quad (4)$$

The SM contributions to the quantity (3) are known up to the leading terms at the three-loop level [7]. However, the consistent SM subtraction in equation (4) only requires the one-loop SM result. In the same vein, we are allowed to make the replacement  $m_Z^2 = m_W^2/c_W^2$  in equation (4), writing instead

$$\Delta\rho = \left[ \frac{A_{WW}(0) - c_W^2 A_{ZZ}(0)}{m_W^2} \right]_{\text{SM extension}} - \left[ \frac{A_{WW}(0) - c_W^2 A_{ZZ}(0)}{m_W^2} \right]_{\text{SM}}. \quad (5)$$

Here and in the following, we use the abbreviations  $c_W = \cos \theta_W$ ,  $s_W = \sin \theta_W$ .

At one loop, the contributions of new physics to the self-energies constitute intrinsically divergent Feynman diagrams, but the divergent parts cancel out among different diagrams, between  $A_{WW}(0)$  and  $c_W^2 A_{ZZ}(0)$ , and also, eventually, through the subtraction of the SM contributions laid out in equation (5). If the new-physics model is renormalizable, then  $\Delta\rho$  is finite. The cancellations finally leave either a quadratic or a logarithmic dependence of  $\Delta\rho$  on the masses of the new-physics particles. The pronounced effects of large masses are what renders the parameter  $\Delta\rho$  so interesting for probing physics beyond the Standard Model.

The functions  $A_{VV}(q^2)$  contain more information about new physics than that just provided by  $\Delta\rho$ . In fact, for new physics much above the electroweak scale, a detailed analysis of the so-called oblique corrections leads to the identification of three relevant observables, which were called  $S$ ,  $T$  and  $U$  in [8] and  $\epsilon_1$ ,  $\epsilon_2$  and  $\epsilon_3$  in [9]<sup>6</sup>. While these two sets of observables differ in their precise definitions, the quantity of interest in this paper is simply

$$\Delta\rho = \alpha T = \epsilon_1, \quad (6)$$

where  $\alpha = e^2/(4\pi) = g^2 s_W^2/(4\pi)$  is the fine-structure constant.

It is not straightforward to obtain a bound on  $\Delta\rho$  from electroweak precision data. One possibility is to add the oblique parameters to the SM parameter set and perform fits to the data. However, since the SM Higgs-boson loops themselves resemble oblique effects, one cannot determine the SM Higgs-boson mass  $m_h$  simultaneously with  $S$  and  $T$  [4]. To get a feeling for the order of magnitude allowed for  $\Delta\rho$ , we quote the number

$$T = -0.03 \pm 0.09(+0.09), \quad (7)$$

which was obtained in [4] by fixing  $U = 0$ . For the mean value of  $T$ , the Higgs-boson mass  $m_h = 117$  GeV was assumed; the mean value in parentheses is for  $m_h = 300$  GeV. Equation (7) translates into  $\Delta\rho = -0.0002 \pm 0.0007(+0.0007)$ .

There is a vast literature on the 2HDM: see [11] for a review, [12] for the renormalization of the model, [13, 14] for the possibility of having a light pseudoscalar compatible with all experimental constraints, and [15, 16], and the references therein for other various recent works. However, just as the 2HDM may differ significantly from the SM, a general multi-Higgs-doublet model may be quite different from its minimal version with only two Higgs

<sup>6</sup> For new physics at a mass scale comparable to the electroweak scale three more such ‘oblique parameters’ have been identified in [10].

doublets [17]. Three or more Higgs doublets frequently appear in models with family symmetries through which one wants to explain various features of the fermion masses and mixings; for some examples in the lepton sector see the reviews in [18].

In this paper, we present a calculation of  $\Delta\rho$  in an extension of the SM with an arbitrary number of Higgs doublets and also, in addition, arbitrary numbers of neutral and charged scalar  $SU(2)$  singlets. Our results can be used to check the compatibility of the scalar sector of multi-Higgs models with the constraints resulting from the electroweak precision experiments.

Recently, there has been some interest in ‘dark’ scalars [19, 20]. These are scalars that have no Yukawa couplings, and are thus decoupled from ordinary matter. Furthermore, they have no vacuum expectation values (VEVs) and therefore display truncated couplings to the gauge bosons. However, they would have quadrilinear vector–vector–scalar–scalar and trilinear vector–scalar–scalar (but no vector–vector–scalar) couplings, and would thus also contribute to, and be constrained by,  $\Delta\rho$ .

The plan of the paper is as follows. In section 2 we present a description of our extension of the SM and the final result of the calculation of  $\Delta\rho$ ; this section is self-consistent and the result can be used without the need to consult the rest of the paper. The details of the calculation are laid out in section 3. The application of our  $\Delta\rho$  formula to the general 2HDM is given in section 4. The summary of our study is found in section 5.

## 2. The model and the result for $\Delta\rho$

### 2.1. The model

We consider an  $SU(2) \times U(1)$  electroweak model in which the scalar sector includes  $n_d$   $SU(2)$  doublets with hypercharge  $1/2$ <sup>7</sup>,

$$\phi_k = \begin{pmatrix} \varphi_k^+ \\ \varphi_k^0 \end{pmatrix}, \quad k = 1, 2, \dots, n_d. \quad (8)$$

Moreover, we allow the model to include an arbitrary number and variety of  $SU(2)$ -singlet scalars; in particular,  $n_c$  complex  $SU(2)$  singlets with hypercharge 1,

$$\chi_j^+, \quad j = 1, 2, \dots, n_c \quad (9)$$

and  $n_n$  real  $SU(2)$  singlets with hypercharge 0,

$$\chi_l^0, \quad l = 1, 2, \dots, n_n. \quad (10)$$

In general, our model may include other scalar fields, singlet under the gauge  $SU(2)$ , with different electric charges.

The neutral fields are allowed to have vacuum expectation values (VEVs). Thus,

$$\langle 0 | \varphi_k^0 | 0 \rangle = \frac{v_k}{\sqrt{2}}, \quad (11)$$

$$\langle 0 | \chi_l^0 | 0 \rangle = u_l, \quad (12)$$

the  $v_k$  being, in general, complex. (The  $u_l$  are real since the  $\chi_l^0$  are real fields.) We define as usual  $v = (\sum_{k=1}^{n_d} |v_k|^2)^{1/2} \simeq 246 \text{ GeV}$ . Then, the masses of the  $W^\pm$  and  $Z^0$  gauge bosons

<sup>7</sup> Equivalently, we may consider the model to contain  $SU(2)$  doublets with hypercharge  $-1/2$ , since

$$\tilde{\phi}_k \equiv i\tau_2 \phi_k^* = \begin{pmatrix} \varphi_k^{0*} \\ -\varphi_k^- \end{pmatrix}$$

is also a doublet of  $SU(2)$ .

are, at tree level,  $m_W = gv/2$  and  $m_Z = m_W/c_W$ , respectively<sup>8</sup>. We expand the neutral fields around their VEVs,

$$\varphi_k^0 = \frac{1}{\sqrt{2}} (v_k + \varphi_k^{0'}), \quad (13)$$

$$\chi_l^0 = u_l + \chi_l^{0'}. \quad (14)$$

Altogether, there are  $n = n_d + n_c$  complex scalar fields with electric charge 1 and  $m = 2n_d + n_n$  real scalar fields with electric charge 0. The mass matrices of all these scalar fields will in general lead to their mixing. The physical (mass-eigenstate) charged and neutral scalar fields will be called  $S_a^+$  ( $a = 1, 2, \dots, n$ ) and  $S_b^0$  ( $b = 1, 2, \dots, m$ ), respectively. Note that the fields  $S_b^0$  are real. We use  $m_a$  to denote the mass of  $S_a^\pm$  and  $\mu_b$  to denote the mass of  $S_b^0$ . We have

$$\varphi_k^+ = \sum_{a=1}^n U_{ka} S_a^+, \quad (15)$$

$$\chi_j^+ = \sum_{a=1}^n T_{ja} S_a^+, \quad (16)$$

$$\varphi_k^{0'} = \sum_{b=1}^m V_{kb} S_b^0, \quad (17)$$

$$\chi_l^{0'} = \sum_{b=1}^m R_{lb} S_b^0, \quad (18)$$

the matrices  $U$ ,  $T$ ,  $V$  and  $R$  having dimensions  $n_d \times n$ ,  $n_c \times n$ ,  $n_d \times m$  and  $n_n \times m$ , respectively. The matrix  $R$  is real, the other three are complex. The matrix

$$\tilde{U} \equiv \begin{pmatrix} U \\ T \end{pmatrix} \quad (19)$$

is  $n \times n$  unitary; it is the matrix which diagonalizes the (Hermitian) mass matrix of the charged scalars. The real matrix

$$\tilde{V} \equiv \begin{pmatrix} \text{Re } V \\ \text{Im } V \\ R \end{pmatrix} \quad (20)$$

is  $m \times m$  orthogonal; it diagonalizes the (symmetric) mass matrix of the real components of the neutral scalar fields<sup>9</sup>.

There are, in the spontaneously broken  $SU(2) \times U(1)$  theory, three unphysical Goldstone bosons,  $G^\pm$  and  $G^0$ . For definiteness we assign to them the indices  $a = 1$  and  $b = 1$ , respectively:

$$S_1^\pm = G^\pm, \quad (21)$$

$$S_1^0 = G^0. \quad (22)$$

Thus, only the  $S_a^\pm$  with  $a \geq 2$  are physical and, similarly, only the  $S_b^0$  with  $b \geq 2$  correspond to true particles. In the general 't Hooft gauge that we shall use in our computation, the masses of  $G^\pm$  and  $G^0$  are arbitrary and unphysical, and they cannot appear in the final result for  $\Delta\rho$ .

<sup>8</sup> Since the neutral singlet fields carry no hypercharge, their VEVs  $u_l$  do not contribute to the masses of the gauge bosons.

<sup>9</sup> Our treatment of the mixing of scalars is inspired by [21].

## 2.2. The result

As we shall demonstrate in the next section, the value of  $\Delta\rho$  in the model outlined above is

$$\Delta\rho = \frac{g^2}{64\pi^2 m_W^2} \left\{ \sum_{a=2}^n \sum_{b=2}^m |(U^\dagger V)_{ab}|^2 F(m_a^2, \mu_b^2) \right. \quad (23a)$$

$$\left. - \sum_{b=2}^{m-1} \sum_{b'=b+1}^m [\text{Im}(V^\dagger V)_{bb'}]^2 F(\mu_b^2, \mu_{b'}^2) \right. \quad (23b)$$

$$\left. - 2 \sum_{a=2}^{n-1} \sum_{a'=a+1}^n |(U^\dagger U)_{aa'}|^2 F(m_a^2, m_{a'}^2) \right. \quad (23c)$$

$$\left. + 3 \sum_{b=2}^m [\text{Im}(V^\dagger V)_{1b}]^2 [F(m_Z^2, \mu_b^2) - F(m_W^2, \mu_b^2)] \right. \quad (23d)$$

$$\left. - 3 [F(m_Z^2, m_h^2) - F(m_W^2, m_h^2)] \right\}, \quad (23e)$$

where  $m_a, m_{a'}$  denote the masses of the charged scalars and  $\mu_b, \mu_{b'}$  denote the masses of the neutral scalars. The term (23b) contains a sum over all pairs of *different* physical neutral scalar particles  $S_b^0$  and  $S_{b'}^0$ ; similarly, the term (23c) contains a sum over all pairs of different charged scalars, excluding the Goldstone bosons  $G^\pm$ , i.e.  $2 \leq a < a' \leq n$ . The term (23e) consists of the subtraction, from the rest of  $\Delta\rho$ , of the SM result— $m_h$  is the mass of the sole SM physical neutral scalar, the so-called Higgs particle.

In equation (23), the function  $F$  of two non-negative arguments  $x$  and  $y$  is

$$F(x, y) \equiv \begin{cases} \frac{x+y}{2} - \frac{xy}{x-y} \ln \frac{x}{y} & \Leftarrow x \neq y, \\ 0 & \Leftarrow x = y. \end{cases} \quad (24)$$

This is a non-negative function, symmetrical under the interchange of its two arguments, and vanishing if and only if these two arguments are equal. This function has the important property that it grows linearly with  $\max(x, y)$ , i.e. quadratically with the heaviest-scalar mass, when that mass becomes very large. Unless there are cancellations, this leads to a quadratic divergence of  $\Delta\rho$  for very heavy scalars (Higgs bosons).

If there are in the model any  $SU(2)$ -singlet scalars with electric charge other than 0 or  $\pm 1$ , then the existence of these scalars does not contribute to  $\Delta\rho$ ; they do not modify equation (23), at one-loop level, in any way.

A simplification occurs when in the model there are no  $SU(2)$ -singlet charged scalars  $\chi_j^\pm$ . In that case, there is no matrix  $T$ , hence the matrix  $U$  is unitary by itself, and the term (23c) vanishes.

When in the model there are no  $SU(2)$ -singlet neutral scalars  $\chi_l^0$ , there is no matrix  $R$ , hence  $\text{Re}(V^\dagger V)_{bb'} = (\text{Re } V^T \text{Re } V + \text{Im } V^T \text{Im } V)_{bb'} = \delta_{bb'}$ . Then, in the terms (23b) and (23d) one may write  $|(V^\dagger V)_{bb'}|^2$  instead of  $[\text{Im}(V^\dagger V)_{bb'}]^2$ .

Thus, in an  $n_d$ -Higgs-doublet model *without any scalar singlets*, one has simply

$$\Delta\rho = \frac{g^2}{64\pi^2 m_W^2} \left\{ \sum_{a=2}^{n_d} \sum_{b=2}^{2n_d} |(U^\dagger V)_{ab}|^2 F(m_a^2, \mu_b^2) \right. \\ \left. - \sum_{b=2}^{2n_d-1} \sum_{b'=b+1}^{2n_d} |(V^\dagger V)_{bb'}|^2 F(\mu_b^2, \mu_{b'}^2) \right\}$$

$$\begin{aligned}
& + 3 \sum_{b=2}^{2n_d} |(V^\dagger V)_{1b}|^2 [F(m_Z^2, \mu_b^2) - F(m_W^2, \mu_b^2)] \\
& - 3 [F(m_Z^2, m_h^2) - F(m_W^2, m_h^2)] \Big\}. \tag{25}
\end{aligned}$$

Our general results have been checked to be consistent with specific results for  $\Delta\rho$  in a few models. These include the results for both the CP conserving version [5, 13] and the CP non-conserving version [16] of the 2HDM<sup>10</sup>. It has also been checked against a model containing one doublet and one scalar singlet [22].

### 3. Derivation of the result

This section contains the derivation of equation (23). It may be skipped by those who are not interested in the details of that derivation.

#### 3.1. The Lagrangian

We use the conventions of [23]. The covariant derivative of the doublets is

$$D_\mu \phi_k = \begin{pmatrix} \partial_\mu \phi_k^+ - i \frac{g}{\sqrt{2}} W_\mu^+ \phi_k^0 + i \frac{g(s_W^2 - c_W^2)}{2c_W} Z_\mu \phi_k^+ + ie A_\mu \phi_k^+ \\ \partial_\mu \phi_k^0 - i \frac{g}{\sqrt{2}} W_\mu^- \phi_k^+ + i \frac{g}{2c_W} Z_\mu \phi_k^0 \end{pmatrix} \tag{26}$$

and the covariant derivative of the charged singlets is

$$D_\mu \chi_j^+ = \partial_\mu \chi_j^+ + i \frac{g s_W^2}{c_W} Z_\mu \chi_j^+ + ie A_\mu \chi_j^+. \tag{27}$$

The covariant derivative of the neutral singlets is, of course, just identical to their ordinary derivative. We use the unitarity of  $\tilde{U}$  in equation (19), in particular

$$(T^\dagger T)_{a'a} = \delta_{a'a} - (U^\dagger U)_{a'a}. \tag{28}$$

We also use the orthogonality of  $\tilde{V}$  in equation (20) to arrive at the gauge-kinetic Lagrangian

$$\begin{aligned}
& \sum_{k=1}^{n_d} (D^\mu \phi_k)^\dagger (D_\mu \phi_k) + \sum_{j=1}^{n_c} (D^\mu \chi_j^-) (D_\mu \chi_j^+) + \frac{1}{2} \sum_{l=1}^{n_n} (\partial^\mu \chi_l^0) (\partial_\mu \chi_l^0) \\
& = \sum_{a=1}^n (\partial^\mu S_a^-) (\partial_\mu S_a^+) + \frac{1}{2} \sum_{b=1}^m (\partial^\mu S_b^0) (\partial_\mu S_b^0) \tag{29a}
\end{aligned}$$

$$+ m_W^2 W^{\mu-} W_\mu^+ + m_Z^2 \frac{Z^\mu Z_\mu}{2} \tag{29b}$$

$$+ im_W \sum_{a=1}^n [W_\mu^- (\omega^\dagger U)_a \partial^\mu S_a^+ - W_\mu^+ (U^\dagger \omega)_a \partial^\mu S_a^-] \tag{29c}$$

$$+ m_Z Z_\mu \sum_{b=1}^m \text{Im}(\omega^\dagger V)_b \partial^\mu S_b^0 \tag{29d}$$

<sup>10</sup> There is some discrepancy between our result and that presented in section 4 of [11].

$$- (em_W A^\mu + gs_W^2 m_Z Z^\mu) \sum_{a=1}^n [(\omega^\dagger U)_a W_\mu^- S_a^+ + (U^\dagger \omega)_a W_\mu^+ S_a^-] \quad (29e)$$

$$+ ie A_\mu \sum_{a=1}^n (S_a^+ \partial^\mu S_a^- - S_a^- \partial^\mu S_a^+) \quad (29f)$$

$$+ i \frac{g}{2c_W} Z_\mu \sum_{a,a'=1}^n [2s_W^2 \delta_{aa'} - (U^\dagger U)_{a'a}] (S_a^+ \partial^\mu S_{a'}^- - S_{a'}^- \partial^\mu S_a^+) \quad (29g)$$

$$+ \frac{g}{2c_W} Z_\mu \sum_{b=1}^{m-1} \sum_{b'=b+1}^m \text{Im}(V^\dagger V)_{bb'} (S_b^0 \partial^\mu S_{b'}^0 - S_{b'}^0 \partial^\mu S_b^0) \quad (29h)$$

$$+ i \frac{g}{2} \sum_{a=1}^n \sum_{b=1}^m [(U^\dagger V)_{ab} W_\mu^+ (S_a^- \partial^\mu S_b^0 - S_b^0 \partial^\mu S_a^-) + (V^\dagger U)_{ba} W_\mu^- (S_b^0 \partial^\mu S_a^+ - S_a^+ \partial^\mu S_b^0)] \quad (29i)$$

$$+ g \left( m_W W_\mu^+ W^{\mu-} + \frac{m_Z}{c_W} \frac{Z_\mu Z^\mu}{2} \right) \sum_{b=1}^m S_b^0 \text{Re}(\omega^\dagger V)_b \quad (29j)$$

$$- \left( \frac{eg}{2} A^\mu + \frac{g^2 s_W^2}{2c_W} Z^\mu \right) \sum_{a=1}^n \sum_{b=1}^m S_b^0 [(U^\dagger V)_{ab} W_\mu^+ S_a^- + (V^\dagger U)_{ba} W_\mu^- S_a^+] \quad (29k)$$

$$+ \left( \frac{g^2}{4} W^{\mu-} W_\mu^+ + \frac{g^2}{4c_W^2} \frac{Z^\mu Z_\mu}{2} \right) \sum_{b,b'=1}^m (V^\dagger V)_{b'b} S_{b'}^0 S_b^0 \quad (29l)$$

$$+ \frac{g^2}{2} W^{\mu-} W_\mu^+ \sum_{a,a'=1}^n (U^\dagger U)_{a'a} S_{a'}^- S_a^+ \quad (29m)$$

$$+ 2e^2 \frac{A^\mu A_\mu}{2} \sum_{a=1}^n S_a^- S_a^+ \quad (29n)$$

$$+ \frac{eg}{c_W} A^\mu Z_\mu \sum_{a,a'=1}^n [2s_W^2 \delta_{aa'} - (U^\dagger U)_{a'a}] S_{a'}^- S_a^+ \quad (29o)$$

$$+ \frac{g^2}{2c_W^2} \frac{Z^\mu Z_\mu}{2} \sum_{a,a'=1}^n [4s_W^4 \delta_{aa'} + (1 - 4s_W^2) (U^\dagger U)_{a'a}] S_{a'}^- S_a^+. \quad (29p)$$


In lines (29c)–(29e) and (29j) we have used an  $n_d$  vector  $\omega$  defined by  $\omega_k \equiv v_k/v$ . By identifying lines (29c) and (29d) with the usual terms [23] mixing the  $W^\pm$  and  $Z^0$  gauge bosons with the  $G^\pm$  and  $G^0$  Goldstone bosons, respectively,

$$im_W (W_\mu^- \partial^\mu G^+ - W_\mu^+ \partial^\mu G^-) + m_Z Z_\mu \partial^\mu G^0,$$

we conclude that the components of the Goldstone bosons are given by [21]

$$U_{k1} = \frac{v_k}{v}, \quad \text{hence} \quad T_{j1} = 0, \quad (30)$$

$$V_{k1} = i \frac{v_k}{v}, \quad \text{hence} \quad R_{l1} = 0. \quad (31)$$



**Figure 1.** Three types of Feynman diagrams occurring in the calculation of the vacuum polarizations.

Therefore, we may rewrite line (29e) as

$$-(em_W A^\mu + gs_W^2 m_Z Z^\mu) (W_\mu^- G^+ + W_\mu^+ G^-) \quad (32)$$

and line (29j) as

$$-g \left( m_W W_\mu^+ W^{\mu-} + \frac{m_Z}{c_W} \frac{Z_\mu Z^\mu}{2} \right) \sum_{b=2}^m S_b^0 \text{Im}(V^\dagger V)_{1b}. \quad (33)$$

The sum starts at  $b = 2$  because  $\text{Im}(V^\dagger V)_{11} = 0$ .

If there are in the theory any  $SU(2)$ -singlet scalars  $S^{\pm Q}$  with electric charges  $\pm Q$  other than 0 or  $\pm 1$ , then those scalars do not mix with components of the doublets. Their covariant derivative is

$$D_\mu S^{+Q} = \partial_\mu S^{+Q} + i \frac{gs_W^2 Q}{c_W} Z_\mu S^{+Q} + ie Q A_\mu S^{+Q}. \quad (34)$$

This yields, in particular, the following two interaction terms in the Lagrangian:

$$\mathcal{L} = \dots + i \frac{gs_W^2 Q}{c_W} Z_\mu (S^{+Q} \partial^\mu S^{-Q} - S^{-Q} \partial^\mu S^{+Q}) \quad (35a)$$

$$+ \left( \frac{gs_W^2 Q}{c_W} \right)^2 Z_\mu Z^\mu S^{-Q} S^{+Q}. \quad (35b)$$

### 3.2. The Feynman diagrams

In our model, in the computation of the vacuum polarizations of the gauge bosons  $W^\pm$  and  $Z^0$  there are four types of Feynman diagrams involving scalar fields:

*Type (a) diagrams:* a scalar branches off from the gauge-boson line and loops back to *the same point* in that gauge-boson line—see figure 1(a). When the scalar is neutral, the relevant interaction terms in the Lagrangian are those in line (29l), for  $b' = b$ ; but then the contribution to  $\Delta\rho$  vanishes, since one obtains  $\Pi_{WW}^{\mu\nu} = c_W^2 \Pi_{ZZ}^{\mu\nu}$ . When the scalar is charged, the relevant terms in the Lagrangian are those in line (29m) for  $\Pi_{WW}^{\mu\nu}$  and line (29p) for  $\Pi_{ZZ}^{\mu\nu}$ , in both cases for  $a' = a$ .

*Type (b) diagrams:* the gauge-boson line splits into two scalar lines which later reunite to form a new gauge-boson line—see figure 1(b). The relevant terms in the Lagrangian are those in line (29i) for  $\Pi_{WW}^{\mu\nu}$ , and those in lines (29g) and (29h) for  $\Pi_{ZZ}^{\mu\nu}$ .

*Type (c) diagrams:* a neutral scalar branches off from the gauge-boson line and loops to *a later point* in that gauge-boson line—see figure 1(c). The interaction terms in the Lagrangian responsible for these Feynman diagrams are those in expression (33).






Figure 2. Tadpole diagrams which do not contribute to  $\Delta\rho$ .

Type (d) diagrams: a neutral scalar branches off, with zero momentum, from the gauge-boson line, and produces a loop of some stuff—see figure 2. These ‘tadpole’ Feynman diagrams originate from the interaction terms in expression (33). They yield a vanishing contribution to  $\Delta\rho$  since one obtains  $\Pi_{WW}^{\mu\nu} = c_W^2 \Pi_{ZZ}^{\mu\nu}$ . Hence we may omit the tadpole diagrams altogether.

### 3.3. Computation of the loop diagrams

We use dimensional regularization in the computation of the Feynman diagrams. The dimension of space-time is  $d$ . An unphysical mass  $\mu$  is used to keep the dimension of each integral unchanged when  $d$  varies. We define the divergent quantity

$$\text{div} \equiv \frac{2}{4-d} - \gamma + 1 + \ln(4\pi\mu^2),$$

where  $\gamma$  is Euler’s constant. In the computation of type (a) Feynman diagrams the relevant momentum integral is

$$\mu^{4-d} \int \frac{d^d k}{(2\pi)^d} \frac{g^{\mu\nu}}{k^2 - A + i\epsilon} = \frac{ig^{\mu\nu}}{16\pi^2} A(\text{div} - \ln A), \quad (36)$$

where  $A$  is the mass squared of the scalar particle in the loop. In order to compute the type (b) and type (c) Feynman diagrams we need first to introduce a Feynman parameter  $x$ , which is later integrated over from  $x = 0$  to  $x = 1$ . For type (b) diagrams we have

$$\begin{aligned} \mu^{4-d} \int \frac{d^d k}{(2\pi)^d} \int_0^1 dx \frac{4k^\mu k^\nu}{[k^2 - Ax - B(1-x) + i\epsilon]^2} &= \frac{ig^{\mu\nu}}{16\pi^2} [A(\text{div} - \ln A) \\ &+ B(\text{div} - \ln B) + F(A, B)], \end{aligned} \quad (37)$$

where  $A$  and  $B$  are the masses squared of the scalars in the loop, and the 4-momentum  $q^\mu$  of the external gauge-boson line is taken to obey  $q^2 = 0$ . Note the presence of terms of the form  $A(\text{div} - \ln A)$  in both diagrams of types (a) and (b); we shall soon see that those terms cancel out in the computation of  $\Delta\rho$ , leaving only the  $F$  functions from the type (b) diagrams. For type (c) diagrams the relevant integral is

$$\begin{aligned} \mu^{4-d} \int \frac{d^d k}{(2\pi)^d} \int_0^1 dx \frac{g^{\mu\nu}}{[k^2 - Ax - B(1-x) + i\epsilon]^2} \\ = \frac{ig^{\mu\nu}}{16\pi^2} \frac{1}{A} \left[ A(\text{div} - \ln A) - \frac{A+B}{2} + F(A, B) \right]. \end{aligned} \quad (38)$$

This integral is symmetric under the interchange of  $A$  and  $B$ ; equation (38) presents a seemingly asymmetric form, but it is in fact symmetric. The reason for expressing the integral in this way is that, due to cancellations, only the terms  $F(A, B)$  survive in the end.

### 3.4. The contributions to $\Delta\rho$ from diagrams of types (a) and (b)

Using (29m) and (36), we see that the contribution to  $A_{WW}(q^2)$  of type (a) Feynman diagrams with charged scalars in the loop is

$$A_{WW}^{(a)}(q^2) = -\frac{g^2}{32\pi^2} \sum_{a=1}^n (U^\dagger U)_{aa} m_a^2 (\text{div} - \ln m_a^2). \quad (39)$$

In the same way, using (29p),

$$A_{ZZ}^{(a)}(q^2) = -\frac{g^2}{32\pi^2 c_W^2} \sum_{a=1}^n [4s_W^4 + (1 - 4s_W^2) (U^\dagger U)_{aa}] m_a^2 (\text{div} - \ln m_a^2). \quad (40)$$

Proceeding to the type (b) Feynman diagrams, from (29i) and (37) we find that

$$\begin{aligned} A_{WW}^{(b)}(0) &= \frac{g^2}{64\pi^2} \sum_{a=1}^n \sum_{b=1}^m (U^\dagger V)_{ab} (V^\dagger U)_{ba} [m_a^2 (\text{div} - \ln m_a^2) \\ &\quad + \mu_b^2 (\text{div} - \ln \mu_b^2) + F(m_a^2, \mu_b^2)] \\ &= \frac{g^2}{64\pi^2} \left[ 2 \sum_{a=1}^n (U^\dagger U)_{aa} m_a^2 (\text{div} - \ln m_a^2) \right. \end{aligned} \quad (41a)$$

$$\left. + \sum_{b=1}^m (V^\dagger V)_{bb} \mu_b^2 (\text{div} - \ln \mu_b^2) \right. \quad (41b)$$

$$\left. + \sum_{a=1}^n \sum_{b=1}^m |(U^\dagger V)_{ab}|^2 F(m_a^2, \mu_b^2) \right]. \quad (41c)$$

We have used

$$\sum_{a=1}^n (U^\dagger V)_{ab} (V^\dagger U)_{ba} = (V^\dagger V)_{bb}, \quad (42)$$

which follows from the unitarity of  $\tilde{U}$ , i.e. from [21]

$$UU^\dagger = 1_{n_d \times n_d}. \quad (43)$$

We have also used

$$\sum_{b=1}^m (U^\dagger V)_{ab} (V^\dagger U)_{ba} = 2(U^\dagger U)_{aa}, \quad (44)$$

which follows from the orthogonality of  $\tilde{V}$ , i.e. from [21]

$$\begin{aligned} \text{Re } V \text{ Re } V^T &= \text{Im } V \text{ Im } V^T = 1_{n_d \times n_d}, \\ \text{Re } V \text{ Im } V^T &= \text{Im } V \text{ Re } V^T = 0_{n_d \times n_d}. \end{aligned} \quad (45)$$

Considering now the self-energy of the  $Z^0$  boson, we find

$$\begin{aligned} A_{ZZ}^{(b)}(0) &= \frac{g^2}{64\pi^2 c_W^2} \left\{ \sum_{a,a'=1}^n [2s_W^2 \delta_{aa'} - (U^\dagger U)_{a'a}] [2s_W^2 \delta_{aa'} - (U^\dagger U)_{aa'}] \right. \\ &\quad \left. \times [m_a^2 (\text{div} - \ln m_a^2) + m_{a'}^2 (\text{div} - \ln m_{a'}^2) + F(m_a^2, m_{a'}^2)] \right\} \end{aligned}$$

$$\begin{aligned}
& + \sum_{b=1}^{m-1} \sum_{b'=b+1}^m [\text{Im}(V^\dagger V)_{bb'}]^2 \\
& \times \left[ \mu_b^2 (\text{div} - \ln \mu_b^2) + \mu_{b'}^2 (\text{div} - \ln \mu_{b'}^2) + F(\mu_b^2, \mu_{b'}^2) \right] \Big\} \\
= & \frac{g^2}{64\pi^2 c_W^2} \left\{ 2 \sum_{a=1}^{n-1} \sum_{a'=a+1}^n |(U^\dagger U)_{aa'}|^2 F(m_a^2, m_{a'}^2) \right. \tag{46a}
\end{aligned}$$

$$+ 2 \sum_{a=1}^n [4s_W^4 + (1 - 4s_W^2)(U^\dagger U)_{aa}] m_a^2 (\text{div} - \ln m_a^2) \tag{46b}$$

$$+ \sum_{b=1}^{m-1} \sum_{b'=b+1}^m [\text{Im}(V^\dagger V)_{bb'}]^2 F(\mu_b^2, \mu_{b'}^2) \tag{46c}$$

$$+ \left. \sum_{b=1}^m (V^\dagger V)_{bb} \mu_b^2 (\text{div} - \ln \mu_b^2) \right\}. \tag{46d}$$

We have used

$$\sum_{b'=1}^m [\text{Im}(V^\dagger V)_{bb'}]^2 = (V^\dagger V)_{bb}, \tag{47}$$

which follows from equations (45).

Putting everything together, we see that

the  $A_{WW}^{(a)}(q^2)$  of equation (39) cancels out the line (41a) of  $A_{WW}^{(b)}(0)$ ;

the  $A_{ZZ}^{(a)}(q^2)$  of equation (40) cancels out the line (46b) of  $A_{ZZ}^{(b)}(0)$ ;

the line (41b) of  $A_{WW}^{(b)}(0)$  cancels out the line (46d) of  $A_{ZZ}^{(b)}(0)$  in the subtraction  $A_{WW} - c_W^2 A_{ZZ}$ .

In this way we finally obtain

$$A_{WW}^{(a+b)}(0) - c_W^2 A_{ZZ}^{(a+b)}(0) = \frac{g^2}{64\pi^2} \left\{ \sum_{a=1}^n \sum_{b=1}^m |(U^\dagger V)_{ab}|^2 F(m_a^2, \mu_b^2) \right. \tag{48a}$$

$$- 2 \sum_{a=1}^{n-1} \sum_{a'=a+1}^n |(U^\dagger U)_{aa'}|^2 F(m_a^2, m_{a'}^2) \tag{48b}$$

$$- \left. \sum_{b=1}^{m-1} \sum_{b'=b+1}^m [\text{Im}(V^\dagger V)_{bb'}]^2 F(\mu_b^2, \mu_{b'}^2) \right\}. \tag{48c}$$

The positive term (48a) originates from  $A_{WW}^{(b)}$  while the negative terms (48b) and (48c) come from  $A_{ZZ}^{(b)}$ .

If there are in the electroweak theory any scalar  $SU(2)$  singlets with electric charges other than 0 or  $\pm 1$ , then the relevant terms in the Lagrangian are those in equation (35). The term (35b) generates a type (a) Feynman diagram which exactly cancels the type (b) Feynman diagram generated by the term (35a)<sup>11</sup>. Thus, scalar  $SU(2)$  singlets with electric charge different from 0 and  $\pm 1$  do not affect  $\Delta\rho$  at all.

<sup>11</sup> This cancellation is analogous to that between equation (40) and line (46b).

The sums in equation (48) include contributions from the Goldstone bosons  $G^\pm = S_1^\pm$  and  $G^0 = S_1^0$ . These Goldstone bosons have unphysical masses  $m_1$  and  $\mu_1$ , respectively, which are arbitrary in a 't Hooft gauge. The terms which depend on those masses are, explicitly,

$$|(U^\dagger V)_{11}|^2 F(m_1^2, \mu_1^2) \quad (49a)$$

$$+ \sum_{b=2}^m |(U^\dagger V)_{1b}|^2 F(m_1^2, \mu_b^2) \quad (49b)$$

$$+ \sum_{a=2}^n |(U^\dagger V)_{a1}|^2 F(m_a^2, \mu_1^2) \quad (49c)$$

$$- 2 \sum_{a=2}^n |(U^\dagger U)_{1a}|^2 F(m_1^2, m_a^2) \quad (49d)$$

$$- \sum_{b=2}^m [\text{Im}(V^\dagger V)_{1b}]^2 F(\mu_1^2, \mu_b^2). \quad (49e)$$

One may eliminate some of these terms by using equations (30) and (31). Indeed,  $(U^\dagger U)_{1a} = -(T^\dagger T)_{1a} = 0$  when  $a \neq 1$ , because  $T_{j1} = 0$  for any  $j$ ; also,  $(U^\dagger V)_{a1} = i(U^\dagger U)_{a1} = 0$  for  $a \neq 1$ . Therefore, the terms (49c) and (49d) vanish. In the term (49a),  $(U^\dagger V)_{11} = i$ . In the term (49b) one may write

$$(U^\dagger V)_{1b} = i(V^\dagger V)_{1b} = -\text{Im}(V^\dagger V)_{1b} \Leftrightarrow b \neq 1, \quad (50)$$

since  $\text{Re}(V^\dagger V)_{1b} = (\text{Re } V^T \text{Re } V + \text{Im } V^T \text{Im } V)_{1b} = -(R^T R)_{1b} = 0$ . In this way, the terms (49) are reduced to

$$F(m_1^2, \mu_1^2) + \sum_{b=2}^m [\text{Im}(V^\dagger V)_{1b}]^2 [F(m_1^2, \mu_b^2) - F(\mu_1^2, \mu_b^2)]. \quad (51)$$

The term  $F(m_1^2, \mu_1^2)$  is independent of the number of scalar doublets and singlets, hence it is eliminated when one subtracts the SM result from the multi-Higgs-doublet model one. The other terms in expression (51) are cancelled out by the diagrams of type (c), as we shall see next.

### 3.5. The contributions to $\Delta\rho$ from the diagrams of type (c)

To compensate for the unphysical masses of the Goldstone bosons, the propagators of gauge bosons  $W^\pm$  and  $Z^0$  with 4-momentum  $k^\mu$  are, in a 't Hooft gauge,

$$-\frac{k_\mu k_\nu}{m_W^2} \frac{i}{k^2 - m_1^2} + \left( -g_{\mu\nu} + \frac{k_\mu k_\nu}{m_W^2} \right) \frac{i}{k^2 - m_W^2}, \quad (52)$$

$$-\frac{k_\mu k_\nu}{m_Z^2} \frac{i}{k^2 - \mu_1^2} + \left( -g_{\mu\nu} + \frac{k_\mu k_\nu}{m_Z^2} \right) \frac{i}{k^2 - m_Z^2}, \quad (53)$$

respectively, i.e. they contain a piece with a pole on the unphysical masses squared  $m_1^2$  and  $\mu_1^2$ , respectively.

Using these propagators to compute the type (c) Feynman diagrams, one obtains

$$A_{WW}^{(c)}(0) = \frac{g^2}{64\pi^2} \sum_{b=2}^m [\text{Im}(V^\dagger V)_{1b}]^2 [-m_1^2 (\text{div} - \ln m_1^2) - 3m_W^2 (\text{div} - \ln m_W^2) + 2(m_W^2 + \mu_b^2) - F(m_1^2, \mu_b^2) - 3F(m_W^2, \mu_b^2)], \quad (54)$$

$$A_{ZZ}^{(c)}(0) = \frac{g^2}{64\pi^2 c_W^2} \sum_{b=2}^m [\text{Im}(V^\dagger V)_{1b}]^2 [-\mu_1^2 (\text{div} - \ln \mu_1^2) - 3m_Z^2 (\text{div} - \ln m_Z^2) + 2(m_Z^2 + \mu_b^2) - F(\mu_1^2, \mu_b^2) - 3F(m_Z^2, \mu_b^2)]. \quad (55)$$

The factors 3 originate in a partial cancellation between the contributions from the pieces  $-g_{\mu\nu}$  and  $k_\mu k_\nu / m_V^2$  in the propagator of the gauge boson  $V$ , the former contribution being four times larger than, and with opposite sign relative to, the latter one, cf equations (37) and (38). Performing the subtraction relevant for  $\Delta\rho$ , one obtains

$$A_{WW}^{(c)}(0) - c_W^2 A_{ZZ}^{(c)}(0) = \frac{g^2}{64\pi^2} \sum_{b=2}^m [\text{Im}(V^\dagger V)_{1b}]^2 \times [-m_1^2 (\text{div} - \ln m_1^2) + \mu_1^2 (\text{div} - \ln \mu_1^2) - 3m_W^2 (\text{div} - \ln m_W^2) + 3m_Z^2 (\text{div} - \ln m_Z^2) + 2(m_W^2 - m_Z^2) - F(m_1^2, \mu_b^2) + F(\mu_1^2, \mu_b^2) - 3F(m_W^2, \mu_b^2) + 3F(m_Z^2, \mu_b^2)]. \quad (56a)$$

$$- 3m_W^2 (\text{div} - \ln m_W^2) + 3m_Z^2 (\text{div} - \ln m_Z^2) \quad (56b)$$

$$+ 2(m_W^2 - m_Z^2) \quad (56c)$$

$$- F(m_1^2, \mu_b^2) + F(\mu_1^2, \mu_b^2) \quad (56d)$$

$$- 3F(m_W^2, \mu_b^2) + 3F(m_Z^2, \mu_b^2)]. \quad (56e)$$

The terms (56a)–(56c) are independent of the number of scalar doublets. They disappear when one subtracts the Standard-Model result from the multi-Higgs-doublet model one, since

$$\sum_{b=2}^m [\text{Im}(V^\dagger V)_{1b}]^2 = (V^\dagger V)_{11} = 1. \quad (57)$$

The terms (56d), which involve the masses of the Goldstone bosons, cancel out the terms in (51) except the first one, which is cancelled by the subtraction of the SM result.

We have thus finished the derivation of equation (23) for  $\Delta\rho$ .

#### 4. The 2HDM and the Zee model

In this section we give, as examples of the application of our general formulae, the expressions for  $\Delta\rho$  in the 2HDM and also in the model of Zee [24] for the radiative generation of neutrino masses, which has one singly-charged  $SU(2)$  singlet together with the two doublets.

In the study of the 2HDM it is convenient to use the so-called ‘Higgs basis’, in which only the first Higgs doublet has a vacuum expectation value. In that basis,

$$\phi_1 = \begin{pmatrix} G^+ \\ (v + H + iG^0)/\sqrt{2} \end{pmatrix}, \quad \phi_2 = \begin{pmatrix} S_2^+ \\ (R + iI)/\sqrt{2} \end{pmatrix}. \quad (58)$$

Here,  $G^+ \equiv S_1^+$  and  $G^0 \equiv S_1^0$  are the Goldstone bosons, while  $S_2^+$  is the physical charged scalar, which has mass  $m_2$ . Thus, the matrix  $U$ , which connects the charged components of

$\phi_1$  and  $\phi_2$  to the eigenstates of mass, is in the Higgs basis of the 2HDM equal to the unit matrix. On the other hand,  $H$ ,  $R$  and  $I$ , which are real fields, must be rotated through a  $3 \times 3$  orthogonal matrix  $O$  to obtain the three physical neutral fields  $S_{2,3,4}^0$ :

$$\begin{pmatrix} H \\ R \\ I \end{pmatrix} = O \begin{pmatrix} S_2^0 \\ S_3^0 \\ S_4^0 \end{pmatrix}. \quad (59)$$

Without lack of generality we choose  $\det O = +1$ . Thus, the  $2 \times 4$  matrix  $V$ , defined through

$$\begin{pmatrix} H + iG^0 \\ R + iI \end{pmatrix} = V \begin{pmatrix} G^0 \\ S_2^0 \\ S_3^0 \\ S_4^0 \end{pmatrix}, \quad (60)$$

is

$$V = \begin{pmatrix} i & O_{11} & O_{12} & O_{13} \\ 0 & O_{21} + iO_{31} & O_{22} + iO_{32} & O_{23} + iO_{33} \end{pmatrix}. \quad (61)$$

Therefore,

$$V^\dagger V = \begin{pmatrix} 1 & -iO_{11} & -iO_{12} & -iO_{13} \\ iO_{11} & 1 & iO_{13} & -iO_{12} \\ iO_{12} & -iO_{13} & 1 & iO_{11} \\ iO_{13} & iO_{12} & -iO_{11} & 1 \end{pmatrix}. \quad (62)$$

The value of  $\Delta\rho$  in the 2HDM is therefore, using our formula in equation (25),

$$\begin{aligned} \Delta\rho = \frac{g^2}{64\pi^2 m_W^2} & \left\{ \sum_{b=2}^4 (1 - O_{1b-1}^2) F(m_b^2, \mu_b^2) \right. \\ & - O_{13}^2 F(\mu_2^2, \mu_3^2) - O_{12}^2 F(\mu_2^2, \mu_4^2) - O_{11}^2 F(\mu_3^2, \mu_4^2) \\ & \left. + 3 \sum_{b=2}^4 O_{1b-1}^2 [F(m_Z^2, \mu_b^2) - F(m_W^2, \mu_b^2) - F(m_Z^2, m_h^2) + F(m_W^2, m_h^2)] \right\}, \quad (63) \end{aligned}$$

where  $\mu_{2,3,4}$  denote the masses of  $S_{2,3,4}^0$ , respectively, while  $m_h$  is the mass of the Higgs boson of the SM. Equation (63) reproduces, in a somewhat simplified form, the result for  $\Delta\rho$  in the 2HDM previously given in [16].

A special case of the 2HDM is the model with one ‘dark’ scalar doublet. This means that a second doublet is added to the SM, but that doublet has no VEV and it does not mix with the standard Higgs doublet [19]. We should then identify  $H$  with the usual Higgs particle. Thus,  $O_{11} = 1$  and  $\mu_2 = m_h$ . Equation (63) then simplifies to [20, 25]

$$\Delta\rho = \frac{g^2}{64\pi^2 m_W^2} \left[ \sum_{b=3}^4 F(m_b^2, \mu_b^2) - F(\mu_3^2, \mu_4^2) \right]. \quad (64)$$

This quantity is small if the three masses  $m_2$ ,  $\mu_3$  and  $\mu_4$  are close together. Note that in this case of a ‘dark’ scalar doublet there are no vector–vector–scalar couplings involving the additional doublet, hence  $\Delta\rho$  stems exclusively from the type (a) and type (b) Feynman diagrams.

In the model of Zee there is, besides the two scalar  $SU(2)$  doublets

$$\phi_1 = \begin{pmatrix} G^+ \\ (v + H + iG^0)/\sqrt{2} \end{pmatrix}, \quad \phi_2 = \begin{pmatrix} H^+ \\ (R + iI)/\sqrt{2} \end{pmatrix}, \quad (65)$$

also one scalar  $SU(2)$  singlet  $\chi^+$  with unit electric charge. Therefore there is a  $2 \times 2$  unitary matrix  $K$  such that

$$\begin{pmatrix} H^+ \\ \chi^+ \end{pmatrix} = K \begin{pmatrix} S_2^+ \\ S_3^+ \end{pmatrix}, \quad (66)$$

where  $S_2^+$  and  $S_3^+$  are the physical charged scalars, which have masses  $m_2$  and  $m_3$ , respectively. So, now the matrix  $U$  of equation (15) is

$$U = \begin{pmatrix} 1 & 0 & 0 \\ 0 & K_{11} & K_{12} \end{pmatrix}, \quad (67)$$

so that

$$U^\dagger U = \begin{pmatrix} 1 & 0 & 0 \\ 0 & |K_{11}|^2 & K_{11}^* K_{12} \\ 0 & K_{11} K_{12}^* & |K_{12}|^2 \end{pmatrix}. \quad (68)$$

Equations (61) and (62) retain their validity, and

$$U^\dagger V = \begin{pmatrix} i & O_{11} & O_{12} & O_{13} \\ 0 & K_{11}^* (O_{21} + iO_{31}) & K_{11}^* (O_{22} + iO_{32}) & K_{11}^* (O_{23} + iO_{33}) \\ 0 & K_{12}^* (O_{21} + iO_{31}) & K_{12}^* (O_{22} + iO_{32}) & K_{12}^* (O_{23} + iO_{33}) \end{pmatrix}. \quad (69)$$

Therefore, using our general formula (23) for  $\Delta\rho$ , we see that, in the model of Zee,

$$\begin{aligned} \Delta\rho = & \frac{g^2}{64\pi^2 m_W^2} \left\{ \sum_{b=2}^4 (1 - O_{1b-1}^2) \sum_{a=2}^3 |K_{1a-1}|^2 F(m_a^2, \mu_b^2) \right. \\ & - 2 |K_{11} K_{12}|^2 F(m_2^2, m_3^2) \\ & - O_{13}^2 F(\mu_2^2, \mu_3^2) - O_{12}^2 F(\mu_2^2, \mu_4^2) - O_{11}^2 F(\mu_3^2, \mu_4^2) \\ & \left. + 3 \sum_{b=2}^4 O_{1b-1}^2 [F(m_Z^2, \mu_b^2) - F(m_W^2, \mu_b^2) - F(m_Z^2, m_h^2) + F(m_W^2, m_h^2)] \right\}. \quad (70) \end{aligned}$$

## 5. Summary

In this paper we have derived the formula for the parameter  $\Delta\rho$ , as defined in equation (4), in an extension of the Standard Model characterized by an arbitrary number of scalar  $SU(2)$  doublets (with hypercharge  $\pm 1/2$ ) and singlets (with arbitrary hypercharges). Our formalism is completely general, using only the masses of the scalars and their mixing matrices, which ensures that our formulae are always applicable. The computation has been carried out in a general  $R_\xi$  gauge, thereby demonstrating that the final result is independent of the masses of the unphysical scalars. We have also explicitly demonstrated that all infinities cancel out in the final result for  $\Delta\rho$ . In order to ease the consultation of this paper, the formulae for  $\Delta\rho$  given in section 2 have been completely separated from their derivation presented in section 3. Our results can be applied either to check the viability of a model or to constrain its parameter space, by comparing the  $\Delta\rho$ , calculated in that model, with numerical bounds on  $\Delta\rho$  obtained from a fit to precision data—for instance, the bound (7) found in [4]. As an illustration of our general formulae, in section 4 we have worked out the specific cases of the two-Higgs-doublet model, with and without one extra charged scalar singlet.

## Acknowledgments

WG thanks S Dittmaier, W Hollik, M Krawczyk and H Neufeld for helpful discussions. The work of LL was supported by the Portuguese *Fundação para a Ciência e a Tecnologia* through the project U777–Plurianual. WG and LL acknowledge support from EU under the MRTN-CT-2006-035505 network programme.

## References

- [1] Ross D A and Veltman M J G 1975 Neutral currents in neutrino experiments *Nucl. Phys. B* **95** 135  
Veltman M J G 1977 Second threshold in weak interactions *Acta Phys. Pol. B* **8** 475  
Veltman M J G 1977 Limit on mass differences in the Weinberg model *Nucl. Phys. B* **123** 89
- [2] Einhorn M B, Jones D R T and Veltman M J G 1981 Heavy particles and the rho parameter in the Standard Model *Nucl. Phys. B* **191** 146
- [3] Yao W M *et al* (Particle Data Group) 2006 Review of particle physics *J. Phys. G: Nucl. Part. Phys.* **33** 1
- [4] Erler J and Langacker P 2006 Review of particle physics *J. Phys. G: Nucl. Part. Phys.* **33** 119
- [5] Bertolini S 1986 Quantum effects in a two-Higgs-doublet model of the electroweak interactions *Nucl. Phys. B* **272** 77
- [6] Denner A, Guth R J and Kühn J H 1990 Relaxation of top-mass limits in the two-Higgs-doublet model *Phys. Lett. B* **240** 438  
Froggatt C D, Moorhouse R G and Knowles I G 1992 Leading radiative corrections in two-scalar-doublet models *Phys. Rev. D* **45** 2471
- [7] Boughezal R, Tausk J B and van der Bij J J 2005 Three-loop electroweak correction to the rho parameter in the large Higgs mass limit *Nucl. Phys. B* **713** 278 (Preprint [hep-ph/0410216](#)) and references therein
- [8] Peskin M E and Takeuchi T 1990 A new constraint on a strongly interacting Higgs sector *Phys. Rev. Lett.* **65** 964  
Peskin M E and Takeuchi T 1992 Estimation of oblique electroweak corrections *Phys. Rev. D* **46** 381
- [9] Altarelli G and Barbieri R 1991 Vacuum-polarization effects of new physics on electroweak processes *Phys. Lett. B* **253** 161  
Altarelli G, Barbieri R and Jadach S 1992 Toward a model-independent analysis of electroweak data *Nucl. Phys. B* **369** 3  
Altarelli G, Barbieri R and Jadach S 1992 Toward a model-independent analysis of electroweak data *Nucl. Phys. B* **376** 444 (erratum)
- [10] Maksymyk I, Burgess C P and London D 1994 Beyond S, T, and U *Phys. Rev. D* **50** 529 (Preprint [hep-ph/9306267](#))
- [11] Gunion J F, Haber H E, Kane G L and Dawson S 1989 *The Higgs Hunter's Guide* (Reading, MA: Addison-Wesley)
- [12] Hollik W 1986 Nonstandard Higgs bosons in  $SU(2) \times U(1)$  radiative corrections *Z. Phys. C* **32** 291  
Hollik W 1988 Radiative corrections with two Higgs doublets at LEP/SLC and HERA *Z. Phys. C* **37** 569  
Denner A, Guth R J, Hollik W and Kühn J H 1991 The Z width in the two-Higgs-doublet model *Z. Phys. C* **51** 695
- [13] Chankowski P H, Krawczyk M and Żochowski J 1999 Implications of the precision data for very light Higgs boson scenarios in 2HDM(II) *Eur. Phys. J. C* **11** 661 (Preprint [hep-ph/9905436](#))
- [14] Chankowski P H, Farris T, Grzadkowski B, Gunion J F, Kalinowski J and Krawczyk M 2000 Do precision electroweak constraints guarantee  $e^+e^-$  collider discovery of at least one Higgs boson of a two-Higgs-doublet model? *Phys. Lett. B* **496** 195 (Preprint [hep-ph/0009271](#))  
Krawczyk M and Temes D 2005 Large 2HDM(II) one-loop corrections in leptonic tau decays *Eur. Phys. J. C* **44** 435 (Preprint [hep-ph/0410248](#))
- [15] Gunion J F and Haber H E 2003 The CP-conserving two-Higgs-doublet model: the approach to the decoupling limit *Phys. Rev. D* **67** 075019 (Preprint [hep-ph/0207010](#))  
Cheung K and Kong O C W 2003 Can the two-Higgs-doublet model survive the constraints from the muon anomalous magnetic moment as suggested? *Phys. Rev. D* **68** 053003 (Preprint [hep-ph/0302111](#))
- [16] El Kaffas A W, Khater W, OGREID O M and Osland P 2007 Consistency of the two-Higgs-doublet model and CP violation in top production at the LHC *Nucl. Phys. B* **775** 45 (Preprint [hep-ph/0605142](#))
- [17] Grossman Y 1994 Phenomenology of models with more than two Higgs doublets *Nucl. Phys. B* **426** 355 (Preprint [hep-ph/9401311](#))
- [18] Grimus W 2004 *Neutrino Physics—Theory (Lectures on Flavor Physics)* ed U-G Meißner and W Plessas (Berlin: Springer) p 169 (Preprint [hep-ph/0307149](#))



- Grimus W and Lavoura L 2003 Models of maximal atmospheric neutrino mixing *Acta Phys. Pol. B* **34** 5393 (Preprint [hep-ph/0310050](#))
- Mondragón A 2006 Models of flavour with discrete symmetries *AIP Conf. Proc.* **857B** 266 (Preprint [hep-ph/0609243](#))
- [19] Ma E 2006 Verifiable radiative seesaw mechanism of neutrino mass and dark matter *Phys. Rev. D* **73** 077301 (Preprint [hep-ph/0601225](#))
- [20] Barbieri R, Hall L J and Rychkov V S 2006 Improved naturalness with a heavy Higgs: an alternative road to LHC physics *Phys. Rev. D* **74** 015007 (Preprint [hep-ph/0603188](#))
- [21] Grimus W and Neufeld H 1989 Radiative neutrino masses in an  $SU(2) \times U(1)$  model *Nucl. Phys. B* **325** 18
- Grimus W and Lavoura L 2002 Soft lepton-flavor violation in a multi-higgs-doublet seesaw model *Phys. Rev. D* **66** 014016 (Preprint [hep-ph/0204070](#))
- [22] Profumo S, Ramsey-Musolf M J and Shaughnessy G 2007 Singlet Higgs phenomenology and the electroweak phase transition *J. High Energy Phys.* **JHEP08(2007)010** (Preprint [0705.2425](#))
- [23] Branco G C, Lavoura L and Silva J P 1999 *CP Violation* (Oxford: Oxford University Press) chapter 11
- [24] Zee A 1980 A theory of lepton-number violation and neutrino Majorana masses *Phys. Lett.* **93B** 389
- Zee A 1980 A theory of lepton-number violation and neutrino Majorana masses *Phys. Lett.* **95B** 461 (erratum)
- Zee A 1985 Charged scalar field and quantum number violations *Phys. Lett. B* **161** 141
- [25] Lavoura L and Li L-F 1993 Mechanism for obtaining a negative  $T$  oblique parameter *Phys. Rev. D* **48** 234