

Benchmark scenarios for the NMSSM

**A. Djouadi,^{abc} M. Drees,^c U. Ellwanger,^a R. Godbole,^d C. Hugonie,^e S.F. King,^b
S. Lehti,^f S. Moretti,^{ab} A. Nikitenko,^{g*} I. Rottländer,^c M. Schumacher^h and
A.M. Teixeira^a**

^a*Laboratoire de Physique Théorique, Université Paris-Sud,
F-91405 Orsay Cedex, France*

^b*School of Physics and Astronomy, University of Southampton,
Highfield, SO17 1BJ, U.K.*

^c*Physikalisches Institut, University of Bonn,
Nussallee 12, 53115 Bonn, Germany*

^d*Center for High Energy Physics, Indian Institute of Science,
Bangalore 560 012, India*

^e*LPTA, Université de Montpellier II, 34095 Montpellier, France*

^f*Helsinki Institute of Physics, P.O. Box 64, FIN-00014, University of Helsinki, Finland*

^g*Physics Department, Imperial College, Prince Consort Road, London SW7 2AZ, U.K.*

^h*Fachbereich Physik, University of Siegen, Walter Flex Str. 3, 57068 Siegen, Germany*

*E-mail: Abdelhak.Djouadi@th.u-psud.fr, Drees@th.physik.uni-bonn.de,
Ulrich.Ellwanger@th.u-psud.fr, rohini@cts.iisc.ernet.in,
Cyril.Hugonie@free.fr, sfk@hep.phys.soton.ac.uk, sami.lehti@helsinki.fi,
stefano@phys.soton.ac.uk, anikiten@mail.cern.ch,
rottlaender@physik.uni-bonn.de,
markus.schumacher@hep.physik.uni-siegen.de, Ana.Teixeira@th.u-psud.fr*

ABSTRACT: We discuss constrained and semi-constrained versions of the next-to-minimal supersymmetric extension of the Standard Model (NMSSM) in which a singlet Higgs superfield is added to the two doublet superfields that are present in the minimal extension (MSSM). This leads to a richer Higgs and neutralino spectrum and allows for many interesting phenomena that are not present in the MSSM. In particular, light Higgs particles are still allowed by current constraints and could appear as decay products of the heavier Higgs states, rendering their search rather difficult at the LHC. We propose benchmark scenarios which address the new phenomenological features, consistent with present constraints from colliders and with the dark matter relic density, and with (semi-)universal soft terms at the GUT scale. We present the corresponding spectra for the Higgs particles, their couplings to gauge bosons and fermions and their most important decay branching ratios. A brief survey of the search strategies for these states at the LHC is given.

KEYWORDS: Supersymmetric Standard Model, Supersymmetry Phenomenology.

*On leave of absence from ITEP, Moscow, Russia

Contents

1. Introduction	1
2. The NMSSM and its particle spectrum	3
2.1 The unconstrained NMSSM	3
2.2 The constrained NMSSM	4
2.3 The Higgs and SUSY spectra	6
3. The benchmark points	7
4. Implications for the LHC	14
4.1 Dominant production processes	14
4.2 Summary of available studies	15
4.3 ATLAS strategy for NMSSM $h_1^0 \rightarrow a_1^0 a_1^0$ searches	16
4.4 Prospects for the CMS experiment	18
5. Conclusions	19

1. Introduction

The next-to-minimal supersymmetric extension of the standard model (NMSSM), in which the spectrum of the minimal supersymmetric extension (MSSM) is extended by one singlet superfield, has been discussed since the early days of supersymmetry (SUSY) model-building [1–3]. In the last decade, the NMSSM gained a renewed interest in view of its positive features as compared to the widely studied MSSM. Firstly, the NMSSM naturally solves in an elegant way the so-called μ problem [4] of the MSSM: to have an acceptable phenomenology, a value in the vicinity of the electroweak or SUSY breaking scale is needed for the supersymmetric Higgs mass parameter μ ; this is automatically achieved in the NMSSM, since the μ -parameter is dynamically generated when the singlet Higgs field acquires a vacuum expectation value (vev) of the order of the SUSY breaking scale, leading to a fundamental Lagrangian that contains no dimensionful parameters apart from the soft SUSY breaking terms. Secondly, as compared to the MSSM, the parameter space of the NMSSM is somewhat less constrained in the Higgs and neutralino sectors by collider and dark matter (DM) experiments: on the one hand, heavier Higgs states can decay into lighter ones with sizable rates [5–16] and, on the other hand, a new possibility appears for achieving the correct cosmological relic density [17] through the so-called “singlino”, i.e. the fifth neutralino of the model, which can have weaker-than-usual couplings to standard model (SM) particles. Thirdly, the NMSSM needs somewhat less fine tuning [8, 18] (although some fine tuning is still required [19]): the upper limit on the mass of the lightest

CP-even Higgs particle is larger than in the MSSM, and therefore more SUSY parameter space survives the bounds imposed by the negative Higgs boson searches at LEP; furthermore, possible unconventional decays of the SM-like Higgs scalar allow it to have a relatively small mass, well below the SM Higgs mass limit of 114 GeV, still consistent with LEP constraints.

Given the possibly quite different phenomenology in the Higgs sector as compared to the SM and the MSSM [20], it is important to address the question of whether such NMSSM specific scenarios will be probed at the LHC. In particular, it would be important to extend the so-called “no-lose theorem” of the MSSM [21], which states that at least one MSSM Higgs particle should be observed at the LHC for the planned integrated luminosity, to the NMSSM [22, 6] or try to define regions of the NMSSM parameter space where more Higgs states are visible at the LHC than those available within the MSSM [23]. However, a potential drawback of the NMSSM, at least in its non constrained versions, is that it leads to a larger number of input parameters in the Higgs sector to deal with, compared to the MSSM. In particular, it is clearly unfeasible to perform multi-dimensional “continuous” scans over the free inputs of the NMSSM, especially if each sampled point is subject to a complete simulation in order to be as close as possible to the experimental conditions.

An alternative approach, acknowledged by both the theoretical and experimental communities, is that of resorting to the definition of so-called “benchmark points” (or slopes, or surfaces) in the SUSY parameter space [24]. These consist of a few “discrete” parameter configurations of a given SUSY model, which embody the most peculiar/representative phenomenological features of the model’s parameter space. Using discrete points avoids scanning the entire parameter space, focusing instead on representative choices that reflect the new interesting features of the model, such as new signals, peculiar mass spectra, etc. A reduced number of points can then be subject to full experimental investigation, without loss of substantial theoretical information.

While several such benchmark scenarios have been devised for the MSSM [24, 25] and thoroughly studied in both the collider and the DM contexts, limited progress has been made so far in this direction for the case of the NMSSM. In ref. [9], an earlier attempt was made to address the possibility of establishing a no-lose theorem for the NMSSM Higgs sector at the LHC through benchmark points (in a non-universal low energy setup in which the SUSY particles are very heavy). However, the corresponding spectrum was not consistent with DM constraints (in particular on the lightest SUSY particle, LSP), which were not yet established for the NMSSM at the time. In addition, many of the points discussed in ref. [9] are now ruled out due to the new lower value of the top quark mass, and to more stringent constraints from collider searches and precision measurements. The tools to calculate the Higgs and SUSY particle spectra have also been upgraded since then [26, 27].

In this report, we build on the experience of ref. [9] and define benchmark points which fulfill the present collider and cosmological constraints using the most up-to-date tools to calculate the particle spectra. However, in contrast to ref. [9], we work in the framework of a (semi-)constrained NMSSM parameter space, henceforth called cNMSSM, where the soft SUSY-breaking parameters are defined at the Grand Unified (GUT) scale:

on the one hand, this approach leads to a much more plausible sparticle spectrum and allows one to relate features of the Higgs sector to properties of the neutralino sector (and hence of the LSP); on the other hand, this restricted parameter space still contains the phenomenological features of the NMSSM that are very distinctive from those of the MSSM, and is suitable for intensive phenomenological investigation by the experimental collaborations. The emphasis will primarily be on the different possible scenarios within the Higgs sector as light pseudoscalars, scalars below 114 GeV decaying dominantly into light pseudoscalars or having a large singlet component, and the implication for Higgs searches at the LHC. All these scenarios can be realized within the (semi-)constrained NMSSM considered below, and comply with the cosmological relic density of the lightest neutralino DM candidate¹ (which, in one case, is dominantly singlino like). Finally, we will describe the tools used to define such benchmark scenarios.

The report is organized as follows. In the next section, we define the NMSSM and its particle content with some emphasis on constrained scenarios defined in terms of soft terms specified at the GUT scale, discuss the tools that allow to calculate the particle spectrum, and the various constraints that should be imposed on the latter. In section 3, we propose five benchmark points which lead to Higgs sectors that are different from those of the MSSM and discuss their main features. In section 4, we outline the possible search strategies at the LHC for the Higgs particles in these scenarios. A brief outlook is given in section 5.

2. The NMSSM and its particle spectrum

2.1 The unconstrained NMSSM

In this paper, we confine ourselves to the NMSSM with a scale invariant superpotential. Alternative generalizations of the MSSM — known as the minimal non-minimal supersymmetric SM (MNSSM), new minimally-extended supersymmetric SM or nearly-minimal supersymmetric SM (nMSSM) or with additional U(1)' gauge symmetries — exist [28], but these will not be considered here, nor the case of explicit CP violation [29]. The scale invariant superpotential of the NMSSM is given, in terms of (hatted) superfields, by

$$\mathcal{W} = \lambda \widehat{S} \widehat{H}_u \widehat{H}_d + \frac{\kappa}{3} \widehat{S}^3 + h_t \widehat{Q} \widehat{H}_u \widehat{t}_R^c - h_b \widehat{Q} \widehat{H}_d \widehat{b}_R^c - h_\tau \widehat{L} \widehat{H}_d \widehat{\tau}_R^c \quad (2.1)$$

in which only the third generation fermions have been included (possible neutrino Yukawa couplings have been set to zero), and \widehat{Q}, \widehat{L} stand for superfields associated with the (t, b) and (τ, ν_τ) SU(2) doublets. The first two terms substitute the $\mu \widehat{H}_u \widehat{H}_d$ term in the MSSM superpotential, while the three last terms are the usual generalization of the Yukawa interactions. The soft SUSY breaking terms consist of the scalar mass terms for the Higgs and sfermion scalar fields which, in terms of the fields corresponding to the complex scalar

¹In this analysis we cover practically all the scenarios involving a light neutral Higgs boson. We do not expect that new scenarios, with a Higgs phenomenology that strongly differs from the cases here discussed, could be found by relaxing the cosmological relic density constraint.

components of the superfields, are given by

$$\begin{aligned}
 -\mathcal{L}_{\text{mass}} = & m_{H_u}^2 |H_u|^2 + m_{H_d}^2 |H_d|^2 + m_S^2 |S|^2 \\
 & + m_{\tilde{Q}}^2 |\tilde{Q}^2| + m_{\tilde{t}_R}^2 |\tilde{t}_R^2| + m_{\tilde{b}_R}^2 |\tilde{b}_R^2| + m_{\tilde{L}}^2 |\tilde{L}^2| + m_{\tilde{\tau}_R}^2 |\tilde{\tau}_R^2|, \quad (2.2)
 \end{aligned}$$

and the trilinear interactions between the sfermion and Higgs fields,

$$-\mathcal{L}_{\text{tril}} = \lambda A_\lambda H_u H_d S + \frac{1}{3} \kappa A_\kappa S^3 + h_t A_t \tilde{Q} H_u \tilde{t}_R^c - h_b A_b \tilde{Q} H_d \tilde{b}_R^c - h_\tau A_\tau \tilde{L} H_d \tilde{\tau}_R^c + \text{h.c.} \quad (2.3)$$

In an unconstrained NMSSM with non-universal soft terms at the GUT scale, the three SUSY breaking masses squared for H_u , H_d and S appearing in $\mathcal{L}_{\text{mass}}$ can be expressed in terms of their vevs through the three minimization conditions of the scalar potential. Thus, in contrast to the MSSM (where one has only two free parameters at the tree level, generally chosen to be the ratio of Higgs vacuum expectation values $\tan \beta$ and the mass of the pseudoscalar Higgs boson), the Higgs sector of the NMSSM is described by the six parameters

$$\lambda, \kappa, A_\lambda, A_\kappa, \tan \beta = \langle H_u \rangle / \langle H_d \rangle \text{ and } \mu_{\text{eff}} = \lambda \langle S \rangle. \quad (2.4)$$

One can choose sign conventions such that the parameters λ and $\tan \beta$ are positive, while the parameters κ , A_λ , A_κ and μ_{eff} can have both signs.

In addition to the above parameters of the Higgs sector, one needs to specify the soft SUSY breaking mass terms in eq. (2.2) for the scalars, the trilinear couplings in eq. (2.3) as well as the gaugino soft SUSY breaking mass parameters to describe the model completely,

$$-\mathcal{L}_{\text{gauginos}} = \frac{1}{2} \left[M_1 \tilde{B} \tilde{B} + M_2 \sum_{a=1}^3 \tilde{W}^a \tilde{W}_a + M_3 \sum_{a=1}^8 \tilde{G}^a \tilde{G}_a + \text{h.c.} \right]. \quad (2.5)$$

Clearly, in the limit $\lambda \rightarrow 0$ with finite μ_{eff} , the NMSSM turns into the MSSM with a decoupled singlet sector. Whereas the phenomenology of the NMSSM for $\lambda \rightarrow 0$ could still differ somewhat from the MSSM in the case where the lightest SUSY particle is the singlino (and hence with the possibility of a long lived next-to-lightest SUSY particle [30]), we will not consider this situation here.

2.2 The constrained NMSSM

As the number of input parameters is rather large, one can attempt to define a constrained model, hereafter called the cNMSSM, in which the soft SUSY-breaking parameters are fixed at the GUT scale, leading to only a handful of free input parameters to cope with. This approach is motivated by the fact that in many schemes for SUSY-breaking, the soft SUSY breaking parameters are universal at a very high energy scale. For example, in the cMSSM or mSUGRA scenario, one imposes a common gaugino mass $M_{1/2}$, a scalar mass m_0 and a trilinear coupling A_0 at M_{GUT} , leading to only four continuous free parameters (together with $\tan \beta$) and the sign of μ . The values of the numerous soft SUSY-breaking parameters at low energies are then obtained through the renormalization group evolution (RGE). Analogously, the cNMSSM allows to describe the entire sparticle spectrum, including the

chargino and neutralino sectors, in terms of a small number of free parameters. This is in contrast to the naïve procedure where one postulates universal sfermion masses directly at the weak scale.

In analogy to the cMSSM, one can impose unification of the soft SUSY-breaking gaugino masses, sfermion and Higgs masses and trilinear couplings at the scale M_{GUT} :

$$M_{1,2,3} \equiv M_{1/2}, \quad m_{\tilde{F}_i} = m_{H_i} \equiv m_0, \quad A_i \equiv A_0. \quad (2.6)$$

The cMSSM has two additional parameters (μ and B) beyond those in eq. (2.6), but the correct value of M_{Z^0} imposes one constraint (typically used to determine $|\mu|$) and B can be replaced by $\tan\beta$. Likewise, the fully constrained NMSSM has two additional parameters (λ and κ) beyond the three parameters in eq. (2.6), and the correct value of M_{Z^0} imposes one constraint. Hence, a priori the number of free parameters in the cMSSM and the fully constrained NMSSM is the same.

In principle, one could minimize the effective potential of the cNMSSM with respect to H_u , H_d and S , and determine the overall scale of the soft terms in eq. (2.6) from the correct value of M_{Z^0} : this approach has been pursued in ref. [2]. However, since $\tan\beta$ is then obtained as output (while the top quark Yukawa coupling h_t is an input), it becomes very difficult to obtain the correct value for $m_{t_{\text{top}}}$. Also, the numerical minimization of the effective potential including radiative corrections is quite computer-time consuming.

Therefore it is much more convenient to employ the analytic form of the three minimization conditions of the effective potential of the NMSSM: for given M_{Z^0} , $\tan\beta$, λ and all soft terms at the weak scale (except for m_S^2), these can be solved for $|\mu_{\text{eff}}|$ (or $|\langle S \rangle|$), κ and m_S^2 . As in the cMSSM, the sign of μ_{eff} can still be chosen at will; in some sense, the determination of the parameter B through the minimization conditions of the MSSM is replaced by the determination of κ in the NMSSM and the additional minimization w.r.t. $\langle S \rangle$ in the NMSSM leads to the determination of m_S^2 .

Assuming that all weak scale soft scalar masses (apart from m_S^2) arise from a unified scalar mass m_0^2 , the determined weak scale value of m_S^2 — when run up to the GUT scale — will not coincide with m_0^2 in general. One could confine oneself to regions in parameter space where the difference between m_S^2 and m_0^2 is negligibly small, but we found that the phenomenology of this fully constrained NMSSM hardly differs (at least in the Higgs sector on which we are focusing here) from the one of the cMSSM once present LEP constraints are imposed. Hence we will relax the hypothesis of complete unification of the soft terms in the singlet sector since the singlet could play a special rôle,² and we will allow for both $m_S^2 \neq m_0^2$ and $A_\kappa \neq A_0$ at M_{GUT} . Note that, although m_S^2 at M_{GUT} can be negative, this does *not* signal an instability of the potential: the direction $\langle S \rangle \rightarrow \infty$ is always protected by a quartic self coupling $\sim \kappa^2$, which leads to $\langle S \rangle \sim M_{\text{SUSY}}$ at the minimum of the potential.

In addition, for some of the benchmark points (see points P4 and P5 below), we will also relax the unification hypothesis for $m_{H_u}^2$ and $m_{H_d}^2$, in analogy to corresponding scenarios within the MSSM (called non-universal Higgs model or NUHM [32]). The hypothesis

²An example is when the singlet is mixed with or identified with a radion originating from a 5d brane world [31]; however, here we do not consider a Kaluza-Klein mass scale below M_{GUT} .

$A_\lambda = A_0$ will also be relaxed for point 5. Such points in parameter space can have additional unconventional properties, whose phenomenology should also be investigated.

2.3 The Higgs and SUSY spectra

Following the procedure employed by the routine NMSPEC within NMSSMTools [27], which calculates the spectra of the Higgs and SUSY particles in the NMSSM, a point in the parameter space of the cNMSSM is defined by the soft SUSY breaking terms at M_{GUT} (except for the parameter m_S^2), $\tan\beta$ at the weak scale, λ at the SUSY scale (defined as an average of the first generation squark masses) and the sign of the parameter μ_{eff} . The parameters κ , m_S^2 and $|\mu_{\text{eff}}|$ are determined at the SUSY scale in terms of the other parameters through the minimization equations of the scalar potential.

The RGEs for the gauge and Yukawa couplings have to be integrated from the weak scale up to M_{GUT} (defined by the unification of the gauge couplings g_1 and g_2), and the RGEs for the soft terms from M_{GUT} down to the weak scale. Since κ and m_S^2 are computed at the weak scale, and threshold effects for the gauge and Yukawa couplings depend on the soft terms at the SUSY scale, some iterations are necessary in order to satisfy the desired boundary conditions for the soft terms at the GUT scale, but usually the procedure converges quite rapidly (at least for not too large values of $\tan\beta$ or λ). For the most relevant SM parameters, the strong coupling and the top/bottom quark masses, we chose $\alpha_s(M_{Z^0}) = 0.1172$, $m_b(m_b)^{\overline{\text{MS}}} = 4.214$ GeV and $m_{\text{top}}^{\text{pole}} = 171.4$ GeV.

After RGE running is completed, the gluino, chargino, neutralino and sfermion masses are computed including dominant one loop corrections to their pole masses. The lightest scalar Higgs pole mass is determined to the following accuracy (in a notation where its tree level mass is proportional to $\mathcal{O}(g^2)$, where g denotes any of the electroweak gauge couplings): one loop corrections of $\mathcal{O}(h_{t,b}^4)$ and $\mathcal{O}(h_{t,b}^2 g^2)$ are computed exactly; one loop corrections of the orders g^4 , $g^2 \lambda^2$, λ^4 , $g^2 \kappa^2$, κ^4 and $\lambda^2 \kappa^2$ include only terms involving large logarithms $\ln(M_i^2/M_{Z^0}^2)$, where M_i are potentially large Higgs or sparticle masses, while two loop corrections of $\mathcal{O}(h_{t,b}^6)$ and $\mathcal{O}(h_{t,b}^4 \alpha_s)$ include only terms with two powers of large logs.

Once the pole masses are known, all Higgs decay branching ratios into SM and SUSY particles (an adaptation of the decays in the MSSM [33] to the NMSSM case) are determined including dominant (mainly QCD) radiative corrections [34], as well as sparticle loops which contribute to the couplings of a neutral Higgs to two photons or gluons [35].

When the spectrum and the couplings of the Higgs and SUSY particles are computed, available Tevatron and LEP constraints are checked. The results of the four LEP collaborations, combined by the LEP Higgs working group, are included [36]. More specifically, the following experimental constraints are taken into account:

- (i) The masses of the neutralino as well as their couplings to the Z^0 boson are compared with the LEP constraints from direct searches and from the invisible Z^0 boson width;
- (ii) Direct bounds from LEP and Tevatron on the masses of the charged particles (h^\pm , χ^\pm , \tilde{q} , \tilde{l}) and of the gluino are taken into account;

- (iii) Constraints on the Higgs production rates from all channels studied at LEP. These include in particular $Z^0 h_i^0$ production, h_i^0 being any of the CP-even Higgs particles, with all possible two body decay modes of h_i^0 (into b quarks, τ leptons, jets, photons or invisible), and all possible decay modes of h_i^0 of the form $h_i^0 \rightarrow a_j^0 a_j^0$, a_j^0 being any of the CP-odd Higgs particles, with all possible combinations of a_j^0 decays into b quarks, c quarks, τ leptons and jets. Also considered is the associated production mode $e^+e^- \rightarrow h_i^0 a_j^0$ with, possibly, $h_i^0 \rightarrow a_j^0 a_j^0$. (In practice, for our purposes, only combinations of $i = 1, 2$ with $j = 1$ are phenomenologically relevant.)

We stress that light Higgs scalars (with $M_{h_{1,2}^0} \lesssim 114$ GeV) can still be allowed by LEP constraints, if either (i) the Z^0 - Z^0 - $h_{1,2}^0$ coupling is heavily suppressed (if, for instance, the state h_i^0 is dominantly a gauge singlet); or (ii) $M_{a_1^0} \lesssim 11$ GeV such that the state $h_{1,2}^0$ decays dominantly into $a_1^0 a_1^0$ states, but the $b\bar{b}$ decay of a_1^0 is impossible; constraints from the decays $h_{1,2}^0 \rightarrow a_1^0 a_1^0 \rightarrow 4\tau$ allow for $M_{h_{1,2}^0}$ down to ~ 86 GeV. It is important to note, however, that LEP constraints are implemented only for individual processes, and that combinations of different processes could potentially rule out seemingly viable scenarios.

Finally, experimental constraints from B physics [37] such as the branching ratios of the rare decays $\text{BR}(B \rightarrow X_s \gamma)$, $\text{BR}(B_s \rightarrow \mu^+ \mu^-)$ and $\text{BR}(B^+ \rightarrow \tau^+ \nu_\tau)$ and the mass differences ΔM_s and ΔM_d , are also implemented; compatibility of each point in parameter space with the current experimental bounds is required at the two sigma level. (Notably constraints from $\text{BR}(B_s \rightarrow \mu^+ \mu^-)$ can rule out too light pseudoscalars.)

The new features of the NMSSM have an impact on the properties of the lightest neutralino as a dark matter candidate [17]. For instance, given the presence of a fifth neutralino (singlino), the composition of the annihilating WIMPs can be significantly different from those within the MSSM in wide regions of the parameter space. In particular, the LSP can have a large singlino component, in which case one has new couplings of the LSP to singlet-like Higgs states whose mass can be substantially lighter than the Higgs states within the MSSM. In the presence of light h_1^0 and a_1^0 states, there are new channels through which neutralino annihilation can occur in the NMSSM: $Z^0 h_1^0$, $h_1^0 h_1^0$, $h_1^0 a_1^0$ and $a_1^0 a_1^0$, either via s -channel Z^0 , a_i^0 , h_i^0 or t -channel neutralino exchange [17]. Among the proposed benchmark points, we will find examples of these new features.

Technically, the relic abundance of the NMSSM dark matter candidate χ_1^0 is evaluated via a link to MicrOMEGAS [38]. All the relevant cross sections for the lightest neutralino annihilation and co-annihilation are computed. The density evolution equation is numerically solved and the relic density of χ_1^0 is calculated. The result is compared with the ‘‘WMAP’’ constraint $0.094 \lesssim \Omega_{\text{CDM}} h^2 \lesssim 0.136$ at the 2σ level [39].

3. The benchmark points

As already mentioned in the introduction, in view of the upcoming LHC, quite some work has been dedicated to probe the NMSSM Higgs sector over the recent years. In the NMSSM, two different types of scenarios have been pointed out, depending on whether Higgs-to-Higgs decays are kinematically allowed or forbidden. Within the first category,

where Higgs-to-Higgs decays are kinematically allowed and can be dominant, there are two possibilities, each associated with light scalar/pseudoscalar Higgs states:

(i) The lightest pseudoscalar Higgs boson a_1^0 is rather light, $M_{a_1^0} \lesssim 40\text{-}50$ GeV, and the lightest CP-even Higgs particle h_1^0 has enough phase space for the decay into two pseudoscalar Higgs particles, $h_1^0 \rightarrow a_1^0 a_1^0$, to be kinematically accessible. In this case, the branching ratio for the decay $h_1^0 \rightarrow a_1^0 a_1^0$ is typically very large and this new decay channel is the dominant one. Concerning the lightest pseudoscalar decays, there are two further possibilities: either $M_{a_1^0} \gtrsim 10$ GeV and the a_1^0 boson decays into a pair of b quarks or a pair of tau leptons (with the former decay being in general dominant), leading to $h_1^0 \rightarrow a_1^0 a_1^0 \rightarrow 4\tau, 4b$ and $2\tau 2b$ final states, or $M_{a_1^0} \lesssim 10$ GeV and the dominant decay mode of the a_1^0 boson is into a pair of tau leptons, leading to $h_1^0 \rightarrow a_1^0 a_1^0 \rightarrow 4\tau$ final states. In this latter scenario, one can still distinguish two situations for the h_1^0 boson: its mass is either close to its theoretical upper limit of 130 GeV (within the cNMSSM), or to the lower limit of 90 GeV, still in agreement with constraints from Higgs boson searches at LEP2.

(ii) The lightest CP-even Higgs boson is relatively light, $M_{h_1^0} \lesssim 50$ GeV, and decays into $b\bar{b}$ pairs. (The situation where $M_{h_1^0} \lesssim 10$ GeV, such that the latter channel is kinematically closed and the decay $h_1^0 \rightarrow \tau^+ \tau^-$ dominates, is very constrained by LEP data.) In this case, the next-to-lightest CP even Higgs boson h_2^0 is SM-like, and has a mass below ~ 150 GeV. Even so, it can still decay into two h_1^0 bosons leading to the final topologies $h_2^0 \rightarrow h_1^0 h_1^0 \rightarrow 4\tau, 2\tau 2b$ and $4b$, the latter final state being largely dominating.

The second category of scenarios, where Higgs-to-Higgs decays are suppressed, includes regions of the parameter space where all Higgs particles are relatively light with masses in the range 90-200 GeV. This opens the possibility of producing all the five neutral and the charged Higgs bosons at the LHC. This scenario is similar to the so-called ‘‘intense coupling regime’’ of the MSSM [40, 41] (which has been shown to be rather difficult to be fully covered at the LHC), but with two more neutral Higgs particles.

In the context of the general NMSSM (without unification constraints), there exists also a decoupling regime in which all the Higgs bosons are heavy and decouple from the spectrum except for one CP-even Higgs particle, whose mass can be up to ~ 140 GeV [42]. Such decoupling scenarios are difficult to realize within the cNMSSM and, in any case, they would be very difficult to disentangle from an MSSM ‘‘ M_h^{max} ’’-scenario at the LHC, since the separation of the two would require a determination of the parameter $\tan\beta$ which is notoriously difficult to achieve at the LHC.

In the following, we propose five benchmark points of the NMSSM parameter space, P1 to P5, in which the above discussed scenarios are realized. Each point is representative of distinctive NMSSM features: points P1 to P3 exemplify scenarios where h_1^0 decays into light pseudoscalar states (decaying, in turn, into $b\bar{b}$ or $\tau^+ \tau^-$ final states), P4 illustrates the NMSSM possibility of a very light h_1^0 , while P5 corresponds to the case where all Higgs bosons are rather light. In all cases, the input parameters as well as the resulting Higgs masses and the relevant decay information are given in table 1. In the subsequent table 2, we summarize the most relevant features of each point regarding the cosmological relic density. In table 1, R_i denote the couplings of the corresponding Higgs scalars to gauge bosons (normalized with respect to the coupling of a SM Higgs scalar), $t_i^{(\prime)}$ and $b_i^{(\prime)}$ denote




Figure 1: Allowed values of A_κ , $M_{h_1^0}$ and $M_{a_1^0}$ (in GeV) as well as $\Omega_{\text{CDM}}h^2$ as a function of λ . We take $m_0 = 174$ GeV, $M_{1/2} = 500$ GeV, $A_0 = -1500$ GeV and $\tan\beta = 10$; $m_{\tilde{g}}^2$ is determined from the electroweak symmetry breaking conditions.

the corresponding normalized couplings to up- and down-quarks. In table 2, N_{1i} denote the components of the LSP in the basis (bino, wino, higgsino-down, higgsino-up, singlino).

As discussed in ref. [27], the scenarios where the pseudoscalar Higgs boson is light and the decay $h_1^0 \rightarrow a_1^0 a_1^0$ is dominant, can be realized within the cNMSSM with nearly universal soft terms at the GUT scale, the exception being the parameters $m_{\tilde{g}}^2$ and A_κ , which are chosen independently. A corresponding region in parameter space, containing P1, P2 and P3, is shown in figure 1, where we chose $m_0 = 174$ GeV, $M_{1/2} = 500$ GeV, $A_0 = -1500$ GeV, $\tan\beta = 10$ and the sign of μ_{eff} is positive. As a function of the parameter λ , we show the values of A_κ that are allowed by LEP constraints, the masses of the h_1^0 and a_1^0 bosons (where larger values of $M_{a_1^0}$ correspond to larger values of the trilinear coupling $|A_\kappa|$), and the dark matter relic density $\Omega_{\text{CDM}}h^2$. The benchmark points P1, P2 and P3 discussed below are chosen at the upper and lower boundaries of this region of the cNMSSM parameter space. The lower limit on the pseudoscalar a_1^0 mass, $M_{a_1^0} \gtrsim 8$ GeV, in the last plot of figure 1 originates from constraints from B physics; see ref. [37] for details.

In figure 2, we show the obtained results from a scan in the $[\lambda, A_\kappa]$ parameter space (with the values of the other parameters fixed to those used in figure 1) for the masses of the CP-even h_1^0 and CP-odd a_1^0 states, as well as of the branching ratios of the decays $h_1^0 \rightarrow a_1^0 a_1^0$ and $a_1^0 \rightarrow \tau^+ \tau^-$. In the two upper frames for the Higgs masses, a scan is




Figure 2: Scans in the $[\lambda, A_\kappa]$ parameter space for the h_1^0, a_1^0 masses, and the branching ratios for the decay modes $h_1^0 \rightarrow a_1^0 a_1^0$ and $a_1^0 \rightarrow \tau^+ \tau^-$. The two upper frames include P1 to P3 while the remaining frames include only P1 and P2; other parameters as in figure 1.

performed over the same $[\lambda, A_\kappa]$ range as in figure 1, which therefore includes the three points P1, P2 and P3. In the lower frames for the Higgs masses and for the branching ratios, a finer scan zooms on the $[\lambda, A_\kappa]$ range which involves only the two points P1 and P2. Theoretically excluded are regions with an unstable scalar potential (negative masses squared for a_1 or h_1). Experimental constraints originate from the LEP bound on m_{h_1} .

A very light h_1^0 state (and, thus, with the decay $h_2^0 \rightarrow h_1^0 h_1^0$ being the dominant channel) is represented by P4, which can be obtained once one relaxes the universality conditions on the soft SUSY-breaking Higgs mass terms, $m_{H_d} \neq m_{H_u} \neq m_0$. A scenario as in P5, in which all NMSSM Higgs bosons are light, is possible if one allows additionally for $A_\lambda \neq A_0$.

For all benchmark points the numerical value of the pseudoscalar Higgs boson mass $M_{a_1^0}$ is quite sensitive not only to the NMSSM input parameters (notably to the trilinear coupling A_κ), but also to the employed SM parameters and to the precision with which radiative corrections are computed. Since the numerical values of $M_{a_1^0}$ are phenomenologically much more important than, for instance, A_κ at M_{GUT} , we consider the benchmark points to be defined in terms of $M_{a_1^0}$ rather than in terms of A_κ at M_{GUT} . Next, we summarize the most relevant phenomenological properties of the benchmark points.

In the first two points P1 and P2, the lightest CP-even Higgs boson has a mass of

$M_{h_1^0} \simeq 120$ GeV and is SM-like, as reflected by the corresponding couplings to gauge bosons R_1 , top quarks t_1 and bottom quarks b_1 , which are almost equal to unity: see table 1. The lightest CP-odd Higgs boson has masses of 40.5 GeV and 9.1 GeV, which are obtained by choosing A_κ at M_{GUT} as -33.9 GeV and -33.4 GeV, respectively; the remaining parameters of the Higgs sector at M_{SUSY} , like $\kappa, A_\lambda, A_\kappa, \mu_{\text{eff}}$ and the gaugino mass parameter M_2 , are given in the second part of table 1.

In both cases P1 and P2, the decay channel $h_1^0 \rightarrow a_1^0 a_1^0$ is largely dominating with a branching rate very close to 90%, while the decays $h_1^0 \rightarrow b\bar{b}$ and $\tau^+\tau^-$ are suppressed by an order of magnitude as compared to the SM. The most relevant difference between the two scenarios concerns the mass and decays of the lightest pseudoscalar state. In P1 the pseudoscalar a_1^0 decays into b quarks and τ leptons, with branching fractions of $\sim 90\%$ and $\sim 10\%$, respectively. On the other hand, in P2 the pseudoscalar a_1^0 with its mass $M_{a_1^0} \simeq 9.1$ GeV decays dominantly into $\tau^+\tau^-$ pairs, with a rate close to 90%.

For point P3, the same inputs of points P1 and P2 are chosen except for the A_κ and λ parameters, which are now varied as to have a lighter h_1^0 state. The particular value for A_κ leads again to a pseudoscalar Higgs boson below $b\bar{b}$ threshold, $M_{a_1^0} \simeq 9.8$ GeV, which decays almost exclusively into $\tau^+\tau^-$ final states. The difference between P3 and P2 is the lightest CP-even Higgs boson h_1^0 , which has a mass $M_{h_1^0} \simeq 90$ GeV, lower than in scenarios P1 and P2. In this case, and although h_1^0 is still SM-like, exhibiting couplings to gauge bosons, top and bottom quarks that are very close to those of the SM Higgs boson, it decays nevertheless almost exclusively into a_1^0 pairs, with a rate close to 100%. Another difference between P2 and P3 is that in the former case the decay mode $h_1^0 \rightarrow a_1^0 Z^0$ is kinematically possible but the branching ratio for this new interesting channel that is not listed in the table is rather small, $\text{BR}(h_1^0 \rightarrow a_1^0 Z^0) \sim 3\%$.

Note that in all these first three points, the heaviest neutral Higgs particles h_2^0, h_3^0 and a_2^0 , as well as the charged Higgs states h^\pm , all have masses close to or above 1 TeV. The main decay modes are into $b\bar{b}$ and $t\bar{t}$ for the neutral and tb for the charged states, as $\tan\beta$ is not too large and the $t\bar{t}$ -Higgs couplings are not very strongly suppressed, while the branching fractions for the neutral Higgs-to-Higgs decays, in particular the channels $h_2^0 \rightarrow h_1^0 h_1^0$ and $h_2^0 \rightarrow a_1^0 a_1^0$, are very tiny, not exceeding the permille level.

Regarding the properties of the DM candidate, P1, P2 and P3 exhibit a lightest neutralino which is bino-like, and whose mass $m_{\chi_1^0} \simeq 208$ GeV. In all three cases, the correct relic density, $\Omega_{\text{CDM}} h^2 \simeq 0.1$, is achieved through the co-annihilation with the $\tilde{\tau}_1$ slepton, which has a mass comparable to that of the LSP, $m_{\tilde{\tau}_1} \simeq 213, 213$ and 215 GeV, respectively, for P1, P2 and P3. (In fact, the requirement of the correct relic density determined the choice of m_0 .) In each benchmark scenario, the dominant co-annihilation channels are thus $\chi_1^0 \tilde{\tau}_1 \rightarrow \gamma\tau$ ($\sim 33\%$) and $\chi_1^0 \tilde{\tau}_1 \rightarrow Z^0\tau$ ($\sim 10\%$).

The benchmark point P4 corresponds to a scenario in which the CP-even boson h_1^0 is very light, $M_{h_1^0} \simeq 32$ GeV, and singlet-like, with a singlet component above 99%. The lightest scalar Higgs particle predominantly decays into $b\bar{b}$ pairs, with $\text{BR}(h_1^0 \rightarrow b\bar{b}) \simeq 92\%$, and to a smaller extent into τ pairs with $\text{BR}(h_1^0 \rightarrow \tau^+\tau^-) \simeq 7\%$. The CP-even h_2^0 boson has a mass of $M_{h_2^0} \simeq 123$ GeV and is SM-like, with normalized couplings to W/Z^0 and t/b

Point	P1	P2	P3	P4	P5
GUT/input parameters					
$\text{sign}(\mu_{\text{eff}})$	+	+	+	-	+
$\tan\beta$	10	10	10	2.6	6
m_0 (GeV)	174	174	174	775	1500
$M_{1/2}$ (GeV)	500	500	500	760	175
A_0	-1500	-1500	-1500	-2300	-2468
A_λ	-1500	-1500	-1500	-2300	-800
A_κ	-33.9	-33.4	-628.568	-1170	60
NUHM: M_{H_d} (GeV)	-	-	-	880	-311
NUHM: M_{H_u} (GeV)	-	-	-	2195	1910
Parameters at the SUSY scale					
λ (input parameter)	0.1	0.1	0.4	0.53	0.016
κ	0.11	0.11	0.31	0.12	-0.0029
A_λ (GeV)	-982	-982	-629	-510	45.8
A_κ (GeV)	-1.63	-1.14	-11.4	220	60.2
M_2 (GeV)	392	392	393	603	140
μ_{eff} (GeV)	968	968	936	-193	303
CP even Higgs bosons					
$m_{h_1^0}$ (GeV)	120.2	120.2	89.9	32.2	90.8
R_1	1.00	1.00	0.998	0.036	-0.315
t_1	1.00	1.00	0.999	0.084	-0.305
b_1	1.018	1.018	0.975	-0.289	-0.645
$\text{BR}(h_1^0 \rightarrow b\bar{b})$	0.072	0.056	7×10^{-4}	0.918	0.895
$\text{BR}(h_1^0 \rightarrow \tau^+\tau^-)$	0.008	0.006	7×10^{-5}	0.073	0.088
$\text{BR}(h_1^0 \rightarrow a_1^0 a_1^0)$	0.897	0.921	0.999	0.0	0.0
$m_{h_2^0}$ (GeV)	998	998	964	123	118
R_2	-0.0018	-0.0018	0.005	0.999	0.927
t_2	-0.102	-0.102	-0.095	0.994	0.894
b_2	10.00	10.00	9.99	1.038	2.11
$\text{BR}(h_2^0 \rightarrow b\bar{b})$	0.30	0.30	0.14	0.080	0.87
$\text{BR}(h_2^0 \rightarrow t\bar{t})$	0.11	0.11	0.046	0.0	0.0
$\text{BR}(h_2^0 \rightarrow a_1^0 Z^0)$	0.22	0.22	0.72	0.0	0.0
$m_{h_3^0}$ (GeV)	2142	2142	1434	547	174
CP odd Higgs bosons					
$m_{a_1^0}$ (GeV)	40.5	9.1	9.8	185	99.7
t'_1	0.0053	0.0053	0.0142	0.0513	-0.00438
b'_1	0.529	0.528	1.425	0.347	-0.158
$\text{BR}(a_1^0 \rightarrow b\bar{b})$	0.91	0.	0.	0.61	0.91
$\text{BR}(a_1^0 \rightarrow \tau^+\tau^-)$	0.085	0.88	0.81	0.071	0.090
$m_{a_2^0}$ (GeV)	1003	1003	996	546	170
Charged Higgs boson					
m_{h^\pm} (GeV)	1005	1005	987	541	188

Table 1: Input and output parameters for the five benchmark NMSSM points.

states close to unity. However, since $M_{h_2^0} > 2M_{h_1^0}$, it mostly decays into two h_1^0 bosons, with $\text{BR}(h_2^0 \rightarrow h_1^0 h_1^0) \simeq 88\%$, and the dominant SM-like $b\bar{b}$ decay mode occurs only at a rate smaller than 10%. The lightest CP-odd particle is not very heavy, $M_{a_1^0} = 185$ GeV, and decays mostly into fermion pairs $\text{BR}(a_1^0 \rightarrow b\bar{b}) \sim 61\%$ and $\text{BR}(a_1^0 \rightarrow \tau^+\tau^-) \sim 7\%$.

Point	P1	P2	P3	P4	P5
Dark matter					
LSP (χ_1^0) mass (GeV)	208	208	208	100	70.4
N_{11}	0.999	0.999	0.999	-0.039	0.977
N_{12}	-0.008	-0.008	-0.009	0.043	-0.098
N_{13}	0.048	0.048	0.050	-0.028	0.178
N_{14}	-0.015	-0.015	-0.016	0.405	-0.068
N_{15}	0	0	0.003	0.912	-0.003
$\Omega_{\text{CDM}} h^2$	0.097	0.097	0.126	0.099	0.106

Table 2: LSP properties and relic density for the five benchmark NMSSM points.

The other dominant decay is the interesting channel $a_1^0 \rightarrow h_1^0 Z^0$ which has a branching ratio of the order of 30%. Finally, the heaviest CP even h_3^0 and CP-odd a_2^0 states and the charged h^\pm particles have masses in the 500 GeV range and will mostly decay, as $\tan\beta$ is small, into $t\bar{t}/tb$ final states for the neutral/charged states. All these features make the phenomenology of point P4 rather different from that of points P1 to P3 discussed above.

To achieve a correct cosmological relic density, the common sfermion and gaugino mass parameter m_0 and $M_{1/2}$ at the GUT scale are set close to 1 TeV. At the SUSY scale, one thus finds a higgsino-singlino-like neutralino LSP, whose mass is $m_{\chi_1^0} \sim 100$ GeV. LSP annihilation essentially occurs via two channels: $\chi_1^0 \chi_1^0 \rightarrow W^+ W^-$ (60%) and $\chi_1^0 \chi_1^0 \rightarrow Z^0 h_1^0$ (20%), mediated by s -channel Z^0 and Higgs boson exchange.

Finally, point P5 is characterized by having all Higgs particles relatively light with masses in the range 90 to 190 GeV. Here, the small value for the coupling λ leads to a small value $\kappa \simeq -0.003$. The three CP-even Higgs bosons with masses of 91, 118 and 174 GeV, respectively, share the couplings of the SM Higgs boson to SM gauge bosons with the dominant component being taken by the h_2^0 state. The reduced couplings of the state h_3^0 for this point, not given in the table 1, are $R_3 = -0.20$, $t_3 = -0.37$ and $b_3 = 5.7$. The pseudoscalar Higgs bosons have masses $M_{a_1^0} \simeq 100$ GeV and $M_{a_2^0} \simeq 170$ GeV, while the charged Higgs particle is the heaviest one with a mass $M_{h^\pm} \simeq 188$ GeV. Here, all the neutral Higgs-to-Higgs decays are kinematically forbidden. This is also the case of neutral Higgs decays into lighter Higgs states with opposite parity and gauge bosons. The only non-fermionic two-body Higgs decays are thus $h^\pm \rightarrow W h_1^0$ and $h_3^0 \rightarrow WW$, but as the involved Higgs-gauge boson couplings are small, the branching ratios are tiny.

In this last point P5, the LSP with a mass $m_{\chi_1^0} \sim 70$ GeV, is a bino-like neutralino but has a small non-negligible higgsino component. The correct cosmological relic density $\Omega_{\text{CDM}} h^2 \simeq 0.1$ is achieved through the exchange of Higgs bosons in the s -channel in the annihilation processes $\chi_1^0 \chi_1^0 \rightarrow b\bar{b}, \tau^+ \tau^-$ which occur at a rate of $\sim 90\%$ and 10% , respectively, i.e. comparable to the Higgs decay branching ratios.

More detailed properties of the benchmark points such as more branching ratios of the Higgs scalars are available on the web site of ref. [27].

4. Implications for the LHC

4.1 Dominant production processes

In the cases discussed here, at least one CP-even Higgs particle h_i^0 has strong enough couplings to massive gauge bosons or top/bottom quarks, i.e. $R_i, t_i/b_i \sim 1$, to allow for the production at the LHC in one of the main channels which are advocated for the search of the SM Higgs particle [20]:

- (i) gluon-gluon fusion, $gg \rightarrow h_i^0$, occurring through a loop of heavy top quarks which couple strongly to the Higgs boson,
- (ii) vector boson fusion, $qq \rightarrow qqW^*W^*, qqZ^{0*}Z^{0*} \rightarrow qqh_i^0$, which leads to two forward jets and a centrally decaying Higgs boson,
- (iii) the Higgs-strahlung processes, $q\bar{q}' \rightarrow Wh_i^0$ and $q\bar{q} \rightarrow Z^0h_i^0$, which lead to a Higgs and a massive gauge boson in the final state,
- (iv) associated production with heavy quark pairs $q\bar{q}/gg \rightarrow Q\bar{Q}h_i^0$, with $Q = t, b$.

In scenarios P1 to P3, this CP-even particle is the h_1^0 boson which has $R_1 \simeq t_1 \simeq b_1 \simeq 1$, but which decays predominantly into a pair of light pseudoscalar Higgs particles, $h_1^0 \rightarrow a_1^0 a_1^0$, which subsequently decay into light fermion pairs, $a_1^0 \rightarrow b\bar{b}$ and $\tau^+\tau^-$. In scenario P4, this particle is the h_2^0 boson which decays most of the time into a pair of h_1^0 particles, $h_2^0 \rightarrow h_1^0 h_1^0$, which again decay into light fermion pairs. In these four cases, the backgrounds for the modes $gg \rightarrow h_i^0 \rightarrow 4f$ and $qq/gg \rightarrow t\bar{t}h_i^0 \rightarrow t\bar{t} + 4f$ with $f = b, \tau$, will be extremely large and only the vector boson fusion (owing to the forward jet tagging) and possibly the Higgs-strahlung (due to the leptons coming from the decays of the gauge bosons) can probably be viable at the LHC. Note that in the case of scenario P4, one can try taking advantage of the relatively light a_1^0 and of the large rate for the interesting and clean decay mode $a_1^0 \rightarrow h_1^0 Z^0$, which occurs at the level of 2% if the Z^0 boson decays into electrons and muons. However, the cross sections for a_1^0 production in the $gg \rightarrow a_1^0$ and $gg/q\bar{q} \rightarrow t\bar{t}a_1^0$ processes, which are the dominant ones for the production of the MSSM CP-odd Higgs boson [20], are very small as the reduced $a_1^0 tt$ coupling is very tiny, $t_1' \sim 0.05$. One would have then to rely on associated production of the a_1^0 state with a CP-even Higgs boson, such as $q\bar{q} \rightarrow Z^{0*} \rightarrow a_1^0 h_1^0$, but the cross section is not expected to be very large.

In scenario P5, the particle which has couplings to gauge bosons and top quarks that are close to those of the SM Higgs boson is the h_2^0 boson which decays into $b\bar{b}$ and $\tau^+\tau^-$ final states with branching ratios close to 90% and 10%, respectively. Here again, the gg fusion and presumably associated production with top quarks cannot be used since the interesting decays such as $h_2^0 \rightarrow WW^*, Z^0 Z^{0*}$ and $\gamma\gamma$ are suppressed when compared to the SM case (by at least a factor of four as $b_2 \sim 2$). Thus, in this case, only the channels $qq \rightarrow qqh_2^0 \rightarrow qq\tau^+\tau^-$ and possibly $q\bar{q}' \rightarrow Wh_2^0 \rightarrow \ell\nu b\bar{b}$ seem feasible. The state h_1^0 has still non-negligible couplings to gauge bosons and top quarks which lead to cross sections that are “only” one order of magnitude smaller than in the SM. Since here, only the decays $h_1^0 \rightarrow b\bar{b}$ (90%) and $\tau^+\tau^-$ (10%) are again relevant, the only processes which can be used

are the vector boson fusion and Higgs-strahlung processes discussed above. Even so, one needs a luminosity 10 times larger to have the same event samples as in the SM. Note that for this point, one can also consider associated CP-even and CP-odd Higgs production, $q\bar{q} \rightarrow Z^{0*} \rightarrow h_i^0 a_i^0$ but the cross sections are still small. Finally, one should note that for this point, the value of $\tan \beta$ is moderate and thus, the charged Higgs coupling to tb states is the smallest possible, and does not guarantee the detection of the h^\pm bosons.

4.2 Summary of available studies

Let us now briefly summarize the few detailed studies (possibly including Monte Carlo simulations) of the LHC potential for the NMSSM Higgs sector that have been performed for scenarios close to the ones discussed here.³

In scenarios with a very light pseudoscalar Higgs boson, preliminary LHC studies focused on the $qq \rightarrow qqWW, qqZ^0Z^0 \rightarrow qqh_1^0 \rightarrow qqa_1^0a_1^0$ detection mode, i.e. via vector boson fusion (VBF) with forward/backward jet tagging [6]. The hope was that such NMSSM specific scenarios would be visible, particularly if the lightest CP-odd Higgs boson mass allowed for abundant $a_1^0a_1^0 \rightarrow b\bar{b}\tau^+\tau^-$ decays, with both τ -leptons being detected via their e, μ leptonic decays. At high luminosity, the VBF signal may be detectable at the LHC as a bump in the tail of a rapidly falling mass distribution of the $2\tau 2j$ system (without b tagging). However, this procedure relies on the background shape to be accurately predictable. These analyses were based on Monte Carlo (MC) event generation via the SUSY routines of the HERWIG code [43] and the toy detector simulation GETJET. Further analyses based on the PYTHIA generator [44] and the more adequate ATLAS detector simulation ATLFast [45] found that the original selection procedures may need improvement in order to extract a signal.

Also considered was the Higgs-strahlung process with the $q\bar{q}' \rightarrow W^* \rightarrow W^\pm h_1^0 \rightarrow W^\pm a_1^0 a_1^0$ signature, exploiting leptonic decays of gauge bosons with a subleading component from $q\bar{q} \rightarrow Z^{0*} \rightarrow Z^0 h_1^0 \rightarrow Z^0 a_1^0 a_1^0$ and, more marginally, associated production with top quarks, $q\bar{q}, gg \rightarrow t\bar{t}h_1^0 \rightarrow t\bar{t}a_1^0 a_1^0$ [12]. Regardless of the decay modes of the pseudoscalar Higgs bosons, it has been shown in this parton-level analysis, but in which the efficiency to trigger on the signal is included, that at least the Higgs-strahlung process $q\bar{q} \rightarrow Wh_1^0$ should be taken into account along with the VBF process to improve the overall signal efficiency. This is particularly true for h_1^0 masses below ~ 90 GeV where the Higgs-strahlung cross section exceeds that from VBF production. In another parton-level study [16], Higgs-strahlung with $h_1^0 \rightarrow a_1^0 a_1^0 \rightarrow 4b$ was considered, and it was claimed that with a charged lepton for the W boson and the fully tagged $4b$ final state with Higgs mass reconstruction, the signal could be disentangled from the background. Nevertheless, these results need to be confirmed by MC and detector simulations.

Note that in ref. [13], the $h_1^0 \rightarrow 4\gamma$ final state topology was advocated to be useful for the LHC at very high luminosity if the branching ratio of the decay exceeds the level of 10^{-4} . However, in general, this decay has a smaller branching ratio and again, a detailed simulation which takes into account the experimental environment is lacking here. The

³The prospects for the Tevatron have been discussed in refs. [11, 13, 15] and seem quite dim.

scope of other decays, such as $a_1^0 a_1^0 \rightarrow jjjj$ and $jj \tau^+ \tau^-$ where j represents a light quark jet, is expected to be very much reduced, while the possibilities from $a_1^0 a_1^0 \rightarrow 4\tau$ are currently being explored, in both vector boson and Higgs-strahlung production processes.

In scenarios with a very light scalar Higgs boson such as our point P4, constraints from Higgs searches at LEP do not allow for h_1^0 masses below about 10 GeV, hence the main decay mode would be $h_1^0 \rightarrow b\bar{b}$ while the decay $h_1^0 \rightarrow \tau^+ \tau^-$ would have a branching ratio of the order of 7-8%. The studies discussed above for very light pseudoscalar Higgs bosons but with the SM-like Higgs boson being the h_2^0 state, which then decays into two h_1^0 bosons, can therefore be adapted to this case. In particular, the situation for point P4 would be very similar to that of point P1 as the production cross sections, the decay branching ratios and the masses of the involved primary and secondary Higgs bosons are very similar.

Ref. [46] considered a particularly challenging NMSSM scenario, with a Higgs spectrum very similar to that of the MSSM, i.e., nearly degenerate (doublet dominated), heavy charged, scalar and pseudoscalar states and a light scalar Higgs boson at around 120–140 GeV, but including an additional singlet-dominated scalar and a pseudoscalar; such a scenario is somewhat similar to our point P4. Despite of having a reasonably large production cross-sections at the LHC, this light Higgs boson would be difficult to observe since its main hadronic decays cannot be easily disentangled from the QCD backgrounds.

In addition, some studies performed in the CP-violating MSSM, in which the h_1^0 boson can also be very light with reduced couplings to gauge bosons, can also be adapted to this context. Note however, that in the case of the CP-violating MSSM, one would also have a light h^\pm boson, while in the NMSSM this state can be very heavy.

Finally, *in scenarios in which all NMSSM Higgs bosons are relatively light* as in point P5, no detailed analysis has been performed. To our knowledge, the only study that is available is the one performed in ref. [9] in which the ATLAS and CMS signal significances for the MSSM, assuming a high luminosity of 300 fb^{-1} , were rescaled to take into account the reduced couplings of the various Higgs particles to gauge bosons and top quarks. The effect of almost overlapping resonances, new decay modes or production channels not present in the MSSM analyses of the ATLAS and CMS collaborations have not been considered. In fact, the situation in this scenario looks similar to that of the intense coupling regime of the MSSM [40], in which the three neutral Higgs bosons have masses in the range 100-140 GeV. It was shown in the detailed simulation of ref. [41] that while it would be very difficult to resolve all Higgs resonances, it would be at least possible to detect one or two Higgs states. However, in the latter case, the value of $\tan\beta$ is assumed to be large, leading to strongly enhanced cross sections in the gg fusion and $b\bar{b}$ Higgs processes, while in the NMSSM one could have relatively moderate $\tan\beta$ values as in P5 and thus, smaller production rates.

4.3 ATLAS strategy for NMSSM $h_1^0 \rightarrow a_1^0 a_1^0$ searches

At the ATLAS collaboration [47], current efforts to find suitable search channels for special NMSSM phenomenological scenarios are being focused on the vector boson fusion production of a scalar Higgs boson with relatively low mass and subsequent decay via a pair of pseudoscalar Higgs bosons into four τ -leptons, $h_1^0 \rightarrow a_1^0 a_1^0 \rightarrow 4\tau$. This decay chain is

avored in points P2 and P3 proposed here. The main emphasis is presently given to the case where all four τ -leptons decay leptonically.

A typical feature of the vector boson fusion production mode is the so-called tagging jets that are produced from the quarks that are scattered off the heavy vector bosons and merge to give the Higgs boson. These jets typically have high energies and lie in different hemispheres in the forward- and backward regions of the detector. Cutting on this signature is an important means to suppress background processes. Since there is no color flow between the quarks in the vector boson fusion process, jet production in the central detector region is suppressed. In contrast, central emission is favored in QCD interactions which constitute important background processes at the LHC [48]. Experimentally, this can be exploited by vetoing on additional jets in this region.

The decay products of the Higgs boson typically lie in the central detector region. In general, leptons from the same pseudoscalar Higgs boson form pairs that lie close to each other in the detector, the separation being sensitive to the pseudoscalar mass. The invariant mass of the lepton pair has to be lower than the pseudoscalar mass, and is thus much lower than the Z^0 -mass. For background processes including Z^0 -bosons, the photon interference therefore needs to be considered. In the experimentally most simple case, all four τ -leptons decay to muons in the process $h_1^0 \rightarrow a_1^0 a_1^0 \rightarrow 4\tau \rightarrow 4\mu + 4\nu_\mu + 4\nu_\tau$. Since muons do not deposit considerable energy in the detector, muons from a very close pair can be also classified as isolated. Decay channels including electrons need more consideration, as their energy is deposited in the calorimeters by electromagnetic showering. The possibility of separating nearby electrons or finding un-isolated muons inside an electromagnetic shower from an electron needs careful study with a full ATLAS detector simulation.

The transverse momentum of the stable leptons in this channel is rather low, since a large part of the energy is carried away by the eight neutrinos in the final state. Therefore, not in all cases will all four leptons be identified. It might therefore prove favorable to require only three leptons to be found. For triggering, two muons (electrons) with $p_T > 10$ GeV (20 GeV) or one muon (electron) with $p_T > 20$ GeV (25 GeV) are needed. It might also be considered to require a minimum transverse momentum for the remaining leptons to avoid having a high number of lepton fakes. Another feature of this channel is the large missing momentum from the eight neutrinos in the final state.

In spite of the eight neutrinos in the final state and the fact that one lepton might remain unrecognized, it is still possible to reconstruct the mass of the scalar Higgs boson with help of the collinear approximation [49] which is also used for mass reconstruction in the VBF, $Higgs \rightarrow \tau\tau$ channel [50]. Since the pseudoscalar bosons and the τ -leptons from their decays obtain a large Lorentz boost due to the large mass and p_T of the Higgs boson, their decay products are emitted roughly in the direction of the original pseudoscalar. Exploiting momentum conservation in the transverse plane yields the 4-momentum vectors of the two pseudoscalars and thus the invariant mass of the scalar Higgs boson. The performance of this algorithm for the $h_1^0 \rightarrow a_1^0 a_1^0 \rightarrow 4\tau \rightarrow 4\mu + 8\nu$ channel is currently under study at ATLAS.

Possible background processes for this channel are $t\bar{t}$ production, vector boson production in association with bottom or top quarks and production of vector boson pairs with

additional light jets. Here, the leptons can come from decays of the heavy vector bosons or from decays of the bottom quarks. The tagging jets of the vector boson fusion channel might be faked by untagged b -jets or by light jets from other sources in the event. It should be noted that the production of a vector boson pair in association with two light jets contains diagrams that have a structure similar to the vector boson fusion process, with two heavy bosons being scattered off by the incoming quarks and no color flow between the quark lines. Possible methods to separate these background processes from the signal and their performance are currently studied by the ATLAS-collaboration.

4.4 Prospects for the CMS experiment

The final state topology $h_1^0 \rightarrow a_1^0 a_1^0 \rightarrow \tau^+ \tau^- \tau^+ \tau^-$ where the ≈ 100 GeV h_1^0 state is produced in the VBF process and the pseudoscalar a_1^0 boson has a mass $2m_\tau < M_{a_1^0} < 2m_b$ (so that the $a_1^0 \rightarrow \tau^+ \tau^-$ decay mode is dominant) is currently under investigation by the CMS collaboration for the $\mu^\pm \mu^\pm \tau_{\text{jet}}^\mp \tau_{\text{jet}}^\mp$ final state containing two same sign muons and two τ jets. The τ leptons from the decays of the light a_1^0 state are approximately collinear and the non-isolated, di-muon high level trigger is needed to select the signal events. The standard CMS di-muon trigger with the relaxed isolation has the di-muon threshold of 10 GeV on both muons for a luminosity $\mathcal{L} = 2 \times 10^{33} \text{ cm}^{-2} \text{ s}^{-1}$ [52] and, since the muons from the signal events are very soft, the lower thresholds are needed. For instance, with a 7 GeV threshold the efficiency is increased approximately by a factor of two, but the QCD background rate is also increased by the same factor [53] which is not acceptable. In order to cope with the rate, the same-sign relaxed di-muon trigger was recently introduced [54]. A PYTHIA simulation shows that the rate of di-muons from $b\bar{b}$ production is reduced by a factor of four when asking for the two same sign muons to have a threshold of 5 GeV. The off-line selection strategy requires the presence of the two same sign, non-isolated muons with one track in the cone around each muon direction, thus selecting the one prong τ decays.

For a point with masses $M_{h_1^0} \sim 100$ GeV and $M_{a_1^0} \sim 5$ GeV, one applies the following basic event selection cuts at the generation level: *i*) two same sign muons with $p_T > 7$ GeV and $|\eta| < 2.1$ with one track of $p_T > 1$ GeV in a cone 0.3 around each muon; *ii*) opposite charge for the muon and the track; *iii*) two τ jets with $p_T > 10$ GeV and $|\eta| < 2.1$; *iv*) two jets with $p_T > 30$ GeV and $|\eta| < 4.5$. After these cuts, and using the SM VBF Higgs production cross section which is approximately 5.4 pb for the considered h_1^0 mass, the cross section of the signal after all selections is around 2 fb, leading to ≈ 60 events at a luminosity of 30 fb^{-1} . The dominant backgrounds with two non isolated muons from the $b\bar{b}$ and $t\bar{t}$ production processes are under evaluation.

For heavier a_1^0 bosons as in scenarios P2 and P3 where $M_{a_1^0} \sim 10$ GeV, the “non isolation” requirement should be relaxed since two τ leptons from the $a_1^0 \rightarrow \tau\tau$ decay are more separated and one should accept zero or one track in the cone (note that the relaxed di-muon trigger accepts both isolated and non isolated muons). One needs, however, to consider the backgrounds with isolated muons from W and Z^0 decays, in addition to $b\bar{b}$ and $t\bar{t}$.

The $h_1^0 \rightarrow a_1^0 a_1^0 \rightarrow \tau^+ \tau^- \tau^+ \tau^-$ decay with h_1^0 produced in Higgs-strahlung with leptonic decays of the W bosons, which can give a very clean and almost background free signal, is

also being considered by the CMS collaboration. The leptons coming from W decays allow to trigger on the events and the CMS single isolated lepton trigger can be used [53]. The signal is unique as one has two collinear τ leptons due to the large boost of a_1^0 bosons and, potentially, the most interesting final state is when one of the τ s decays hadronically and the other decays into muons.

As for the selection criteria and the muon and τ identification, for each τ -jet candidate there must be a muon in the τ -jet cone. The τ jet and the muon are required to be oppositely charged and the events with two identified τ jets with $E_T > 10$ GeV and two muons with $p_T > 7$ GeV are selected. Unlike in the VBF case, the muons are not required to have the same sign. Finally, the events with extra jets on top of two τ jets are rejected. A preliminary, full detector simulation and reconstruction analysis of the signal events in a scenario with $M_{h_1^0} \sim 100$ GeV and $M_{a_1^0} \sim 5$ GeV shows that after all selection cuts, one expects ≈ 10 events for 30 fb^{-1} data if the SM cross section of ~ 2.6 pb and a branching ratio $\text{BR}(h_1^0 \rightarrow a_1^0 a_1^0) \simeq 0.9$ are assumed. The potential backgrounds are $t\bar{t}$, tW , QCD multijet, W +jets and $b\bar{b}$ events. Preliminary tests with full detector simulation show that the requirement of two τ jets, both having an oppositely charged muon in the jet cone, as well as jet veto, suppresses the backgrounds very efficiently.

5. Conclusions

The NMSSM is a very interesting supersymmetric extension of the SM, as it solves the notorious μ problem of the MSSM and it has less fine tuning. It also leads to an interesting collider phenomenology in some cases, in particular in the Higgs sector, which is extended to contain an additional CP-even and CP-odd state as compared to the MSSM. Hence, the searches for the NMSSM Higgs bosons will be rather challenging at the LHC in scenarios in which some neutral Higgs particles are very light, opening the possibility of dominant Higgs to Higgs decays, or when all Higgs bosons are relatively light but have reduced couplings to the electroweak gauge bosons and to the top quarks, compared to the SM Higgs case. These scenarios require much more detailed phenomenological studies and experimental simulations to ensure that at least one Higgs particle of the NMSSM will be observed at the LHC.

In this note, we have proposed benchmark points in which these difficult scenarios are realized in a semi-unified NMSSM which involves a rather limited number of input parameters at the grand unification scale and which fulfills all present collider and cosmological constraints. In three of the five benchmark scenarios introduced here, the lightest CP-even Higgs boson decays mainly into two very light pseudoscalar Higgs states which subsequently decay into two b quarks or τ leptons, leading to four fermion final states; in a fourth scenario, the next-to-lightest CP-even Higgs boson has a mass of ~ 120 GeV and couplings to fermions and gauge bosons that are SM-like but it decays into pairs of the lightest Higgs state, which then decay into a $b\bar{b}$ pair; in a last scenario, all neutral and charged Higgs particles are light (with masses less than ~ 180 GeV) and have weaker couplings to W/Z^0 bosons than the SM Higgs particle. We have analysed the Higgs and supersymmetric particle spectra of these benchmark scenarios, discussed the various decay

and production rates as well as other phenomenological implications and attempted to set the basis for the search strategies to be followed by the ATLAS and CMS collaborations in order to observe at least one of the neutral Higgs states in these scenarios.

Addendum. After the completion of this paper, discussions of various possibilities of the observation of the cascade decay $h_1^0 \rightarrow a_1^0 a_1^0 \rightarrow \dots$ at hadron colliders appeared in refs. [56].

Acknowledgments

AD is grateful to the Leverhulme Trust (London, UK) for partial financial support in the form a Visiting Professorship to the NExT Institute (Southampton & RAL, UK) and to the Alexander von-Humboldt Foundation (Bonn, Germany). SM acknowledges financial support from The Royal Society (London, UK). We acknowledge support from the FP7 RTN MRTN-CT-2006-035505 (HEPTOOLS), the French ANR project PHYS@COL&COS and the Indo-French Center IFCPAR for the project number 3004-2. We also thank the organisers of the 2007 Les Houches workshop on ‘Physics at TeV Colliders’ for the nice working framework.

References

- [1] P. Fayet, *Supergauge invariant extension of the Higgs mechanism and a model for the electron and its neutrino*, *Nucl. Phys.* **B 90** (1975) 104; *Supersymmetry and weak, electromagnetic and strong interactions*, *Phys. Lett.* **B 64** (1976) 159; *Spontaneously broken supersymmetric theories of weak, electromagnetic and strong interactions*, *Phys. Lett.* **B 69** (1977) 489; H.P. Nilles, M. Srednicki and D. Wyler, *Weak interaction breakdown induced by supergravity*, *Phys. Lett.* **B 120** (1983) 346; J.M. Frere, D.R.T. Jones and S. Raby, *Fermion masses and induction of the weak scale by supergravity*, *Nucl. Phys.* **B 222** (1983) 11; J.P. Derendinger and C.A. Savoy, *Quantum effects and $SU(2) \times U(1)$ breaking in supergravity gauge theories*, *Nucl. Phys.* **B 237** (1984) 307; A.I. Veselov, M.I. Vysotsky and K.A. Ter-Martirosian, *Low-energy supergravity and the light t quark*, *Sov. Phys. JETP* **63** (1986) 489 [*Zh. Eksp. Teor. Fiz.* **90** (1986) 838]; J.R. Ellis, J.F. Gunion, H.E. Haber, L. Roszkowski and F. Zwirner, *Higgs bosons in a nonminimal supersymmetric model*, *Phys. Rev.* **D 39** (1989) 844; M. Drees, *Supersymmetric models with extended higgs sector*, *Int. J. Mod. Phys.* **A 4** (1989) 3635.
- [2] U. Ellwanger, M. Rausch de Traubenberg and C.A. Savoy, *Particle spectrum in supersymmetric models with a gauge singlet*, *Phys. Lett.* **B 315** (1993) 331 [[hep-ph/9307322](#)]; *Higgs phenomenology of the supersymmetric model with a gauge singlet*, *Z. Physik* **C 67** (1995) 665 [[hep-ph/9502206](#)]; *Phenomenology of supersymmetric models with a singlet*, *Nucl. Phys.* **B 492** (1997) 21.
- [3] U. Ellwanger, *Radiative corrections to the neutral Higgs spectrum in supersymmetry with a gauge singlet*, *Phys. Lett.* **B 303** (1993) 271 [[hep-ph/9302224](#)]; P.N. Pandita, *Radiative corrections to the scalar Higgs masses in a nonminimal supersymmetric Standard Model*, *Z. Physik* **C 59** (1993) 575;

- T. Elliott, S.F. King and P.L. White, *Radiative corrections to Higgs boson masses in the next-to-minimal supersymmetric standard model*, *Phys. Rev. D* **49** (1994) 2435 [[hep-ph/9308309](#)];
- S.F. King and P.L. White, *Resolving the constrained minimal and next-to-minimal supersymmetric standard models*, *Phys. Rev. D* **52** (1995) 4183 [[hep-ph/9505326](#)];
- F. Franke and H. Fraas, *Neutralinos and Higgs bosons in the next-to-minimal supersymmetric standard model*, *Int. J. Mod. Phys. A* **12** (1997) 479 [[hep-ph/9512366](#)];
- D.J. Miller, R. Nevzorov and P.M. Zerwas, *The Higgs sector of the next-to-minimal supersymmetric standard model*, *Nucl. Phys. B* **681** (2004) 3 [[hep-ph/0304049](#)].
- [4] J.E. Kim and H.P. Nilles, *The μ problem and the strong CP problem*, *Phys. Lett. B* **138** (1984) 150.
- [5] B.A. Dobrescu, G.L. Landsberg and K.T. Matchev, *Higgs boson decays to CP-odd scalars at the Tevatron and beyond*, *Phys. Rev. D* **63** (2001) 075003 [[hep-ph/0005308](#)];
B.A. Dobrescu and K.T. Matchev, *Light axion within the next-to-minimal supersymmetric standard model*, *JHEP* **09** (2000) 031 [[hep-ph/0008192](#)].
- [6] U. Ellwanger, J.F. Gunion, C. Hugonie and S. Moretti, *Towards a no-lose theorem for NMSSM Higgs discovery at the LHC*, [hep-ph/0305109](#); *NMSSM Higgs discovery at the LHC*, [hep-ph/0401228](#);
HIGGS WORKING GROUP collaboration, K.A. Assamagan et al., *The Higgs working group: summary report 2003*, [hep-ph/0406152](#);
LHC/LC STUDY GROUP collaboration, G. Weiglein et al., *Physics interplay of the LHC and the ILC*, *Phys. Rept.* **426** (2006) 47 [[hep-ph/0410364](#)].
- [7] U. Ellwanger, J.F. Gunion and C. Hugonie, *NMHDECAY: a Fortran code for the Higgs masses, couplings and decay widths in the NMSSM*, *JHEP* **02** (2005) 066 [[hep-ph/0406215](#)].
- [8] R. Dermisek and J.F. Gunion, *Escaping the large fine tuning and little hierarchy problems in the next to minimal supersymmetric model and $h \rightarrow aa$ decays*, *Phys. Rev. Lett.* **95** (2005) 041801 [[hep-ph/0502105](#)]; *The NMSSM solution to the fine-tuning problem, precision electroweak constraints and the largest LEP Higgs event excess*, *Phys. Rev. D* **76** (2007) 095006 [[arXiv:0705.4387](#)].
- [9] U. Ellwanger, J.F. Gunion and C. Hugonie, *Difficult scenarios for NMSSM Higgs discovery at the LHC*, *JHEP* **07** (2005) 041 [[hep-ph/0503203](#)].
- [10] R. Dermisek and J.F. Gunion, *Consistency of LEP event excesses with an $h \rightarrow aa$ decay scenario and low-fine-tuning NMSSM models*, *Phys. Rev. D* **73** (2006) 111701 [[hep-ph/0510322](#)]; *The NMSSM close to the R-symmetry limit and naturalness in $h \rightarrow aa$ decays for $m(a) < 2m(b)$* , *Phys. Rev. D* **75** (2007) 075019 [[hep-ph/0611142](#)];
S. Chang, P.J. Fox and N. Weiner, *Naturalness and Higgs decays in the MSSM with a singlet*, *JHEP* **08** (2006) 068 [[hep-ph/0511250](#)].
- [11] P.W. Graham, A. Pierce and J.G. Wacker, *Four taus at the Tevatron*, [hep-ph/0605162](#).
- [12] S. Moretti, S. Munir and P. Poulose, *Another step towards a no-lose theorem for NMSSM Higgs discovery at the LHC*, *Phys. Lett. B* **644** (2007) 241 [[hep-ph/0608233](#)];
C. Hugonie and S. Moretti, *Higgs sector of non-minimal supersymmetric models at future hadron colliders*, [hep-ph/0110241](#).
- [13] S. Chang, P.J. Fox and N. Weiner, *Visible cascade Higgs decays to four photons at hadron colliders*, *Phys. Rev. Lett.* **98** (2007) 111802 [[hep-ph/0608310](#)].

- [14] R. Dermisek and J.F. Gunion, *The NMSSM close to the R-symmetry limit and naturalness in $h \rightarrow aa$ decays for $m(a) < 2m(b)$* , *Phys. Rev. D* **75** (2007) 075019 [[hep-ph/0611142](#)].
- [15] T. Stelzer, S. Wiesenfeldt and S. Willenbrock, *Higgs at the Tevatron in extended supersymmetric models*, *Phys. Rev. D* **75** (2007) 077701 [[hep-ph/0611242](#)];
U. Aglietti et al., *Tevatron-for-LHC Report: Higgs*, [hep-ph/0612172](#).
- [16] K. Cheung, J. Song and Q.-S. Yan, *Role of $h \rightarrow \eta\eta$ in intermediate-mass Higgs boson searches at the Large Hadron Collider*, *Phys. Rev. Lett.* **99** (2007) 031801 [[hep-ph/0703149](#)].
- [17] G. Bélanger, F. Boudjema, C. Hugonie, A. Pukhov and A. Semenov, *Relic density of dark matter in the NMSSM*, *JCAP* **09** (2005) 001 [[hep-ph/0505142](#)];
J.F. Gunion, D. Hooper and B. McElrath, *Light neutralino dark matter in the NMSSM*, *Phys. Rev. D* **73** (2006) 015011 [[hep-ph/0509024](#)];
D.G. Cerdeno, E. Gabrielli, D.E. Lopez-Fogliani, C. Muñoz and A.M. Teixeira, *Phenomenological viability of neutralino dark matter in the NMSSM*, *JCAP* **06** (2007) 008 [[hep-ph/0701271](#)];
V. Barger et al., *Recoil detection of the lightest neutralino in MSSM singlet extensions*, *Phys. Rev. D* **75** (2007) 115002 [[hep-ph/0702036](#)];
C. Hugonie, G. Bélanger and A. Pukhov, *Dark matter in the constrained NMSSM*, *JCAP* **11** (2007) 009 [[arXiv:0707.0628](#)].
- [18] M. Bastero-Gil, C. Hugonie, S.F. King, D.P. Roy and S. Vempati, *Does LEP prefer the NMSSM?*, *Phys. Lett. B* **489** (2000) 359 [[hep-ph/0006198](#)].
- [19] P.C. Schuster and N. Toro, *Persistent fine-tuning in supersymmetry and the NMSSM*, [hep-ph/0512189](#).
- [20] For reviews of the SM and MSSM Higgs sectors, see A. Djouadi, *The anatomy of electro-weak symmetry breaking. I: the Higgs boson in the standard model*, *Phys. Rept.* **457** (2008) 1 [[hep-ph/0503172](#)]; *The anatomy of electro-weak symmetry breaking. II: the Higgs bosons in the minimal supersymmetric model*, *Phys. Rept.* **459** (2008) 1 [[hep-ph/0503173](#)].
- [21] J. Dai, J.F. Gunion and R. Vega, *Guaranteed detection of a minimal supersymmetric model Higgs boson at hadron supercolliders*, *Phys. Lett. B* **315** (1993) 355 [[hep-ph/9306319](#)]; *LHC detection of neutral MSSM Higgs bosons via $gg \rightarrow b\bar{b}h \rightarrow b\bar{b}b\bar{b}$* , *Phys. Lett. B* **345** (1995) 29 [[hep-ph/9403362](#)];
D. Denegri et al., *Summary of the CMS discovery potential for the MSSM SUSY Higgses*, [hep-ph/0112045](#);
M. Schumacher, *Investigation of the discovery potential for Higgs bosons of the minimal supersymmetric extension of the standard model (MSSM) with ATLAS*, [hep-ph/0410112](#).
- [22] J.F. Gunion, H.E. Haber and T. Moroi, *Will at least one of the Higgs bosons of the next-to-minimal supersymmetric extension of the standard model be observable at LEP2 or the LHC?*, [hep-ph/9610337](#);
U. Ellwanger, J.F. Gunion and C. Hugonie, *Establishing a no-lose theorem for NMSSM Higgs boson discovery at the LHC*, [hep-ph/0111179](#);
G. Azuelos et al., *The beyond the standard model working group: summary report*, [hep-ph/0204031](#).
- [23] C. Hugonie and S. Moretti, *Higgs sector of non-minimal supersymmetric models at future hadron colliders*, [hep-ph/0110241](#);
S. Moretti and S. Munir, *Di-photon Higgs signals at the LHC in the next-to-minimal supersymmetric standard model*, *Eur. Phys. J. C* **47** (2006) 791 [[hep-ph/0603085](#)].

- [24] See for instance, B.C. Allanach et al., *The Snowmass points and slopes: benchmarks for SUSY searches*, hep-ph/0202233;
 J.A. Aguilar-Saavedra et al., *Supersymmetry parameter analysis: sPA convention and project*, *Eur. Phys. J. C* **46** (2006) 43 [hep-ph/0511344].
- [25] See for instance, M.S. Carena, S. Heinemeyer, C.E.M. Wagner and G. Weiglein, *Suggestions for improved benchmark scenarios for Higgs- boson searches at LEP2*, hep-ph/9912223;
 S. Heinemeyer, *MSSM Higgs physics at higher orders*, *Int. J. Mod. Phys. A* **21** (2006) 2659 [hep-ph/0407244];
 LEP WORKING GROUP FOR HIGGS BOSON SEARCHES collaboration, R. Barate et al., *Search for the standard model Higgs boson at LEP*, *Phys. Lett. B* **565** (2003) 61 [hep-ex/0306033].
- [26] U. Ellwanger and C. Hugonie, *NMHDECAY 2.0: an updated program for sparticle masses, Higgs masses, couplings and decay widths in the NMSSM*, *Comput. Phys. Commun.* **175** (2006) 290 [hep-ph/0508022].
- [27] U. Ellwanger and C. Hugonie, *NMSPEC: a Fortran code for the sparticle and Higgs masses in the NMSSM with GUT scale boundary conditions*, *Comput. Phys. Commun.* **177** (2007) 399 [hep-ph/0612134];
 see also <http://www.th.u-psud.fr/NMHDECAY/nmssmtools.html>.
- [28] M. Cvetič, D.A. Demir, J.R. Espinosa, L.L. Everett and P. Langacker, *Electroweak breaking and the mu problem in supergravity models with an additional U(1)*, *Phys. Rev. D* **56** (1997) 2861 [*Erratum ibid.* **58** (1998) 119905] [hep-ph/9703317];
 P. Langacker and J. Wang, *U(1)' symmetry breaking in supersymmetric E_6 models*, *Phys. Rev. D* **58** (1998) 115010 [hep-ph/9804428];
 C. Panagiotakopoulos and K. Tamvakis, *Stabilized NMSSM without domain walls*, *Phys. Lett. B* **446** (1999) 224 [hep-ph/9809475]; *New minimal extension of MSSM*, *Phys. Lett. B* **469** (1999) 145 [hep-ph/9908351];
 C. Panagiotakopoulos and A. Pilaftsis, *Higgs scalars in the minimal non-minimal supersymmetric standard model*, *Phys. Rev. D* **63** (2001) 055003 [hep-ph/0008268];
 A. Dedes, C. Hugonie, S. Moretti and K. Tamvakis, *Phenomenology of a new minimal supersymmetric extension of the standard model*, *Phys. Rev. D* **63** (2001) 055009 [hep-ph/0009125];
 A. Menon, D.E. Morrissey and C.E.M. Wagner, *Electroweak baryogenesis and dark matter in the NMSSM*, *Phys. Rev. D* **70** (2004) 035005 [hep-ph/0404184];
 T. Han, P. Langacker and B. McElrath, *The Higgs sector in a U(1)' extension of the MSSM*, *Phys. Rev. D* **70** (2004) 115006 [hep-ph/0405244];
 D.A. Demir, G.L. Kane and T.T. Wang, *The minimal U(1)' extension of the MSSM*, *Phys. Rev. D* **72** (2005) 015012 [hep-ph/0503290];
 S.F. King, S. Moretti and R. Nevzorov, *Exceptional supersymmetric standard model*, *Phys. Lett. B* **634** (2006) 278 [hep-ph/0511256]; *Theory and phenomenology of an exceptional supersymmetric standard model*, *Phys. Rev. D* **73** (2006) 035009 [hep-ph/0510419];
 V. Barger, P. Langacker, H.-S. Lee and G. Shaughnessy, *Higgs sector in extensions of the MSSM*, *Phys. Rev. D* **73** (2006) 115010 [hep-ph/0603247].
- [29] M. Matsuda and M. Tanimoto, *Explicit CP-violation of the Higgs sector in the next-to-minimal supersymmetric standard model*, *Phys. Rev. D* **52** (1995) 3100 [hep-ph/9504260];
 N. Haba, *Explicit CP-violation in the Higgs sector of the next-to-minimal supersymmetric standard model*, *Prog. Theor. Phys.* **97** (1997) 301 [hep-ph/9608357];

- S.W. Ham, S.K. Oh and D. Son, *Neutral Higgs sector of the next-to-minimal supersymmetric standard model with explicit CP-violation*, *Phys. Rev. D* **65** (2002) 075004 [[hep-ph/0110052](#)];
M. Boz, *The Higgs sector and electron electric dipole moment in next-to-minimal supersymmetry with explicit CP-violation*, *Mod. Phys. Lett. A* **21** (2006) 243 [[hep-ph/0511072](#)];
S.W. Ham, S.H. Kim, S.K. OH and D. Son, *Higgs bosons of the NMSSM with explicit CP-violation at the ILC*, *Phys. Rev. D* **76** (2007) 115013 [[arXiv:0708.2755](#)].
- [30] U. Ellwanger and C. Hugonie, *Neutralino cascades in the $(M + 1)$ SSM*, *Eur. Phys. J. C* **5** (1998) 723 [[hep-ph/9712300](#)]; *Topologies of the $(M + 1)$ SSM with a singlino LSP at LEP2*, *Eur. Phys. J. C* **13** (2000) 681 [[hep-ph/9812427](#)];
V. Barger, P. Langacker and G. Shaughnessy, *Neutralino signatures of the singlet extended MSSM*, *Phys. Lett. B* **644** (2007) 361 [[hep-ph/0609068](#)]; *Collider signatures of singlet extended Higgs sectors*, *Phys. Rev. D* **75** (2007) 055013 [[hep-ph/0611239](#)].
- [31] J.A. Casas, J.R. Espinosa and I. Navarro, *Unconventional low-energy SUSY from warped geometry*, *Nucl. Phys. B* **620** (2002) 195 [[hep-ph/0109127](#)];
A. Birkedal, Z. Chacko and Y. Nomura, *Relaxing the upper bound on the mass of the lightest supersymmetric Higgs boson*, *Phys. Rev. D* **71** (2005) 015006 [[hep-ph/0408329](#)].
- [32] For recent analyses, see for instance, J.R. Ellis, S. Heinemeyer, K.A. Olive and G. Weiglein, *Phenomenological indications of the scale of supersymmetry*, *JHEP* **05** (2006) 005 [[hep-ph/0602220](#)];
J.R. Ellis, S. Heinemeyer, K.A. Olive and G. Weiglein, *Light heavy MSSM Higgs bosons at large $\tan\beta$* , *Phys. Lett. B* **653** (2007) 292 [[arXiv:0706.0977](#)].
- [33] A. Djouadi, J. Kalinowski, P. Ohmann and P.M. Zerwas, *Heavy SUSY Higgs bosons at e^+e^- linear colliders*, *Z. Physik C* **74** (1997) 93 [[hep-ph/9605339](#)].
- [34] A. Djouadi, M. Spira and P.M. Zerwas, *Production of Higgs bosons in proton colliders: QCD corrections*, *Phys. Lett. B* **264** (1991) 440; *QCD Corrections to Hadronic Higgs Decays*, *Z. Physik C* **70** (1996) 427 [[hep-ph/9511344](#)];
M. Spira, A. Djouadi, D. Graudenz and P.M. Zerwas, *Higgs boson production at the LHC*, *Nucl. Phys. B* **453** (1995) 17 [[hep-ph/9504378](#)];
A. Djouadi, J. Kalinowski and M. Spira, *HDECAY: a program for Higgs boson decays in the standard model and its supersymmetric extension*, *Comput. Phys. Commun.* **108** (1998) 56 [[hep-ph/9704448](#)].
- [35] J.F. Gunion and H.E. Haber, *Higgs bosons in supersymmetric models. 1*, *Nucl. Phys. B* **272** (1986) 1 [*Erratum ibid.* **402** (1993) 567];
P. Kalyniak, R. Bates and J.N. Ng, *Two photon decays of scalar and pseudoscalar bosons in supersymmetry*, *Phys. Rev. D* **33** (1986) 755;
R. Bates, J.N. Ng and P. Kalyniak, *Two photon decay widths of Higgs bosons in minimal broken supersymmetry*, *Phys. Rev. D* **34** (1986) 172;
J.F. Gunion, G. Gamberini and S.F. Novaes, *Can the Higgs bosons of the minimal supersymmetric model be detected at a hadron collider via two photon decays?*, *Phys. Rev. D* **38** (1988) 3481;
A. Djouadi, V. Driesen, W. Hollik and J.I. Illana, *The coupling of the lightest SUSY Higgs boson to two photons in the decoupling regime*, *Eur. Phys. J. C* **1** (1998) 149 [[hep-ph/9612362](#)];
A. Djouadi, *Squark effects on Higgs boson production and decay at the LHC*, *Phys. Lett. B* **435** (1998) 101 [[hep-ph/9806315](#)].

- [36] ALEPH collaboration, S. Schael et al., *Search for neutral MSSM Higgs bosons at LEP*, *Eur. Phys. J. C* **47** (2006) 547 [[hep-ex/0602042](#)].
- [37] G. Hiller, *B-physics signals of the lightest CP-odd Higgs in the NMSSM at large $\tan\beta$* , *Phys. Rev. D* **70** (2004) 034018 [[hep-ph/0404220](#)];
F. Domingo and U. Ellwanger, *Updated constraints from B physics on the MSSM and the NMSSM*, *JHEP* **12** (2007) 090 [[arXiv:0710.3714](#)].
- [38] G. Bélanger, F. Boudjema, A. Pukhov and A. Semenov, *MicrOMEGAs: a program for calculating the relic density in the MSSM*, *Comput. Phys. Commun.* **149** (2002) 103 [[hep-ph/0112278](#)]; *MicrOMEGAs: version 1.3*, *Comput. Phys. Commun.* **174** (2006) 577 [[hep-ph/0405253](#)].
- [39] J. Hamann, S. Hannestad, M.S. Sloth and Y. Wong, *How robust are inflation model and dark matter constraints from cosmological data?*, *Phys. Rev. D* **75** (2007) 023522 [[astro-ph/0611582](#)].
- [40] E. Boos, A. Djouadi, M. Muhlleitner and A. Vologdin, *The MSSM Higgs bosons in the intense-coupling regime*, *Phys. Rev. D* **66** (2002) 055004 [[hep-ph/0205160](#)];
E. Boos, V. Bunichev, A. Djouadi and H.J. Schreiber, *Prospects of mass measurements for neutral MSSM Higgs bosons in the intense-coupling regime at a linear collider*, *Phys. Lett. B* **622** (2005) 311 [[hep-ph/0412194](#)];
A. Djouadi and Y. Mambrini, *The Higgs intense-coupling regime in constrained SUSY models and its astrophysical implications*, *JHEP* **12** (2006) 001 [[hep-ph/0609234](#)].
- [41] E. Boos, A. Djouadi and A. Nikitenko, *Detection of the neutral MSSM Higgs bosons in the intense-coupling regime at the LHC*, *Phys. Lett. B* **578** (2004) 384 [[hep-ph/0307079](#)].
- [42] U. Ellwanger and C. Hugonie, *The upper bound on the lightest Higgs mass in the NMSSM revisited*, *Mod. Phys. Lett. A* **22** (2007) 1581 [[hep-ph/0612133](#)].
- [43] S. Moretti, K. Odagiri, P. Richardson, M.H. Seymour and B.R. Webber, *Implementation of supersymmetric processes in the HERWIG event generator*, *JHEP* **04** (2002) 028 [[hep-ph/0204123](#)];
G. Corcella et al., *HERWIG 6: an event generator for hadron emission reactions with interfering gluons (including supersymmetric processes)*, *JHEP* **01** (2001) 010 [[hep-ph/0011363](#)].
- [44] T. Sjöstrand, L. Lönnblad and S. Mrenna, *PYTHIA 6.2: physics and manual*, [hep-ph/0108264](#).
- [45] S. Baffioni, *$h \rightarrow AA$ in NMSSM*, talk presented at *GdR Supersymétrie 2004*, 5–7 July 2004, Clermont-Ferrand, France;
S. Baffioni and D. Zerwas, private communication.
- [46] D.J. Miller and S. Moretti, *An interesting NMSSM scenario at the LHC and LC*, [hep-ph/0403137](#).
- [47] ATLAS collaboration, *ATLAS detector and physics performance. Technical design report*, CERN-LHCC-99-014; *ATLAS detector and physics performance. Technical design report 2*, CERN-LHCC-99-015.
- [48] V.D. Barger, R.J.N. Phillips and D. Zeppenfeld, *Mini-jet veto: a tool for the heavy Higgs search at the LHC*, *Phys. Lett. B* **346** (1995) 106 [[hep-ph/9412276](#)];
D.L. Rainwater and D. Zeppenfeld, *Searching for $H \rightarrow \gamma\gamma$ in weak boson fusion at the LHC*, *JHEP* **12** (1997) 005 [[hep-ph/9712271](#)].

- [49] R.K. Ellis, I. Hinchliffe, M. Soldate and J.J. van der Bij, *Higgs decay to $\tau^+\tau^-$: a possible signature of intermediate mass Higgs bosons at the SSC*, *Nucl. Phys. B* **297** (1988) 221.
- [50] S. Asai et al., *Prospects for the search for a standard model Higgs boson in ATLAS using vector boson fusion* *Eur. Phys. J. C* **35S2** (2004) 19 [[hep-ph/0402254](#)].
- [51] I. Rottländer et al., to appear.
- [52] CMS collaboration, *CMS physics : technical design report 2*, CERN-LHCC-2006-021, CMS-TDR-008-2.
- [53] CMS collaboration, *The CMS trigger and data acquisition project, volume II*, CERN-LHCC-2002-26.
- [54] CMS collaboration, *The CMS high level trigger*, CERN-LHCC-2007-021, LHCC-G-134.
- [55] S. Lehti et al., to appear.
- [56] M. Carena, T. Han, G.-Y. Huang and C.E.M. Wagner, *Higgs signal for $h \rightarrow aa$ at hadron colliders*, *JHEP* **04** (2008) 092 [[arXiv:0712.2466](#)];
J.R. Forshaw, J.F. Gunion, L. Hodgkinson, A. Papaefstathiou and A.D. Pilkington, *Reinstating the 'no-lose' theorem for NMSSM Higgs discovery at the LHC*, *JHEP* **04** (2008) 090 [[arXiv:0712.3510](#)];
R. Barbieri, L.J. Hall, A.Y. Papaioannou, D. Pappadopulo and V.S. Rychkov, *An alternative NMSSM phenomenology with manifest perturbative unification*, *JHEP* **03** (2008) 005 [[arXiv:0712.2903](#)].